

COMPOSITION OF ANALYTIC PARAPRODUCTS

ALEXANDRU ALEMAN, CARME CASCANTE, JOAN FÀBREGA,
DANIEL PASCUAS, AND JOSÉ ÁNGEL PELÁEZ

ABSTRACT. For a fixed analytic function g on the unit disc \mathbb{D} , we consider the analytic paraproducts induced by g , which are defined by $T_g f(z) = \int_0^z f(\zeta)g'(\zeta) d\zeta$, $S_g f(z) = \int_0^z f'(\zeta)g(\zeta) d\zeta$, and $M_g f(z) = f(z)g(z)$. The boundedness of these operators on various spaces of analytic functions on \mathbb{D} is well understood. The original motivation for this work is to understand the boundedness of compositions of two of these operators, for example T_g^2 , $T_g S_g$, $M_g T_g$, etc. Our methods yield a characterization of the boundedness of a large class of operators contained in the algebra generated by these analytic paraproducts acting on the classical weighted Bergman and Hardy spaces in terms of the symbol g . In some cases it turns out that this property is not affected by cancellation, while in others it requires stronger and more subtle restrictions on the oscillation of the symbol g than the case of a single paraproduct.

1. INTRODUCTION

Let $\mathcal{H}(\mathbb{D})$ be the space of analytic functions on the unit disc \mathbb{D} of the complex plane. For $\alpha > -1$ and $0 < p < \infty$, the weighted Bergman space A_α^p consists of the functions $f \in \mathcal{H}(\mathbb{D})$ such that

$$\|f\|_{\alpha,p}^p := (\alpha + 1) \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\alpha dA(z) < \infty,$$

where dA is the the normalized area measure on \mathbb{D} . Let H^p , $0 < p \leq \infty$, denote the classical Hardy space of analytic functions in \mathbb{D} . To simplify the notations, we shall write $A_{-1}^p := H^p$ and $\|\cdot\|_{-1,p} := \|\cdot\|_{H^p}$, $0 < p < \infty$. Given $g \in \mathcal{H}(\mathbb{D})$, let us consider the multiplication operator $M_g f = fg$ and the integral operators

$$T_g f(z) = \int_0^z f(\zeta)g'(\zeta) d\zeta \quad S_g f(z) = \int_0^z f'(\zeta)g(\zeta) d\zeta \quad (z \in \mathbb{D}).$$

Due to the integration by parts relation

$$(1.1) \quad M_g f = T_g f + S_g f + g(0)f(0),$$

we call these operators *analytic paraproducts*.

Date: October 16, 2021.

2020 Mathematics Subject Classification. 30H10, 30H20, 47G10.

Key words and phrases. Analytic paraproduct, Hardy spaces, weighted Bergman spaces, Bloch space, BMOA space.

The research of the second, third and fourth author was supported in part by Ministerio de Economía y Competitividad, Spain, project MTM2017-83499-P, and Generalitat de Catalunya, project 2017SGR358. The research of the fifth author was supported in part by Ministerio de Economía y Competitividad, Spain, projects PGC2018-096166-B-100; La Junta de Andalucía, projects FQM210 and UMA18-FEDERJA-002.

It is well-known [2, 3, 4, 10] that T_g is bounded on A_α^p if and only if g belongs to the Bloch space \mathcal{B} when $\alpha > -1$, and $g \in BMOA$ in the Hardy space case $\alpha = -1$. In particular, these results show that cancellation plays a key role in the boundedness of the integral operator T_g . This is very different from the case of M_g and S_g , whose boundedness on these spaces is equivalent to the boundedness of g in \mathbb{D} (see Proposition 2.4 below and the references following it).

Throughout the paper the spaces of bounded and compact linear operators on A_α^p are denoted by $\mathcal{B}(A_\alpha^p)$ and $\mathcal{K}(A_\alpha^p)$, respectively. Moreover, if $T : A_\alpha^p \rightarrow A_\alpha^p$ is a linear map, we write $\|T\|_{\alpha,p} = \sup_{\|f\|_{\alpha,p} \leq 1} \|Tf\|_{\alpha,p}$, and we refer to this quantity as the operator norm of T on A_α^p , despite that A_α^p is not a normed space for $0 < p < 1$.

The primary aim of this paper is to study the boundedness of compositions (products) of analytic paraproducts acting on A_α^p . In order to provide some intuition and motivation for this circle of problems, let us have a brief look at compositions of two such paraproducts. Clearly, $M_g^2 = M_g M_g$ is bounded on these spaces if and only if $g \in H^\infty$ and we shall show (Theorem 1.2) that the same holds for S_g^2 . On the other hand, it turns out that $T_g^2 \in \mathcal{B}(A_\alpha^p)$ if and only if $T_g \in \mathcal{B}(A_\alpha^p)$ (Theorem 1.1). Regarding mixed products, a simple computation reveals that $S_g T_g = T_g M_g = \frac{1}{2} T_g^2$, so that both compositions are bounded on A_α^p if and only if $g^2 \in \mathcal{B}$, when $\alpha > -1$, or $g^2 \in BMOA$, when $\alpha = -1$. This last condition is strictly stronger than $g \in \mathcal{B}$ or $g \in BMOA$, respectively (see Proposition 2.1 below). The compositions in reversed order raise additional problems because they cannot be expressed as a single paraproduct. They can be related to the previous ones using (1.1):

$$(1.2) \quad M_g T_g = S_g T_g + T_g^2, \quad T_g S_g = S_g T_g - T_g^2 - g(0)(g - g(0))\delta_0,$$

where $\delta_0 f = f(0)$. Intuitively, from above it appears that $S_g T_g = \frac{1}{2} T_g^2$ is the “dominant term” in both sums, but a priori it is not clear whether such sums are affected by cancellation or not. Thus we are led in a natural way to consider sums of compositions of paraproducts rather than only compositions. A similar situation occurs when dealing with $M_g S_g$ and $S_g M_g$. Due to these preliminary observations we turn our attention to the full algebra \mathcal{A}_g generated by the operators M_g , S_g , and T_g . The operators in \mathcal{A}_g will be called *g-operators*. In this general framework we begin by showing that any *g-operator* L has a representation

$$(1.3) \quad L = \sum_{k=0}^n S_g^k T_g P_k(T_g) + S_g P_{n+1}(S_g) + g(0) P_{n+2}(g - g(0)) \delta_0,$$

for some $n \in \mathbb{N}$, where the P_k 's are polynomials. This representation is essentially unique, see §3.2. If $P_k = 0$, for $0 \leq k \leq n+1$, we will say that L is a *trivial operator*. With this representation in hand we can derive a fairly surprising necessary condition for the boundedness of general operators in this algebra.

Theorem 1.1. *Let $g \in \mathcal{H}(\mathbb{D})$, $0 < p < \infty$ and $\alpha \geq -1$. Let L be a *g-operator* written in the form (1.3). Then:*

- a) *If L is a non-trivial operator and $L \in \mathcal{B}(A_\alpha^p)$, then $T_g \in \mathcal{B}(A_\alpha^p)$, that is, $g \in \mathcal{B}$, when $\alpha > -1$, and $g \in BMOA$, when $\alpha = -1$.*

- b) If L is a non-zero trivial operator, then $L \in \mathcal{B}(A_\alpha^p)$ if and only if $g^{\deg P_{n+2}} \in A_\alpha^p$.

Note that the result applies directly to $T_g S_g$ and $M_g T_g$ via (1.2) and justifies the intuition that these operators are bounded on A_α^p if and only if $g^2 \in \mathcal{B}$, when $\alpha > -1$, or $g^2 \in BMOA$, when $\alpha = -1$. In fact, in §4.3 we provide a complete characterization of the boundedness of compositions of two analytic paraproducts (see also Corollary 1.4 below). The theorem can be used to characterize the boundedness of more complicated g -operators. In addition, it provides a crucial preliminary step in the proof of our characterization of boundedness of certain g -operators which we now state.

Theorem 1.2. *Let $g \in \mathcal{H}(\mathbb{D})$, $0 < p < \infty$ and $\alpha \geq -1$. Let L be a g -operator written in the form (1.3). Then:*

- a) If $P_{n+1} \neq 0$, $L \in \mathcal{B}(A_\alpha^p)$ if and only if $g \in H^\infty$.
b) If $P_{n+1} = 0$ and $P_n = 1$, $L \in \mathcal{B}(A_\alpha^p)$ if and only if $T_{g^{n+1}} \in \mathcal{B}(A_\alpha^p)$, or equivalently, $g^{n+1} \in \mathcal{B}$, when $\alpha > -1$, and $g^{n+1} \in BMOA$, when $\alpha = -1$.
c) If $\alpha > -1$, $P_{n+1} = 0$, and $P_n(0) \neq 0$, $L \in \mathcal{B}(A_\alpha^p)$ if and only if $g^{n+1} \in \mathcal{B}$.

We have not been able to extend part c) of this theorem to the H^p -case. One direction follows directly from Proposition 2.1, but the remaining one is, in our opinion, an interesting and challenging open question.

Question 1.3. *Let $g \in \mathcal{H}(\mathbb{D})$, $0 < p < \infty$, and let L be a g -operator written in the form (1.3) with $P_{n+1} = 0$, and $P_n(0) \neq 0$, which is bounded on H^p . Is it true that $g^{n+1} \in BMOA$?*

When dealing with operators in \mathcal{A}_g , an initial hurdle can be easily recognized, namely that these operators are formally defined as sums of products of possibly unbounded operators on the spaces in question. One way to overcome this difficulty is to consider dilations of the symbol g , which are defined, for $\lambda \in \mathbb{D}$, by $g_\lambda(z) = g(\lambda z)$. In Proposition 4.3 we prove that if $L_g \in \mathcal{A}_g \cap \mathcal{B}(A_\alpha^p)$ then $\|L_{g_\lambda}\|_{\alpha,p} \leq \|L_g\|_{\alpha,p}$ for all $\lambda \in \mathbb{D}$. This fact will be repeatedly used in the proofs of the results stated above. Other key ingredients for the proof of Theorem 1.1 are the estimates

$$\|T_g\|_{\alpha,p}^n \leq c_n \|T_g^n\|_{\alpha,p},$$

which will be established in Proposition 4.1, together with the analysis of iterated commutators of T_g and S_g^k , $k \in \mathbb{N}$. A sample of this set of ideas can be found in Corollary 4.9 below. The proof of Theorem 1.2 is somewhat more involved, in particular, it makes use of the boundary behaviour of A_α^p -valued functions of the form $\lambda \rightarrow L_{g_\lambda} f$, $\lambda \in \mathbb{D}$, $f \in A_\alpha^p$.

In order to discuss the class of g -operators covered by Theorem 1.2 b) it is convenient to introduce the following terminology. An n -letter g -word is a g -operator of the form $L = L_1 \cdots L_n$, where each L_j is either M_g , S_g or T_g . For $n \in \mathbb{N}$, let $\mathcal{A}_g^{(n)}$ be the linear span of g -words with no more than n letters and define the *order* of a g -operator L to be the least $n \in \mathbb{N}$ such that $L \in \mathcal{A}_g^{(n)}$. It turns out that if $L \in \mathcal{A}_g^{(n)}$ then the words involved in

its representation (1.3) have length at most n . For example, g -operators of order two have the form

$$(1.4) \quad L = a_1 T_g + a_2 T_g^2 + a_3 S_g T_g + a_4 S_g + a_5 S_g^2 + g(0)P(g - g(0))\delta_0,$$

where $a_j \in \mathbb{C}$, $1 \leq j \leq 5$, and P is a polynomial of degree smaller than 2. These operators are covered by Theorem 1.2, and we have the following complete characterization of their boundedness.

Corollary 1.4. *Let $g \in \mathcal{H}(\mathbb{D})$, $0 < p < \infty$ and $\alpha \geq -1$. If L is a g -operator of order two written in the form (1.4), then:*

- a) *When either $a_4 \neq 0$ or $a_5 \neq 0$, $L \in \mathcal{B}(A_\alpha^p)$ if and only if $g \in H^\infty$.*
- b) *When $a_3 \neq 0$ and $a_4 = a_5 = 0$, $L \in \mathcal{B}(A_\alpha^p)$ if and only if $g^2 \in \mathcal{B}$, for $\alpha > -1$, and $g^2 \in BMOA$, for $\alpha = -1$.*
- c) *When $a_3 = a_4 = a_5 = 0$ and either $a_1 \neq 0$ or $a_2 \neq 0$, $L \in \mathcal{B}(A_\alpha^p)$ if and only if $g \in \mathcal{B}$, for $\alpha > -1$, and $g \in BMOA$, for $\alpha = -1$.*

On the other hand, our main result does not cover g -operators with $P_{n+1} = 0$ and $P_n(0) = 0$ in the representation (1.3). An example of this type, where the condition for the boundedness is different, follows from the second identity in (1.2). This together with $S_g T_g = \frac{1}{2} T_{g^2}$ implies

$$\frac{1}{4} T_{g^2}^2 = S_g(T_g S_g) T_g = S_g^2 T_g^2 - S_g T_g^3,$$

i.e. the operator on the right is the representation (1.3) of $\frac{1}{4} T_{g^2}^2$. In view of Theorem 1.2 one might expect that the presence of S_g^2 forces the boundedness of T_{g^3} , but by Theorem 1.1 this operator is bounded on A_α^p if and only if $g^2 \in \mathcal{B}$, for $\alpha > -1$, and $g^2 \in BMOA$, for $\alpha = -1$.

There are also g -operators of order 3 with $P_{n+1} = 0$ and $P_n(0) = 0$ in the representation (1.3). The simplest example is the 3-letter-word $S_g T_g^2$ and in this case the situation differs even more dramatically to the one described in Theorem 1.2. The following result shows that the boundedness of such g -operators cannot be characterized with conditions of the form $g \in H^\infty$, or $g^n \in \mathcal{B}$ ($BMOA$), with $n \in \mathbb{N}$.

As usual, we denote by \log the principal branch of the logarithm on $\mathbb{C} \setminus (-\infty, 0]$, that is, $\log 1 = 0$. For an open set $U \subset \mathbb{C}$ and an analytic function $h : U \rightarrow \mathbb{C} \setminus (-\infty, 0]$, $\beta \in \mathbb{C}$, we define $h^\beta = \exp(\beta \log h)$.

Theorem 1.5. *Consider the function $g : \mathbb{D} \rightarrow \mathbb{C} \setminus (-\infty, 0]$ defined by*

$$g(z) = \log\left(\frac{e}{1-z}\right) \quad (z \in \mathbb{D}).$$

Then:

- a) *$g \in BMOA$, but for any $\alpha \geq -1$, $p > 0$, we have $S_g T_g^2 \notin \mathcal{B}(A_\alpha^p)$.*
- b) *For $\frac{1}{2} < \beta < \frac{2}{3}$, $g^{2\beta} \notin \mathcal{B}$ (and so $g^{2\beta} \notin BMOA$), but $S_{g^\beta} T_{g^\beta}^2 \in \mathcal{K}(A_\alpha^p)$, for any $\alpha \geq -1$ and $p > 0$.*

The paper is organized as follows. Section 2 contains some preliminary results concerning the Bloch space and $BMOA$, in particular the condition $g^k \in \mathcal{B}(BMOA)$, for some $k \in \mathbb{N}$. In Section 3 we study the vector space structure of the algebra \mathcal{A}_g and prove the representation (1.3). Section 4 is

devoted to the proof of our main results, Theorems 1.1 and 1.2. Finally, in the last section we prove Theorem 1.5.

As usual, \mathbb{N} is the set of positive integers and $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ is the unit circle. For $\lambda \in \mathbb{C}$ and $r > 0$, $D(\lambda, r) = \{z \in \mathbb{C} : |z - \lambda| < r\}$ is the open disc centered at λ with radius r . For two non-negative functions A and B , $A \lesssim B$ ($B \gtrsim A$) means that there is a finite positive constant C , independent of the variables involved, which satisfies $A \leq CB$. Moreover, we will write $A \simeq B$ when $A \lesssim B$ and $B \lesssim A$.

2. THE SPACES OF SYMBOLS

In this section we will recall and prove some preliminary results about $BMOA$ and the Bloch space. For any $a \in \mathbb{D}$, define $\phi_a(z) := \frac{a-z}{1-\bar{a}z}$, and consider the classical $BMOA$ and Bloch spaces endowed with their Garsia's seminorms $\|\cdot\|_{BMOA}$ and $\|\cdot\|_{\mathcal{B}}$ (see, for instance, [6, 8] and the references therein):

$$\begin{aligned} BMOA &:= \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{BMOA}^2 := \sup_{a \in \mathbb{D}} \|f \circ \phi_a - f(a)\|_{H^2}^2 < \infty \right\} \\ \mathcal{B} &:= \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{\mathcal{B}}^2 := \sup_{a \in \mathbb{D}} \|f \circ \phi_a - f(a)\|_{A^2}^2 < \infty \right\}. \end{aligned}$$

For a given Banach space (or a complete metric space) X of analytic functions on \mathbb{D} , a positive Borel measure μ on \mathbb{D} is called a q -Carleson measure for X (vanishing q -Carleson measure for X) if the identity operator $I : X \rightarrow L^q(\mu)$ is bounded (compact). Recall that $f \in \mathcal{B}$ if and only if $\|f\|_{\mathcal{B}} := \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty$, and $f \in BMOA$ if and only if $(1 - |z|^2) |f'(z)|^2 dA(z)$ is a Carleson measure for H^p , $0 < p < \infty$, or equivalently

$$\|f\|_{BMOA}^2 := \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} (1 - |\phi_a|^2) |f'|^2 dA < \infty.$$

Moreover, $\|f\|_{\mathcal{B}} \simeq \|f\|_{\mathcal{B}}$ and $\|f\|_{BMOA} \simeq \|f\|_{BMOA}$.

We also consider the little-oh subspaces of $BMOA$ and \mathcal{B} :

$$\begin{aligned} VMOA &:= \left\{ f \in H^2 : \lim_{|a| \rightarrow 1^-} \|f \circ \phi_a - f(a)\|_{H^2}^2 = 0 \right\} \\ \mathcal{B}_0 &:= \left\{ f \in A^2 : \lim_{|a| \rightarrow 1^-} \|f \circ \phi_a - f(a)\|_{A^2}^2 = 0 \right\}. \end{aligned}$$

For $f \in \mathcal{H}(\mathbb{D})$, recall that $f \in \mathcal{B}_0$ if and only if $\lim_{|z| \rightarrow 1^-} (1 - |z|^2) |f'(z)| = 0$, and $f \in VMOA$ if and only if $(1 - |z|^2) |f'(z)|^2 dA(z)$ is a vanishing Carleson measure for H^p , $0 < p < \infty$, or equivalently

$$\lim_{|a| \rightarrow 1^-} \int_{\mathbb{D}} (1 - |\phi_a|^2) |f'|^2 dA = 0.$$

For $0 < p < \infty$ and $m, n \in \mathbb{N}$, $m \leq n$, Jensen's inequality shows that $\|f^m\|_{\alpha, p}^{1/m} \leq \|f^n\|_{\alpha, p}^{1/n}$. We will show that this result also holds for the Garsia's $BMOA$ and Bloch seminorms.

Proposition 2.1. *Let $m, n \in \mathbb{N}$, $m < n$, and $f \in \mathcal{H}(\mathbb{D})$. Then,*

$$(2.1) \quad \|f^m\|_{BMOA}^{1/m} \leq \|f^n\|_{BMOA}^{1/n}$$

$$(2.2) \quad \|f^m\|_{\mathcal{B}}^{1/m} \leq \|f^n\|_{\mathcal{B}}^{1/n}.$$

In particular, if $f^n \in BMOA$ ($f^n \in \mathcal{B}$), then $f^m \in BMOA$ ($f^m \in \mathcal{B}$). Moreover, if $f^n \in VMOA$ ($f^n \in \mathcal{B}_0$), then $f^m \in VMOA$ ($f^m \in \mathcal{B}_0$).

Bearing in mind that $f \mapsto f \circ \phi_a$ maps H^2 or A^2 to itself, Proposition 2.1 follows from the following lemma.

Lemma 2.2. *Let $m, n \in \mathbb{N}$, $m < n$. Then:*

$$(2.3) \quad \|f^m - f^m(0)\|_{H^2}^{1/m} \leq \|f^n - f^n(0)\|_{H^2}^{1/n} \quad (f \in \mathcal{H}(\mathbb{D}))$$

$$(2.4) \quad \|f^m - f^m(0)\|_{A^2}^{1/m} \leq \|f^n - f^n(0)\|_{A^2}^{1/n} \quad (f \in \mathcal{H}(\mathbb{D})).$$

Proof. We only prove (2.3), the proof of (2.4) is completely analogous replacing H^2 by A^2 . First of all, recall that

$$(2.5) \quad \|f^k\|_{H^2}^2 = \|f\|_{H^{2k}}^{2k} \quad (f \in \mathcal{H}(\mathbb{D}), k \in \mathbb{N}),$$

and, by Jensen's inequality,

$$(2.6) \quad \|f\|_{H^{2n}} \geq \|f\|_{H^{2m}} \quad (f \in \mathcal{H}(\mathbb{D})).$$

Now (2.3), in the case $f(0) = 0$, directly follows from (2.5) and (2.6). Indeed, we have that

$$\|f^n\|_{H^2}^2 = \|f\|_{H^{2n}}^{2n} \geq \|f\|_{H^{2m}}^{2n} = \|f^m\|_{H^2}^{(2n)/m} \quad (f \in \mathcal{H}(\mathbb{D})),$$

and so

$$(2.7) \quad \|f^n\|_{H^2}^{1/n} \geq \|f^m\|_{H^2}^{1/m} \quad (f \in \mathcal{H}(\mathbb{D})),$$

which, in particular, gives (2.3) when $f(0) = 0$.

The general case is a consequence of (2.6), (2.7), and a simple argument. First note that

$$(2.8) \quad \|f^k - f^k(0)\|_{H^2}^2 = \|f^k\|_{H^2}^2 - |f(0)|^{2k} \quad (f \in \mathcal{H}(\mathbb{D}), k \in \mathbb{N}).$$

Then, for any $f \in \mathcal{H}(\mathbb{D})$, we have that

$$\begin{aligned} \|f^n - f^n(0)\|_{H^2}^2 &\stackrel{(*)}{=} \|f^n\|_{H^2}^2 - |f(0)|^{2n} \\ &\stackrel{(\star)}{\geq} \|f^m\|_{H^2}^{(2n)/m} - |f(0)|^{2n} \\ &\stackrel{(*)}{=} (\|f^m - f^m(0)\|_{H^2}^2 + |f(0)|^{2m})^{n/m} - |f(0)|^{2n} \\ &\stackrel{(\diamond)}{\geq} \|f^m - f^m(0)\|_{H^2}^{(2n)/m}, \end{aligned}$$

where $(*)$ and (\star) follow from (2.8) and (2.7), respectively, while (\diamond) is a consequence of the classical superadditivity inequality

$$(x + y)^\alpha \geq x^\alpha + y^\alpha \quad (x, y \geq 0, \alpha \geq 1).$$

(Recall that any convex function $\varphi : [0, \infty) \rightarrow \mathbb{R}$ with $\varphi(0) = 0$ is superadditive, i.e. $\varphi(x + y) \geq \varphi(x) + \varphi(y)$, for any $x, y \geq 0$.)

Hence

$$\|f^n - f^n(0)\|_{H^2}^{1/n} \geq \|f^m - f^m(0)\|_{H^2}^{1/m} \quad (f \in \mathcal{H}(\mathbb{D})),$$

and the proof is complete. \square

The final part of this section recalls the descriptions of the symbols $g \in \mathcal{H}(\mathbb{D})$ such that the operators T_g , S_g and M_g are bounded, or compact, on A_α^p .

Theorem 2.3. *Let $g \in \mathcal{H}(\mathbb{D})$, $0 < p < \infty$ and $\alpha \geq -1$. Then:*

- a) $T_g \in \mathcal{B}(A_\alpha^p)$ if and only if $g \in \mathcal{B}$, when $\alpha > -1$, and $g \in BMOA$, when $\alpha = -1$. Moreover, $\|T_g\|_{\alpha,p} \simeq \|g\|_{\mathcal{B}}$, if $\alpha > -1$, and $\|T_g\|_{\alpha,p} \simeq \|g\|_{BMOA}$, if $\alpha = -1$.
- b) $T_g \in \mathcal{K}(A_\alpha^p)$ if and only if $g \in \mathcal{B}_0$, when $\alpha > -1$, and $g \in VMOA$, when $\alpha = -1$.

Theorem 2.3 is originally proved, for $\alpha = -1$, in [4, Thm. 1, Corollary 1] ($p \geq 1$) and in [2, Thm. 1(ii), Corollary 1(ii)] ($0 < p < 1$) and, for $\alpha > -1$, in [5, Thm. 1] ($p \geq 1$) and in [3, Thm. 4.1(i)] ($0 < p < 1$).

Proposition 2.4. *Let $g \in \mathcal{H}(\mathbb{D})$, $0 < p < \infty$ and $\alpha \geq -1$. Then:*

- a) $S_g \in \mathcal{B}(A_\alpha^p)$ (or $M_g \in \mathcal{B}(A_\alpha^p)$) if and only if $g \in H^\infty$. Moreover, $\|S_g\|_{\alpha,p} \simeq \|M_g\|_{\alpha,p} \simeq \|g\|_{H^\infty}$.
- b) $S_g \in \mathcal{K}(A_\alpha^p)$ (or $M_g \in \mathcal{K}(A_\alpha^p)$) if and only if $g \equiv 0$.

The characterization of the boundedness for M_g follows from a classical result on pointwise multipliers (see [7, Lemma 11] or [14, Lemma 1.10]). The remaining part of Proposition 2.4 is well known for the experts, but unfortunately we have not found any explicit reference. For a sake of completeness we include a sketch of the proof. If $g \in H^\infty$ then $M_g, T_g, g(0)\delta_0 \in \mathcal{B}(A_\alpha^p)$, and so $S_g \in \mathcal{B}(A_\alpha^p)$, by (1.1). In order to prove the converse, recall that the Bergman kernel for A_α^2 is $K_\alpha(z, \lambda) = (1 - \bar{\lambda}z)^{-\alpha-2}$, and, in particular, the analytic function

$$h_\lambda(z) = \frac{(1 - |\lambda|^2)^{\frac{\alpha+2}{p}}}{(1 - \bar{\lambda}z)^{\frac{2\alpha+4}{p}}} \quad (\lambda \in \mathbb{D})$$

satisfies $\|h_\lambda\|_{\alpha,p} = 1$. Thus if $S_g \in \mathcal{B}(A_\alpha^p)$ then

$$|(S_g h_\lambda)'(\lambda)| \leq \frac{c_{\alpha,p}}{(1 - |\lambda|^2)^{\frac{\alpha+2}{p}+1}} \|S_g h_\lambda\|_{\alpha,p} \leq \frac{c_{\alpha,p} \|S_g\|_{\alpha,p}}{(1 - |\lambda|^2)^{\frac{\alpha+2}{p}+1}},$$

from which follows that $\|g\|_{H^\infty} \leq C_{\alpha,p} \|S_g\|_{\alpha,p}$. A similar argument shows that if $M_g \in \mathcal{B}(A_\alpha^p)$ then $g \in H^\infty$.

Using standard arguments on compact operators between spaces of analytic functions (see Lemma 4.10) together with the above estimates it is easy to prove part b) of Proposition 2.4.

3. THE ALGEBRA \mathcal{A}_g GENERATED BY THE OPERATORS T_g , S_g , AND M_g .

The main goal of this section is to show that any operator $L \in \mathcal{A}_g$ has a unique representation of the form (1.3) when g is non constant and $g(0) \neq 0$. A powerful purely algebraic machinery which helps dealing with such questions are the Gröbner bases [1], [13]. However, we have preferred a direct approach, partly for the sake of completeness, but also because our further

arguments need some more specific information about this representation, like for example Proposition 3.7 below.

3.1. Some useful identities. In this section we gather some formulas that will be used later on.

Proposition 3.1. *Let $g \in \mathcal{H}(\mathbb{D})$, and $j, k \in \mathbb{N}$. Then:*

$$\begin{aligned}
(3.1) \quad & M_g = S_g + T_g + g(0) \delta_0 \\
(3.2) \quad & M_g^k = M_{g^k} \\
(3.3) \quad & S_g^k = S_{g^k} \\
(3.4) \quad & S_{g^j} T_{g^k} = \frac{k}{j+k} T_{g^{j+k}} \\
(3.5) \quad & S_{g^j} M_{g^k} = S_{g^{j+k}} + \frac{k}{j+k} T_{g^{j+k}} \\
(3.6) \quad & T_{g^j} M_{g^k} = \frac{j}{j+k} T_{g^{j+k}} \\
(3.7) \quad & T_g S_g = S_g T_g - T_g^2 - g(0)(g - g(0)) \delta_0
\end{aligned}$$

Proof. Let $f \in \mathcal{H}(\mathbb{D})$.

(3.1) Since $(gf)' = g'f + gf'$, we have

$$g(z)f(z) = g(0)f(0) + \int_0^z g'(\zeta)f(\zeta) d\zeta + \int_0^z g(\zeta)f'(\zeta) d\zeta,$$

that is, $M_g f = T_g f + S_g f + g(0) \delta_0 f$.

(3.2) $M_g^k f = g^k f = M_{g^k} f$.

(3.3) We proceed by induction on k . For $k = 1$ there is nothing to prove. Now assume that $S_g^k = S_{g^k}$. Then

$$S_g^{k+1} f(z) = S_g(S_{g^k} f)(z) = \int_0^z g(\zeta)^{k+1} f'(\zeta) d\zeta = S_{g^{k+1}} f(z),$$

that is, $S_g^{k+1} = S_{g^{k+1}}$.

(3.4) It follows by integration from the identity $g^j (T_{g^k} f)' = \frac{k}{j+k} (g^{j+k})' f$.

(3.5) It follows from (3.1), (3.3) and (3.4):

$$S_{g^j} M_{g^k} = S_{g^j} S_{g^k} + S_{g^j} T_{g^k} = S_{g^{j+k}} + \frac{k}{j+k} T_{g^{j+k}}$$

(3.6) It follows by integration from the identity $(g^j)' M_{g^k} f = \frac{j}{j+k} (g^{j+k})' f$.

(3.7) It follows from (3.1), (3.6) and (3.4) :

$$\begin{aligned}
T_g S_g &= T_g M_g - T_g^2 - g(0)(g - g(0)) \delta_0 \\
&= S_g T_g - T_g^2 - g(0)(g - g(0)) \delta_0
\end{aligned}$$

□

Proposition 3.2. *Let $g \in \mathcal{H}(\mathbb{D})$, then*

$$(3.8) \quad T_g (g - g(0))^n = \frac{1}{n+1} (g - g(0))^{n+1} \quad (n \in \mathbb{N} \cup \{0\})$$

$$(3.9) \quad S_g (g - g(0))^n = g(0)(g - g(0))^n + \frac{n}{n+1} (g - g(0))^{n+1} \quad (n \in \mathbb{N})$$

Proof. Identity (3.8) is a direct computation, while (3.9) is easily checked:

$$\begin{aligned} S_g(g - g(0))^n(z) &= n \int_0^z g(\zeta)g'(\zeta)(g(\zeta) - g(0))^{n-1}d\zeta \\ &= n \int_0^z g'(\zeta)(g(\zeta) - g(0))^n d\zeta \\ &\quad + ng(0) \int_0^z g'(\zeta)(g(\zeta) - g(0))^{n-1}d\zeta \\ &= \frac{n}{n+1}(g(z) - g(0))^{n+1} + g(0)(g(z) - g(0))^n. \quad \square \end{aligned}$$

Corollary 3.3. *Let $g \in \mathcal{H}(\mathbb{D})$ and let P be a polynomial of degree n . Then:*

- a) $T_g P(g - g(0)) = Q(g - g(0))$, where $Q(z) = \int_0^z P(\zeta) d\zeta$ is a polynomial of degree $n + 1$.
- b) $S_g P(g - g(0)) = Q(g - g(0))$, with $Q(z) = g(0)(P(z) - P(0)) + \int_0^z \zeta P'(\zeta) d\zeta$, which is a polynomial of degree of $n + 1$.

Proof. Part a) directly follows from (3.8). Part b) is a direct consequence of (3.9) and the fact that $S_g 1 = 0$. \square

Corollary 3.4. *Let $g \in \mathcal{H}(\mathbb{D})$ and let $m, n \in \mathbb{N}$. Then*

$$S_g^{m-j} T_g^j (g - g(0))^n = \frac{n!}{(m+n)(n+j-1)!} (g - g(0))^{m+n} + P(g - g(0)) \quad (0 \leq j \leq m),$$

where P is a polynomial of degree less than $m + n$ and whose coefficients only depend on $g(0)$, m , n and j .

Proof. By (3.8) it is clear that

$$T_g^j (g - g(0))^n = \frac{1}{(n+1)\cdots(n+j)} (g - g(0))^{n+j} = \frac{n!}{(n+j)!} (g - g(0))^{n+j}.$$

But (3.9) gives that

$$S_g^{m-j} (g - g(0))^{n+j} = \frac{n+j}{m+n} (g - g(0))^{m+n} + Q(g - g(0)),$$

where Q is a polynomial of degree less than $m + n$ whose coefficients only depend on $g(0)$, m , n and j . Hence the proof is complete. \square

3.2. Vector space structure of \mathcal{A}_g .

Definition 3.5. Let $L \in \mathcal{A}_g^{(n)}$, where $n \in \mathbb{N}$. We say that L admits an *ST-decomposition* if there exists a polynomial P of degree less than n satisfying

$$L = \sum_{k=1}^n \sum_{j=0}^k c_{j,k} S_g^j T_g^{k-j} + g(0)P(g - g(0)) \delta_0,$$

where $c_{j,k} \in \mathbb{C}$, for any j, k .

Proposition 3.6. *Let $g \in \mathcal{H}(\mathbb{D})$ and $n \in \mathbb{N}$. Then every $L \in \mathcal{A}_g^{(n)}$ admits an *ST-decomposition*.*

Proof. We proceed by induction on n . For $n = 1$ there is nothing to prove because (3.1) holds. Let $n > 1$. Since, by the induction hypothesis, any m -letter g -word, with $m \leq n - 1$, admits an *ST-decomposition*, we will complete the proof by induction once we have checked that $L^{(n)} = L_n L^{(n-1)}$ has an *ST-decomposition*, when L_n is either S_g , T_g , or M_g and $L^{(n-1)}$ is

either $g(0)P(g-g(0))\delta_0$, where P is a polynomial of degree less than $n-1$, or $S_g^j T_g^{k-j}$, where $0 \leq j \leq k$ and $1 \leq k \leq n-1$.

Assume first that $L^{(n-1)} = g(0)P(g-g(0))\delta_0$. By the identity (3.1) we only need to consider the case when L_n is either T_g or S_g . Then, by Corollary 3.3, $L^{(n)} = g(0)Q(g-g(0))\delta_0$, where Q is a polynomial of degree less than n .

Now assume that $L^{(n-1)} = S_g^j T_g^{k-j}$. As above, we only need to consider the cases $L_n = S_g$ and $L_n = T_g$. If $L_n = S_g$ then $L^{(n)} = S_g^{j+1} T_g^{k-j}$, and, in particular, $L^{(n)}$ has an ST -decomposition. Now consider the case $L_n = T_g$. If $j = 0$ then $L^{(n)} = T_g^{k+1}$ and we are done. If $j = k = 1$, then $L^{(n)} = T_g S_g = S_g T_g - T_g^2 - g(0)(g-g(0))\delta_0$, by (3.7), so we also are done. Finally, if $j > 1$ and $k > 1$ then, again by (3.7), we have that

$$L^{(n)} = S_g T_g S_g^{j-1} T_g^{k-j} - T_g^2 S_g^{j-1} T_g^{k-j},$$

because $\delta_0 S_g = 0$. Since $T_g S_g^{j-1} T_g^{k-j}$ and $T_g^2 S_g^{j-1} T_g^{k-j-1}$ are g -words with less than n letters, they admit ST -decompositions, by the induction hypothesis. It directly follows that $L^{(n)}$ also has an ST -decomposition. \square

From now on, in order to simplify the notation, we will write $g_0 = g-g(0)$. By the above proposition, any non-trivial g -operator L can be written as

$$(3.10) \quad L = \sum_{k=0}^n S_g^k T_g P_k(T_g) + S_g P_{n+1}(S_g) + g(0)P_{n+2}(g_0)\delta_0,$$

where $n \in \mathbb{N} \cup \{0\}$ and P_0, \dots, P_{n+2} are polynomials such that $\deg P_{n+2} < n$ and either $P_n \neq 0$ or $P_{n+1} \neq 0$. In other words, the vector space \mathcal{A}_g is spanned by $\{S_g^j T_g^k : j, k \in \mathbb{N} \cup \{0\}, j+k \geq 1\} \cup \{(g_0)^j \delta_0 : j \in \mathbb{N} \cup \{0\}\}$ when $g(0) \neq 0$, and by $\{S_g^j T_g^k : j, k \in \mathbb{N} \cup \{0\}, j+k \geq 1\}$, when $g(0) = 0$.

Our next goal is proving the uniqueness of the ST -decomposition when the symbol g is non constant and $g(0) \neq 0$. We will need two preliminary results.

Proposition 3.7. *Let $g \in \mathcal{H}(\mathbb{D})$, and let $L = L_1 + g(0)P(g-g(0))\delta_0$, where*

$$L_1 = \sum_{k=1}^m \sum_{j=0}^k c_{j,k} S_g^j T_g^{k-j}$$

is a g -operator of order $m \in \mathbb{N}$ and P is a polynomial of degree less than m . Then there exists an increasing sequence $\{n_i\}_i$ in \mathbb{N} such that $L[(g-g(0))^{n_i}] = P_i(g-g(0))$, where P_i is a polynomial of degree $m+n_i$.

Proof. By Corollary 3.4, for $n \in \mathbb{N}$ and $0 \leq k \leq m$, we have

$$L[(g_0)^{n+k+1}] = L_1[(g_0)^{n+k+1}] = \frac{(n+k+1)!}{m+n+k+1} a_{k,n} (g_0)^{m+n+k+1} + P_k(g_0),$$

where P_k is a polynomial of degree less than $m+n+k+1$ and

$$a_{k,n} = \sum_{j=0}^m \frac{c_{j,m}}{(n+m-j+k)!}.$$

Since L_1 has order m , $(c_{0,m}, c_{1,m}, \dots, c_{m,m}) \neq (0, 0, \dots, 0)$, so we have that $(a_{0,n}, a_{1,n}, \dots, a_{m,n}) \neq (0, 0, \dots, 0)$, provided that

$$(3.11) \quad D_n^{(m)} := \det \left(\frac{1}{(n+m-j+k)!} \right)_{j,k=0}^m \neq 0,$$

and, in particular, there is some $0 \leq k \leq m$ such that $Lg_0^{n+k+1} = P(g_0)$, where P is a polynomial of degree $m+n+k+1$. Thus we only have to check (3.11). In order to do that we recall the so called Pochhammer symbols:

$$(k)_0 = 1 \quad (k)_\ell = k(k+1) \cdots (k+\ell-1) \quad (k, \ell \in \mathbb{N}).$$

Since

$$\frac{1}{(n+m-j+k)!} = \frac{1}{(n+m+k)!} (n+m-j+k+1)_j,$$

we have that $D_n^{(m)} = b_{n,m} \Delta_n^{(m)}$, where $b_{n,m} > 0$ and

$$\Delta_n^{(m)} := \begin{vmatrix} (n+m+1)_0 & (n+m+2)_0 & \cdots & (n+2m+1)_0 \\ (n+m)_1 & (n+m+1)_1 & \cdots & (n+2m)_1 \\ \vdots & \vdots & \ddots & \vdots \\ (n+1)_m & (n+2)_m & \cdots & (n+m+1)_m \end{vmatrix}.$$

But $(\ell)_0 = 1$ and $(\ell+1)_{j+1} - (\ell)_{j+1} = (j+1)(\ell+1)_j$, we have

$$\Delta_n^{(m)} = \begin{vmatrix} 1(n+m+1)_0 & \cdots & 1(n+2m)_0 \\ 2(n+m)_1 & \cdots & 2(n+2m-1)_1 \\ \vdots & \ddots & \vdots \\ m(n+2)_{m-1} & \cdots & m(n+m)_{m-1} \end{vmatrix},$$

and so $\Delta_n^{(m)} = m! \Delta_{n+1}^{(m-1)}$. Since $\Delta_{n+m}^{(1)} = 1$, we get (3.11). \square

Lemma 3.8. *Let $g \in \mathcal{H}(\mathbb{D})$. If g is not constant then $\{g^n : n \in \mathbb{N} \cup \{0\}\}$ and $\{(g-g(0))^n : n \in \mathbb{N} \cup \{0\}\}$ are bases for the vector space $\{P(g) : P \text{ polynomial}\}$.*

Proof. It is clear that $\{g^n : n \in \mathbb{N} \cup \{0\}\}$ and $\{(g_0)^n : n \in \mathbb{N} \cup \{0\}\}$ span the vector space $\{P(g) : P \text{ polynomial}\}$. Now we want to prove that $\{g^n : n \in \mathbb{N} \cup \{0\}\}$ is linearly independent, which means that if $P(g) = 0$, for some polynomial P , then $P \equiv 0$. Thus assume that $P(g) = 0$, for some polynomial P . Since g is not constant, g takes infinitely many values. It follows that P has infinitely many zeros, that is, $P \equiv 0$. A similar argument shows that $\{(g_0)^n : n \in \mathbb{N} \cup \{0\}\}$ is linearly independent, so the proof is complete. \square

Proposition 3.9. *Let $g \in \mathcal{H}(\mathbb{D})$.*

- If $g \neq 0$ is constant and I is the identity mapping on $\mathcal{H}(\mathbb{D})$, then $\{I, \delta_0\}$ is a basis for $\mathcal{A}_g^{(n)}$, for every $n \in \mathbb{N}$, and so it is also a basis for \mathcal{A}_g .*
- If g is not constant and $g(0) = 0$, then*

$$(3.12) \quad \{S_g^j T_g^{k-j} : 1 \leq k \leq n, 0 \leq j \leq k\}$$

is a basis for $\mathcal{A}_g^{(n)}$, and so $\{S_g^j T_g^k : j, k \in \mathbb{N} \cup \{0\}, j+k \geq 1\}$ is a basis for \mathcal{A}_g .

c) If g is not constant and $g(0) \neq 0$, then

$$(3.13) \quad \{S_g^j T_g^{k-j} : 1 \leq k \leq n, 0 \leq j \leq k\} \cup \{(g - g(0))^j \delta_0 : 0 \leq j < n\}$$

is a basis for $\mathcal{A}_g^{(n)}$, and so

$$\{S_g^j T_g^k : j, k \in \mathbb{N} \cup \{0\}, j + k \geq 1\} \cup \{(g - g(0))^j \delta_0 : j \in \mathbb{N} \cup \{0\}\}$$

is a basis for \mathcal{A}_g .

Proof. a) Assume $g \equiv c \neq 0$. Then $T_g = 0$, $S_g = cI - c\delta_0$ and $M_g = cI$, so both $\mathcal{A}_g^{(n)}$ and \mathcal{A}_g are spanned (as vector spaces) by I and δ_0 . On the other hand, I and δ_0 are linearly independent. Indeed, if $\alpha I + \beta \delta_0 = 0$, for some $\alpha, \beta \in \mathbb{C}$, then $\alpha f = (\alpha I + \beta \delta_0)f = 0$, for $f(z) = z$, so $\alpha = 0$, and therefore $\beta = (\alpha I + \beta \delta_0)1 = 0$.

b) Assume g is not constant and $g(0) = 0$. Then Proposition 3.6 shows that $\mathcal{A}_g^{(n)}$ is spanned by (3.12). On the other hand, the linear independence of (3.12) follows from Proposition 3.7. Indeed, if

$$(3.14) \quad \sum_{k=1}^n \sum_{j=0}^k c_{j,k} S_g^{k-j} T_g^j = 0,$$

where $c_{j,k} \in \mathbb{C}$, then $c_{j,k} = 0$, for any $1 \leq k \leq n$ and $0 \leq j \leq k$, since otherwise Proposition 3.7 shows that there is some $\ell \in \mathbb{N}$ such that

$$\left(\sum_{k=1}^n \sum_{j=0}^k c_{j,k} S_g^{k-j} T_g^j \right) g^\ell = P(g),$$

where P is a non-constant polynomial, which is absurd, taking into account (3.14) and Lemma 3.8.

c) Assume g is not constant and $g(0) \neq 0$. First, note that (3.1) shows that $\delta_0 = \frac{1}{g(0)}(M_g - S_g - T_g) \in \mathcal{A}_g^{(1)}$, and so (3.8) gives that

$$(g_0)^j \delta_0 = j! (T_g^j 1) \delta_0 = j! T_g^j \delta_0 \in \mathcal{A}_g^{(n)} \quad (0 \leq j < n).$$

On the other hand, since Proposition 3.6 shows that $\mathcal{A}_g^{(n)}$ is spanned by (3.13), we only have to prove the linear independence of (3.13). Assume that

$$(3.15) \quad \sum_{k=1}^n \sum_{j=0}^k c_{j,k} S_g^{k-j} T_g^j + P(g_0) \delta_0 = 0,$$

where $c_{j,k} \in \mathbb{C}$ and P is a polynomial. Then $c_{j,k} = 0$, for any $1 \leq k \leq n$ and $0 \leq j \leq k$, since otherwise Proposition 3.7 shows that there is some $\ell \in \mathbb{N}$ such that

$$\left(\sum_{k=1}^n \sum_{j=0}^k c_{j,k} S_g^{k-j} T_g^j + P(g_0) \delta_0 \right) (g_0)^\ell = Q(g_0),$$

where Q is a non-constant polynomial, which is absurd, taking into account (3.15) and Lemma 3.8. Therefore

$$P(g_0) = P(g_0) \delta_0 1 = 0,$$

and a second application of Lemma 3.8 gives that $P \equiv 0$. \square

We end this section by giving a second application of Propositions 3.6 and 3.7 (and Lemma 3.8) which clarifies the concept of trivial g -operator. We recall that $L \in \mathcal{A}_g$ is trivial if $L = g(0)P(g_0)\delta_0$, for some polynomial P .

Proposition 3.10. *Let $g \in \mathcal{H}(\mathbb{D})$.*

- a) *If $g(0) = 0$ and $L = P(g)\delta_0 \in \mathcal{A}_g$, for some polynomial P , then $L = 0$.*
- b) *A g -operator L is trivial if and only if $L(z^\ell) = 0$, for every $\ell \in \mathbb{N}$.*

Proof. Assume that $g(0) = 0$ and $L = P(g)\delta_0 \in \mathcal{A}_g$, for some polynomial P . If g is constant then $g \equiv 0$, so $M_g = S_g = T_g = 0$, and therefore $\mathcal{A}_g = 0$, which gives that $L = 0$. When g is not constant we proceed by contradiction. Suppose that $L \neq 0$. Then $L \in \mathcal{A}_g^{(m)}$, for some $m \in \mathbb{N}$, so Propositions 3.6 and 3.7 show that there is $n \in \mathbb{N}$ such that $Lg^n = Q(g)$, where Q is a polynomial of degree $m + n$. But, since $g(0) = 0$, $Lg^n = 0$, so $Q(g) = 0$, and Lemma 3.8 implies that $Q \equiv 0$, which is a contradiction and finishes the proof of part a).

Finally, we prove part b). Now assume that $L \in \mathcal{A}_g$. If L is trivial, it is clear that $L(z^\ell) = 0$, for every $\ell \in \mathbb{N}$. On the other hand, if $L(z^\ell) = 0$, for any $\ell \in \mathbb{N}$, then $LP = L(P(0)) = P(0)(L1)$, for any polynomial P . Now the continuity of $L : \mathcal{H}(\mathbb{D}) \rightarrow \mathcal{H}(\mathbb{D})$ implies that $Lf = f(0)(L1)$, for any $f \in \mathcal{H}(\mathbb{D})$, that is, $L = (L1)\delta_0$. But $L1 = P(g_0)$, where P is a polynomial, and, by part a), we conclude that L is trivial. \square

4. MAIN RESULTS

We start this section by studying the behaviour of the iterates of T_g .

Proposition 4.1. *Let $g \in \mathcal{H}(\mathbb{D})$. If $n \in \mathbb{N}$, $n > 1$, and $T_g^n \in \mathcal{B}(A_\alpha^p)$, then $T_g \in \mathcal{B}(A_\alpha^p)$ and there exists a constant $c_n > 0$, which only depends on n , such that*

$$(4.1) \quad \|T_g f\|_{\alpha,p}^n \leq c_n \|T_g^n f\|_{\alpha,p} \|f\|_{\alpha,p}^{n-1} \quad f \in A_\alpha^p,$$

and so

$$(4.2) \quad \|T_g\|_{\alpha,p}^n \leq c_n \|T_g^n\|_{\alpha,p}.$$

In particular, $T_g^n \in \mathcal{B}(A_\alpha^p)$, for some $n \in \mathbb{N}$, if and only if $T_g \in \mathcal{B}(A_\alpha^p)$, for any $n \in \mathbb{N}$.

In order to prove Proposition 4.1 we need the following useful result, which is proved in [2, Thm. 1 (i)] for $\alpha = -1$, while for $\alpha > -1$ it is a direct consequence of Hölder's inequality and the fact that the differentiation operator $f \mapsto f'$ is a topological isomorphism from $A_\alpha^p(0) = \{f \in A_\alpha^p : f(0) = 0\}$ onto $A_{\alpha+p}^p$ [15, Thm. 4.28].

Lemma 4.2. *Let $r, q, s > 0$, $\frac{1}{r} + \frac{1}{s} = \frac{1}{q}$, and $g \in A_\alpha^r$. Then $T_g : A_\alpha^s \rightarrow A_\alpha^q$ is bounded and there exists a constant $c > 0$, independent of g , satisfying that $\|T_g\|_{A_\alpha^s \rightarrow A_\alpha^q} \leq c \|g\|_{\alpha,r}$.*

Proof of Proposition 4.1. Note that $T_g^n 1$ is a polynomial of degree n in g , so that $g^k \in A_\alpha^p$, $1 \leq k \leq n$. Inductively it follows easily that $g^k \in A_\alpha^p$, for all $k \geq 1$. Then using integration by parts we see that $T_g^k f \in A_\alpha^p$ whenever $k \geq 1$ and f is a polynomial. For $k > 1$, we apply Lemma 4.2

with $r = s = p$, $q = p/2$, to conclude that if f is a polynomial and $h \in A_\alpha^p$ then $T_{T_g^k f} h \in A_{\frac{\alpha}{2}}^{\frac{p}{2}}$ with

$$\|T_{T_g^k f} h\|_{\alpha, \frac{p}{2}} \lesssim \|T_g^k f\|_{\alpha, p} \|h\|_{\alpha, p}.$$

Now, for $k \geq 2$, let $h = T_g^{k-2} f$ and note that

$$T_{T_g^k f} h(z) = \int_0^z g'(\zeta) (T_g^{k-2} f)(\zeta) (T_g^{k-1} f)(\zeta) d\zeta = \frac{1}{2} (T_g^{k-1} f)^2(z).$$

Since $\|(T_g^{k-1} f)^2\|_{\alpha, \frac{p}{2}} = \|T_g^{k-1} f\|_{\alpha, p}^2$, this leads to the estimate

$$(4.3) \quad \|T_g^{k-1} f\|_{\alpha, p}^2 \lesssim \|T_g^k f\|_{\alpha, p} \|T_g^{k-2} f\|_{\alpha, p} \quad (k \geq 2).$$

By induction on $j \geq 1$, from (4.3) we obtain

$$(4.4) \quad \|T_g^{k-j} f\|_{\alpha, p}^{j+1} \lesssim \|T_g^k f\|_{\alpha, p} \|T_g^{k-j-1} f\|_{\alpha, p}^j \quad (k \geq j+1).$$

Indeed, assume that

$$(4.5) \quad \|T_g^{k-j+1} f\|_{\alpha, p}^j \lesssim \|T_g^k f\|_{\alpha, p} \|T_g^{k-j} f\|_{\alpha, p}^{j-1} \quad (k \geq j),$$

and we want to obtain (4.4).

By (4.3), for each $k \geq j+1$ we have

$$\|T_g^{k-j} f\|_{\alpha, p}^{2j} \lesssim \|T_g^{k-j+1} f\|_{\alpha, p}^j \|T_g^{k-j-1} f\|_{\alpha, p}^j.$$

Now, by (4.5), we obtain

$$\|T_g^{k-j} f\|_{\alpha, p}^{2j} \lesssim \|T_g^k f\|_{\alpha, p} \|T_g^{k-j} f\|_{\alpha, p}^{j-1} \|T_g^{k-j-1} f\|_{\alpha, p}^j,$$

which proves (4.4).

Finally, the estimates (4.3) and (4.4) for $k = 2$ and $k - j = 2$, respectively, give that

$$\begin{aligned} \|T_g f\|_{\alpha, p}^2 &\lesssim \|T_g^2 f\|_{\alpha, p} \|f\|_{\alpha, p} \\ \|T_g^2 f\|_{\alpha, p}^{k-1} &\lesssim \|T_g^k f\|_{\alpha, p} \|T_g f\|_{\alpha, p}^{k-2} \quad (k \geq 3). \end{aligned}$$

Therefore

$$\|T_g f\|_{\alpha, p}^{2(k-1)} \lesssim \|T_g^2 f\|_{\alpha, p}^{k-1} \|f\|_{\alpha, p}^{k-1} \lesssim \|T_g^k f\|_{\alpha, p} \|T_g f\|_{\alpha, p}^{k-2} \|f\|_{\alpha, p}^{k-1} \quad (k \geq 3),$$

and so

$$\|T_g f\|_{\alpha, p}^k \lesssim \|T_g^k f\|_{\alpha, p} \|f\|_{\alpha, p}^{k-1}, \quad \text{for any polynomial } f \quad (k \geq 2).$$

In particular, if $k = n$, bearing in mind that the polynomials are dense in A_α^p , the preceding estimate shows that (4.1) holds, and, as a consequence, (4.2) also holds. Hence $T_g \in \mathcal{B}(A_\alpha^p)$. \square

For $h \in \mathcal{H}(\mathbb{D})$ and $\lambda \in \overline{\mathbb{D}}$, let us consider the dilated functions

$$h_\lambda(z) := h(\lambda z), \quad z \in \mathbb{D}.$$

The map $h \mapsto h_\lambda$ is a linear contractive operator on A_α^p . Moreover,

$$(4.6) \quad (M_g f)_\lambda = M_{g_\lambda} f_\lambda \quad (S_g f)_\lambda = S_{g_\lambda} f_\lambda \quad (T_g f)_\lambda = T_{g_\lambda} f_\lambda.$$

Now a repeated application of (4.6) shows that

$$(4.7) \quad L_{g_\lambda} f_\lambda = (L_g f)_\lambda \quad (L_g \in \mathcal{A}_g).$$

The following result is a key tool in our study of the boundedness of operators in \mathcal{A}_g .

Proposition 4.3. *Let $g \in \mathcal{H}(\mathbb{D})$ and let $L_g \in \mathcal{A}_g$. If $L_g \in \mathcal{B}(A_\alpha^p)$ then $L_{g_\lambda} \in \mathcal{B}(A_\alpha^p)$ and $\|L_{g_\lambda}\|_{\alpha,p} \leq \|L_g\|_{\alpha,p}$, for any $\lambda \in \overline{\mathbb{D}}$. Moreover, if $\varliminf_{r \nearrow 1} \|L_{g_r}\|_{\alpha,p} < \infty$, then $L_g \in \mathcal{B}(A_\alpha^p)$ and $\|L_g\|_{\alpha,p} = \varliminf_{r \nearrow 1} \|L_{g_r}\|_{\alpha,p}$.*

Proof. First note that, for any $\lambda \in \mathbb{T}$, (4.7) gives that $L_{g_\lambda} f = (L_g f_{\bar{\lambda}})_\lambda$ and, since $f \mapsto f_\lambda$ is an invertible isometry on A_α^p , it follows that $L_{g_\lambda} \in \mathcal{B}(A_\alpha^p)$ and $\|L_{g_\lambda}\|_{\alpha,p} = \|L_g\|_{\alpha,p}$. If $\lambda \in \mathbb{D}$, then $g_\lambda \in \mathcal{H}(\overline{\mathbb{D}})$, so $M_{g_\lambda}, S_{g_\lambda}, T_{g_\lambda} \in \mathcal{B}(A_\alpha^p)$, and, as a consequence, $L_{g_\lambda} \in \mathcal{B}(A_\alpha^p)$.

In order to estimate the operator norm of L_{g_λ} , let f be a polynomial and observe that, for fixed $z \in \mathbb{D}$, the function $\lambda \mapsto L_{g_\lambda} f(z)$ is analytic on \mathbb{D} . Indeed, this is an immediate consequence of the fact that if $(\lambda, z) \mapsto h(\lambda, z)$ is an analytic function on the bidisc \mathbb{D}^2 then

$$M_{g_\lambda} h(\lambda, \cdot)(z), \quad S_{g_\lambda} h(\lambda, \cdot)(z) \quad \text{and} \quad T_{g_\lambda} h(\lambda, \cdot)(z)$$

are also analytic functions of (λ, z) on \mathbb{D}^2 .

Next we are going to show that $F(\lambda) = L_{g_\lambda} f$ defines a continuous mapping from $\overline{\mathbb{D}}$ to A_α^p .

Assume first that $\zeta \in \mathbb{D}$. For each $z \in \mathbb{D}$ the function $\lambda \mapsto L_{g_\lambda} f(z)$ is analytic on \mathbb{D} , which implies that $L_{g_\lambda} f(z) \rightarrow L_{g_\zeta} f(z)$, as $\lambda \rightarrow \zeta$. Since $L_{g_\lambda} f$ is uniformly bounded on \mathbb{D} , for $|\lambda - \zeta| < \frac{1}{2}(1 - |\zeta|)$, the Dominated Convergence Theorem shows that $\|F(\lambda) - F(\zeta)\|_{\alpha,p} \rightarrow 0$, as $\lambda \rightarrow \zeta$.

If $\zeta \in \mathbb{T}$, we write, by abuse of notation, $f_{1/\lambda}(z) = f(z/\lambda)$, which is well defined for a polynomial f and $\lambda \in \mathbb{C} \setminus \{0\}$. Then, by (4.7), for any $\lambda \in \overline{\mathbb{D}} \setminus \{0\}$ we have that

$$F(\lambda) - F(\zeta) = L_{g_\lambda} f - L_{g_\zeta} f = (L_g f_{1/\lambda} - L_g f_{\bar{\zeta}})_\lambda + (L_g f_{\bar{\zeta}})_\lambda - (L_g f_{\bar{\zeta}})_\zeta,$$

and so

$$\begin{aligned} \|F(\lambda) - F(\zeta)\|_{\alpha,p} &\leq c \left[\|(L_g f_{1/\lambda} - L_g f_{\bar{\zeta}})_\lambda\|_{\alpha,p} + \|(L_g f_{\bar{\zeta}})_\lambda - (L_g f_{\bar{\zeta}})_\zeta\|_{\alpha,p} \right] \\ &\leq c \left[\|L_g f_{1/\lambda} - L_g f_{\bar{\zeta}}\|_{\alpha,p} + \|(L_g f_{\bar{\zeta}})_\lambda - (L_g f_{\bar{\zeta}})_\zeta\|_{\alpha,p} \right] \\ &\leq c \left[\|L_g\| \|f_{1/\lambda} - f_{\bar{\zeta}}\|_{\alpha,p} + \|(L_g f_{\bar{\zeta}})_\lambda - (L_g f_{\bar{\zeta}})_\zeta\|_{\alpha,p} \right], \end{aligned}$$

where $c = 1$ if $p \geq 1$, and $c = 2^{1/p}$ if $0 < p < 1$. Recall that f is a polynomial and use the elementary fact that, for $h \in A_\alpha^p$, $\|h_\lambda - h_\zeta\|_{\alpha,p} \rightarrow 0$, as $\lambda \rightarrow \zeta$, to conclude that the right hand side converges to 0 and therefore $\|F(\lambda) - F(\zeta)\|_{\alpha,p} \rightarrow 0$, as $\lambda \rightarrow \zeta$.

Hence we have just proved that $F : \overline{\mathbb{D}} \rightarrow A_\alpha^p$ is continuous, and, as a consequence, the function $u_f : \overline{\mathbb{D}} \rightarrow \mathbb{C}$, defined by

$$u_f(\lambda) = \|F(\lambda)\|_{\alpha,p}^p = \|L_{g_\lambda} f\|_{\alpha,p}^p,$$

is also continuous. Moreover, since, for fixed $z \in \mathbb{D}$, $L_{g_\lambda} f(z)$ is an analytic function on λ , it is clear that u_f is subharmonic in \mathbb{D} . It follows that u_f attains its maximum at some point $\zeta \in \mathbb{T}$, which gives that

$$\|L_{g_\lambda} f\|_{\alpha,p} = u_f(\lambda) \leq \|L_{g_\zeta} f\|_{\alpha,p} \leq \|L_{g_\zeta}\| \|f\|_{\alpha,p} = \|L_g\| \|f\|_{\alpha,p},$$

for any $\lambda \in \overline{\mathbb{D}}$ and for any polynomial f . Since the polynomials are dense in A_α^p , we conclude that $\|L_{g_\lambda}\|_{\alpha,p} \leq \|L_g\|_{\alpha,p}$.

Finally, for any $f \in A_\alpha^p$, Fatou's lemma shows that

$$\begin{aligned} \|L_g f\|_{\alpha,p} &\leq \varliminf_{r \nearrow 1} \|(L_g f)_r\|_{\alpha,p} = \varliminf_{r \nearrow 1} \|L_{g_r} f_r\|_{\alpha,p} \\ &\leq \varliminf_{r \nearrow 1} \|L_{g_r}\|_{\alpha,p} \|f_r\|_{\alpha,p} \leq \varliminf_{r \nearrow 1} \|L_{g_r}\|_{\alpha,p} \|f\|_{\alpha,p}. \quad \square \end{aligned}$$

4.1. Proof of Theorem 1.1. From now on we shall use repeatedly the following elementary fact:

Remark 4.4. If a function $\varphi : [0, \infty) \rightarrow \mathbb{R}$ satisfies $\lim_{x \rightarrow \infty} \varphi(x) = \infty$, then the preimage by φ of any bounded set of real numbers is bounded.

We will also need a couple of preliminary results.

Lemma 4.5. *Let $g \in \mathcal{H}(\mathbb{D})$ and let P be a polynomial of degree $n \geq 1$. If $P(g) \in H^\infty$, then $g \in H^\infty$.*

Proof. Assume that $P(g) \in H^\infty$, where $P(z) = \sum_{k=0}^n a_k z^k$ is a polynomial of degree $n \geq 1$. Then

$$|a_n| |g(z)|^n - \sum_{k=0}^{n-1} |a_k| |g(z)|^k \leq \|P(g)\|_\infty \quad (z \in \mathbb{D}),$$

and so Remark 4.4 completes the proof. \square

Lemma 4.6. *Let $g \in \mathcal{H}(\mathbb{D})$ and let P be a polynomial of degree $n \geq 1$. If $P(T_g) \in \mathcal{B}(A_\alpha^p)$, then $T_g \in \mathcal{B}(A_\alpha^p)$.*

Proof. Assume that $P(T_g) \in \mathcal{B}(A_\alpha^p)$, where $P(z) = \sum_{k=0}^n a_k z^k$ is a polynomial of degree $n \geq 1$. Then, by Proposition 4.3,

$$|a_n| \|T_{g_r}^n\|_{\alpha,p} - c_{n,p} \sum_{k=0}^{n-1} |a_k| \|T_{g_r}\|_{\alpha,p}^k \leq \|P(T_{g_r})\|_{\alpha,p} \leq \|P(T_g)\|_{\alpha,p} \quad (0 < r < 1).$$

Now Proposition 4.1 shows that $\|T_{g_r}^n\|_{\alpha,p} \geq c_n \|T_{g_r}\|_{\alpha,p}^n$, for some constant $c_n > 0$ only dependent on n . Thus $\varphi(\|T_{g_r}\|_{\alpha,p}) \leq \|P(T_g)\|_{\alpha,p}$, for every $0 < r < 1$, where $\varphi(x) = c_n |a_n| x^n - c_{n,p} \sum_{k=0}^{n-1} |a_k| x^k$. Hence Remark 4.4 and Proposition 4.3 end the proof. \square

Proof of Theorem 1.1 b). Let be $P(z) = a_m z^m + Q(z)$, where Q is a polynomial of degree less than m . Then, by (4.7)

$$g(0)^2 P((g_0)_r) = (L_g g)_r = L_{g_r} g_r \quad (0 < r < 1)$$

so, since $\|L_{g_r} g_r\|_{\alpha,p} \leq \|L\|_{\alpha,p} \|g_r\|_{\alpha,p}$ (see Proposition 4.3), we obtain the estimate

$$\begin{aligned} |g(0)|^2 |a_m| \|(g_0)_r^m\|_{\alpha,p} &= \|L_{g_r} g_r - g(0)^2 Q((g_0)_r)\|_{\alpha,p} \\ &\lesssim \sum_{j=0}^{m-1} \|(g_0)_r^j\|_{\alpha,p} \lesssim \sum_{j=0}^{m-1} \|(g_0)_r^m\|_{\alpha,p}^{j/m}. \end{aligned}$$

Therefore Remark 4.4 implies that $\sup_{0 < r < 1} \|(g_0)_r^m\|_{\alpha,p} < \infty$, and hence Fatou's lemma shows that $g_0^m \in A_\alpha^p$, which means that $g^m \in A_\alpha^p$. \square

Prior to proving Theorem 1.1 a) some definitions and results about the theory of iterated commutators are needed. Let $A, B : \mathcal{H}(\mathbb{D}) \rightarrow \mathcal{H}(\mathbb{D})$ be two linear operators. The *commutator* of A and B is the linear operator $[A, B] := AB - BA$. If $C, D : \mathcal{H}(\mathbb{D}) \rightarrow \mathcal{H}(\mathbb{D})$ are linear operators which commute with B then

$$(4.8) \quad [CAD, B] = C[A, B]D.$$

The *iterated commutators* $[A, B]_k$, $k \in \mathbb{N}$, are defined inductively as follows:

$$[A, B]_1 := [A, B] \quad \text{and} \quad [A, B]_{k+1} := [[A, B]_k, B], \quad \text{for } k \in \mathbb{N}.$$

We will use the following formula

$$(4.9) \quad [A, B]_k = \sum_{j=0}^k (-1)^j \binom{k}{j} B^j A B^{k-j} \quad (k \in \mathbb{N}).$$

Proposition 4.7. *Let $g \in \mathcal{H}(\mathbb{D})$ and $k \in \mathbb{N}$, then*

$$(4.10) \quad [S_g^k, T_g] = T_g T_{g^k} + g(0)^k (g - g(0)) \delta_0$$

Proof. By (3.4) and (3.6) we have that $S_g^k T_g = S_{g^k} T_g = T_g M_{g^k}$, so (3.1) gives that

$$S_g^k T_g = T_g T_{g^k} + T_g S_{g^k} + g(0)^k T_g \delta_0 = T_g T_{g^k} + T_g S_g^k + g(0)^k g_0 \delta_0,$$

which is just (4.10). \square

Proposition 4.8. *Let $g \in \mathcal{H}(\mathbb{D})$ and $k \in \mathbb{N}$, then*

$$(4.11) \quad \left[\frac{(g-g(0))^k}{k!} \delta_0, T_g \right]_j = (-1)^j \frac{(g-g(0))^{k+j}}{(k+j)!} \delta_0 \quad (j, k \in \mathbb{N}).$$

Proof. Observe that (4.11) follows by induction on j from (3.8). Indeed, (3.8) directly shows (4.11) for $j = 1$:

$$\left[\frac{(g_0)^k}{k!} \delta_0, T_g \right] = -T_g \left(\frac{(g_0)^k}{k!} \right) \delta_0 = -\frac{(g_0)^{k+1}}{(k+1)!} \delta_0.$$

Moreover, if $\left[\frac{(g_0)^k}{k!} \delta_0, T_g \right]_j = (-1)^j \frac{(g_0)^{k+j}}{(k+j)!} \delta_0$ holds, then (3.8) gives that

$$\left[\frac{(g_0)^k}{k!} \delta_0, T_g \right]_{j+1} = (-1)^{j+1} T_g \left(\frac{(g_0)^{k+j}}{(k+j)!} \right) \delta_0 = (-1)^{j+1} \frac{(g_0)^{k+j+1}}{(k+j+1)!} \delta_0. \quad \square$$

Corollary 4.9. *Let $g \in \mathcal{H}(\mathbb{D})$ and $k \in \mathbb{N}$, then*

$$(4.12) \quad [S_g^k, T_g]_j = \frac{k!}{(k-j)!} T_g^j S_g^{k-j} T_g^j - \frac{(-1)^j}{j!} g(0)^k (g - g(0))^j \delta_0 \quad (1 \leq j \leq k).$$

Moreover,

$$(4.13) \quad [S_g^k, T_g]_j = -\frac{(-1)^j}{j!} g(0)^k (g - g(0))^j \delta_0 \quad (j > k).$$

Proof. We prove (4.12) by induction on j . For $j = 1$, (4.12) is just (4.10). Now fix $1 \leq j < k$ and assume that

$$[S_g^k, T_g]_j = \frac{k!}{(k-j)!} T_g^j S_g^{k-j} T_g^j - \frac{(-1)^j}{j!} g(0)^k (g_0)^j \delta_0.$$

Then (4.8) and (3.8) show that

$$[S_g^k, T_g]_{j+1} = \frac{k!}{(k-j)!} T_g^j [S_g^{k-j}, T_g] T_g^j - \frac{(-1)^{j+1}}{(j+1)!} g(0)^k (g_0)^{j+1} \delta_0.$$

Therefore (4.10) implies that

$$[S_g^k, T_g]_{j+1} = \frac{k!}{(k-j-1)!} T_g^{j+1} S_g^{k-j-1} T_g^{j+1} - \frac{(-1)^{j+1}}{(j+1)!} g(0)^k (g_0)^{j+1} \delta_0.$$

Thus (4.12) is proved. In particular,

$$[S_g^k, T_g]_k = k! T_g^{2k} - \frac{(-1)^k}{k!} g(0)^k (g_0)^k \delta_0,$$

and so, for $j > k$, (4.11) implies that

$$[S_g^k, T_g]_j = -(-1)^k g(0)^k \left[\frac{(g_0)^k}{k!} \delta_0, T_g \right]_{j-k} = -\frac{(-1)^j}{j!} g(0)^k (g_0)^j \delta_0. \quad \square$$

Proof of Theorem 1.1 a). First of all, we observe that if L is not trivial and $L \in \mathcal{B}(A_\alpha^p)$, then $g^k \in A_\alpha^p$, for any $k \in \mathbb{N}$. Indeed, Propositions 3.6 and 3.7 show that there is an strictly increasing sequence $\{k_j\}$ in \mathbb{N} such that $L(g_0^{k_j}) = P_j(g_0)$, where P_j is a polynomial of degree $d_j > k_j$. Then arguing as in the proof of part b) we obtain that $g^{d_j} \in A_\alpha^p$, and consequently, $g^k \in A_\alpha^p$, for every $k \in \mathbb{N}$.

Now we prove part a). Taking into account Proposition 3.6, Lemma 4.6, and the above observation, we may assume that

$$L_g = P_0(T_g) + \sum_{k=1}^n S_g^k P_k(T_g),$$

where P_0, \dots, P_n are polynomials, and P_n has degree $m \geq 1$.

On the other hand, since $P_k(T_{g_r})$ commute with T_{g_r} , (4.8), (4.12) and (4.13) give that

$$\begin{aligned} [L_{g_r}, T_{g_r}]_n &= n! T_{g_r}^{2n} P_n(T_{g_r}) + g(0) Q_0(g_r - g(0)) \delta_0 \\ &= Q_n(T_{g_r}) + g(0) Q_0(g_r - g(0)) \delta_0, \end{aligned}$$

where Q_n and Q_0 are polynomials and Q_n has degree $N = 2n + m > n$. Now (4.9) and Proposition 4.3 imply that

$$\|[L_{g_r}, T_{g_r}]_n\|_{\alpha,p} \leq c_{n,p} \|L_{g_r}\|_{\alpha,p} \|T_{g_r}\|_{\alpha,p}^n \leq c_{n,p} \|L_g\|_{\alpha,p} \|T_{g_r}\|_{\alpha,p}^n.$$

Moreover, $\|Q_0(g_r - g(0)) \delta_0\|_{\alpha,p} \leq \|Q_0(g_0) \delta_0\|_{\alpha,p} = C < \infty$, by Theorem 1.1 b) and Proposition 4.3. On the other hand, if $Q_n(z) = \sum_{k=0}^N a_k z^k$, then, taking into account Proposition 4.1, we have

$$c_N |a_N| \|T_{g_r}\|_{\alpha,p}^N - c'_{N,p} \sum_{k=0}^{N-1} |a_k| \|T_{g_r}\|_{\alpha,p}^k \leq \|Q_n(T_{g_r})\|_{\alpha,p}.$$

Therefore, putting all that together, we get that $\varphi(\|T_{g_r}\|_{\alpha,p}) \leq C$, where

$$\varphi(x) = c_N |a_N| x^N - c'_{N,p} \sum_{k=0}^{N-1} |a_k| x^k - c_{n,p} \|L_g\|_{\alpha,p} x^n.$$

Hence Remark 4.4 and Proposition 4.3 conclude the proof. \square

4.2. Proof of Theorem 1.2. In order to give the proof, we need the following well known characterization of compact operators.

Lemma 4.10 ([12, Lemma 3.7]). *Let X and Y be two Banach (or quasi-Banach) spaces of analytic functions on \mathbb{D} , and let $T : X \rightarrow Y$ be a linear operator. Suppose that the following conditions are satisfied:*

- (a) *The point evaluation functionals on Y are bounded.*
- (b) *The closed unit ball of X is a compact subset of $\mathcal{H}(\mathbb{D})$, where $\mathcal{H}(\mathbb{D})$ is endowed with the topology of uniform convergence on compacta.*
- (c) *$T : X \rightarrow Y$ is continuous, where both X and Y are endowed with the topology of uniform convergence on compacta.*

Then $T : X \rightarrow Y$ is a compact operator if and only if for any bounded sequence $\{f_j\}$ in X such that $f_j \rightarrow 0$ uniformly on compacta, the sequence $\{Tf_j\}$ converges to zero in the norm of Y .

It is worth mentioning that conditions (a) and (b) of the previous lemma hold when $X = Y = A_\alpha^p$, and in such a case any g -operator satisfies (c).

Proof of Theorem 1.2 a). If $g \in H^\infty$, then $S_g, T_g \in \mathcal{B}(A_\alpha^p)$, by Theorem 2.3 a)) and Proposition 2.4 a), and so $L_g \in \mathcal{B}(A_\alpha^p)$. Conversely, assume that $L_g \in \mathcal{B}(A_\alpha^p)$ and apply Proposition 4.3 to conclude that for $r \in (0, 1)$, we have $L_{g_r} \in \mathcal{B}(A_\alpha^p)$ with $\|L_{g_r}\|_{\alpha,p} \leq \|L_g\|_{\alpha,p}$. From

$$L_{g_r} = \sum_{k=0}^n S_{g_r}^k T_{g_r} P_k(T_{g_r}) + S_{g_r} P_{n+1}(S_{g_r}) + g_r(0) P_{n+2}(g_r - g_r(0)) \delta_0,$$

we see that for fixed $r \in (0, 1)$, all operators on the right are compact, except

$$S_{g_r} P_{n+1}(S_{g_r}) = S_{g_r P_{n+1}(g_r)} = M_{g_r P_{n+1}(g_r)} - T_{g_r P_{n+1}(g_r)} - g_r P_{n+1}(g_r)(0) \delta_0.$$

By Theorem 2.3 b) we conclude that

$$L_{g_r} = M_{g_r P_{n+1}(g_r)} + K,$$

where $K \in \mathcal{K}(A_\alpha^p)$ is compact.

Now, for any $\lambda \in \mathbb{D}$, we consider the functions

$$h_\lambda(z) = \frac{(1 - |\lambda|^2)^{\frac{\alpha+2}{p}}}{(1 - \bar{\lambda}z)^{\frac{2\alpha+4}{p}}} \quad (z \in \mathbb{D}).$$

Since $(1 - \bar{\lambda}z)^{-\alpha-2}$, $\lambda, z \in \mathbb{D}$, is the Bergman kernel for A_α^2 , $\|h_\lambda\|_{\alpha,p} = 1$, for any $\lambda \in \mathbb{D}$. (Note that for $\alpha = -1$ the corresponding Bergman kernel is the classical Cauchy kernel.) Moreover, it is clear that, for any $\zeta \in \mathbb{T}$, $h_\lambda \rightarrow 0$, as $\lambda \rightarrow \zeta$, uniformly on compacta. So, by Lemma 4.10, $\|Kh_\lambda\|_{\alpha,p} \rightarrow 0$. On the other hand, note that if $G_r = g_r P_{n+1}(g_r)$ then

$$\|M_{G_r} h_\lambda\|_{\alpha,p}^p = (\alpha+1) \int_{\mathbb{D}} B_\alpha(z, \lambda) |G_r(z)|^p (1 - |z|^2)^\alpha dA(z) \quad (\lambda \in \mathbb{D}, \alpha > -1),$$

and

$$\|M_{G_r} h_\lambda\|_{-1,p}^p = \int_{\mathbb{T}} P(\zeta, \lambda) |G_r(\zeta)|^p \frac{|d\zeta|}{2\pi} \quad (\lambda \in \mathbb{D}),$$

where $B_\alpha(z, \lambda) = \frac{(1 - |\lambda|^2)^{\alpha+2}}{|1 - \bar{\lambda}z|^{2\alpha+4}}$ and $P(\zeta, \lambda) = \frac{1 - |\lambda|^2}{|1 - \bar{\lambda}\zeta|^2}$ are the Poisson-Bergman (or Berezin) kernel and the classical Poisson kernel, respectively. Thus, since $|G_r| = |g_r P_{n+1}(g_r)| \in C(\overline{\mathbb{D}})$, we have that (see, for instance, [9, Prop. 8.2.7])

$$\lim_{\lambda \rightarrow \zeta} \|M_{g_r P_{n+1}(g_r)} h_\lambda\|_{\alpha, p}^p = |g_r P_{n+1}(g_r)(\zeta)|^p \quad (\zeta \in \mathbb{T}).$$

Hence

$$|g_r P_{n+1}(g_r)(\zeta)|^p = \lim_{\lambda \rightarrow \zeta} \|L_{g_r} h_\lambda\|_{\alpha, p}^p \leq \|L_g\|_{\alpha, p}^p \lim_{\lambda \rightarrow \zeta} \|h_\lambda\|_{\alpha, p}^p = \|L_g\|_{\alpha, p}^p,$$

for all $\zeta \in \mathbb{T}$ and $0 < r < 1$, which implies that $g P_{n+1}(g) \in H^\infty$, and so $g \in H^\infty$, by Lemma 4.5. Thus the proof is finished. \square

Proof of Theorem 1.2 b). First of all observe that if $L \in \mathcal{B}(A_\alpha^p)$ then Theorem 1.1 a) gives that $g^k \in A_\alpha^p$ for any $k \in \mathbb{N}$, so $g(0)P_{n+1}(g_0)\delta_0 \in \mathcal{B}(A_\alpha^p)$ and therefore

$$\sum_{k=0}^n S_g^k T_g P_k(T_g) = L - g(0)P_{n+1}(g_0)\delta_0 \in \mathcal{B}(A_\alpha^p).$$

Moreover, if either $g^{n+1} \in \mathcal{B}$, if $\alpha > -1$, or $g^{n+1} \in BMOA$, if $\alpha = -1$, then Proposition 2.1 shows that $g \in A_\alpha^p$, and we deduce that $g(0)P_{n+1}(g_0)\delta_0 \in \mathcal{B}(A_\alpha^p)$.

Thus, without loss generality, from now on we assume that

$$L = \sum_{k=0}^n S_g^k T_g P_k(T_g).$$

If $g^n \in \mathcal{B}$ when $\alpha > -1$, or $g^n \in BMOA$ when $\alpha = -1$, then, by Proposition 2.1, the same holds for g^k , $1 \leq k \leq n$, and all the operators involved in the definition of L are bounded, hence so is L .

Conversely, if $L = L_g$ is bounded, then by Theorem 1.1 a) we have that T_g is bounded. Now, by applying Proposition 4.3, Proposition 2.1 and Theorem 2.3, for every $0 < r < 1$, we obtain

$$\begin{aligned} \|T_{g_r^{n+1}}\| &\leq c \left(\|L_{g_r}\| + \sum_{k=1}^{n-1} \|P_k(T_{g_r})\| \|T_{g_r^{k+1}}\| \right) \\ &\leq c'(g, n, p) \left(\|L_g\| + \sum_{k=1}^{n-1} \|T_{g_r^{n+1}}\|^{\frac{k+1}{n+1}} \right), \end{aligned}$$

where $0 < c' = c'(g, n, p) < \infty$ because T_g is bounded. Then Remark 4.4 and Proposition 4.3 complete the proof. \square

We will use Proposition 2.1 and the following result in the proof of Theorem 1.2 c).

Lemma 4.11. *Let $g \in \mathcal{B}$, and, for any $\lambda \in \mathbb{D} \setminus \{0\}$ and $\gamma > 0$, let*

$$f_{\gamma, \lambda}(z) = \frac{z}{(1 - \bar{\lambda}z)^\gamma} \quad (z \in \mathbb{D}).$$

Then:

a) For any $k \in \mathbb{N}$ and $t \in [0, 1]$, we have that

$$|T_g^k f_{\gamma, \lambda}(t\lambda)| \leq \frac{\|g\|_{\mathcal{B}}^k}{|\lambda|^k \gamma^k (1 - t|\lambda|^2)^\gamma}.$$

b) If $a_0, \dots, a_n \in \mathbb{C}$ and $\gamma|\lambda| > \|g\|_{\mathcal{B}}$, then

$$\left| \sum_{k=0}^n a_k T_g^k f_{\gamma, \lambda}(\lambda) \right| \leq |a_0| \frac{|\lambda|}{(1 - |\lambda|^2)^\gamma} + \left(\sum_{k=1}^n |a_k| \right) \frac{\|g\|_{\mathcal{B}}}{|\lambda| \gamma (1 - |\lambda|^2)^\gamma}.$$

Proof. First we prove a) by induction on k . If $k = 1$ and $s \in [0, 1]$, we use the estimates

$$\begin{aligned} |f_{\gamma, \lambda}(s\lambda)| &= \frac{s|\lambda|}{(1 - s|\lambda|^2)^\gamma} \leq \frac{1}{(1 - s|\lambda|^2)^\gamma} \\ |g'(s\lambda)| &\leq \frac{\|g\|_{\mathcal{B}}}{1 - s^2|\lambda|^2} \leq \frac{\|g\|_{\mathcal{B}}}{1 - s|\lambda|^2} \end{aligned}$$

to conclude that

$$\begin{aligned} |T_g f_{\gamma, \lambda}(t\lambda)| &\leq t|\lambda| \int_0^1 |f_{\gamma, \lambda}(st\lambda)| |g'(st\lambda)| ds \\ &\leq \|g\|_{\mathcal{B}} \int_0^1 \frac{t|\lambda| ds}{(1 - st|\lambda|^2)^{\gamma+1}} \leq \frac{\|g\|_{\mathcal{B}}}{|\lambda| \gamma (1 - t|\lambda|^2)^\gamma}. \end{aligned}$$

If the statement holds for some $k \geq 1$ and all $t \in [0, 1]$, then, as above,

$$\begin{aligned} |T_g^{k+1} f_{\gamma, \lambda}(t\lambda)| &\leq t|\lambda| \int_0^1 |T_g^k f_{\gamma, \lambda}(st\lambda)| |g'(st\lambda)| ds \\ &\leq \frac{\|g\|_{\mathcal{B}}^{k+1}}{|\lambda|^k \gamma^k} \int_0^1 \frac{t|\lambda| ds}{(1 - st|\lambda|^2)^{\gamma+1}} \leq \frac{\|g\|_{\mathcal{B}}^{k+1}}{|\lambda|^{k+1} \gamma^{k+1} (1 - t|\lambda|^2)^\gamma}, \end{aligned}$$

and the result follows. Finally, b) is a straightforward application of a). \square

Proof of Theorem 1.2 c). If $g^{n+1} \in \mathcal{B}$, then $g^k \in \mathcal{B}$, for $1 \leq k \leq n+1$, by Proposition 2.1, and the boundedness of L_g follows from the identity $S_g^k T_g = \frac{1}{k+1} T_{g^{k+1}}$ and Theorem 2.3 a).

Conversely, assume without loss of generality that $P_n(0) = n+1$. If $L_g \in \mathcal{B}(A_\alpha^p)$ then $g \in \mathcal{B}$, by Theorem 1.1 a). Moreover, Proposition 4.3 shows that $L_{g_r} \in \mathcal{B}(A_\alpha^p)$ and $\|L_{g_r}\|_{\alpha, p} \leq \|L_g\|_{\alpha, p}$, for any $r \in (0, 1)$. Now we write $P_n(z) = (n+1)(1 + zQ_n(z))$, and using again the identity $S_g^k T_g = \frac{1}{k+1} T_{g^{k+1}}$ we obtain that

$$L_{g_r} = T_{g_r^{n+1}} + T_{g_r^{n+1}} T_{g_r} Q_n(T_{g_r}) + \sum_{k=0}^{n-1} \frac{1}{k+1} T_{g_r^{k+1}} P_k(T_{g_r}) + g_r(0) Q(g_r - g_r(0)) \delta_0.$$

For $\gamma > \frac{\alpha+2}{p}$ and $\lambda \in \mathbb{D} \setminus \{0\}$, we apply L_{g_r} to the function $f_{\gamma, \lambda}$ from Lemma 4.11. Since $\delta_0(f_{\gamma, \lambda}) = 0$, we obtain that

$$L_{g_r} f_{\gamma, \lambda} = T_{g_r^{n+1}} f_{\gamma, \lambda} + T_{g_r^{n+1}} T_{g_r} Q_n(T_{g_r}) f_{\gamma, \lambda} + \sum_{k=0}^{n-1} \frac{1}{k+1} T_{g_r^{k+1}} P_k(T_{g_r}) f_{\gamma, \lambda}.$$

Now we use the standard estimates

$$|h'(\lambda)| \leq \frac{c_\alpha \|h\|_{\alpha, p}}{(1 - |\lambda|^2)^{\frac{\alpha+2}{p} + 1}}, \quad \|f_{\gamma, \lambda}\|_{\alpha, p} \leq \frac{c_{\alpha, \gamma}}{(1 - |\lambda|^2)^{\gamma - \frac{\alpha+2}{p}}},$$

where $c_\alpha > 0$ and $c_{\alpha,\gamma} > 0$ are constants which depend only on α , and α and γ , respectively, and Proposition 4.3 to infer that

$$|(L_{g_r} f_{\gamma,\lambda})'(\lambda)| \leq \frac{c_\alpha c_{\alpha,\gamma} \|L_g\|_{\alpha,p}}{(1-|\lambda|^2)^{\gamma+1}}.$$

Since

$$\begin{aligned} (L_{g_r} f_{\gamma,\lambda})'(\lambda) &= (g_r^{n+1})'(\lambda) f_{\gamma,\alpha}(\lambda) + (g_r^{n+1})'(\lambda) [T_{g_r} Q_n(T_{g_r}) f_{\gamma,\alpha}](\lambda) \\ &\quad + \sum_{k=0}^{n-1} \frac{(g_r^{k+1})'(\lambda)}{k+1} [P_k(T_{g_r}) f_{\gamma,\lambda}](\lambda), \end{aligned}$$

by the triangle inequality we have

$$(4.14) \quad \frac{|\lambda| |(g_r^{n+1})'(\lambda)|}{(1-|\lambda|^2)^\gamma} \leq |(g_r^{n+1})'(\lambda)| |[T_{g_r} Q_n(T_{g_r}) f_{\gamma,\lambda}](\lambda)| + \frac{c_\alpha c_{\alpha,\gamma} \|L_g\|_{\alpha,p}}{(1-|\lambda|^2)^{\gamma+1}} \\ + \sum_{k=0}^{n-1} \frac{|(g_r^{k+1})'(\lambda)|}{k+1} |[P_k(T_{g_r}) f_{\gamma,\lambda}](\lambda)|.$$

We want to estimate the terms on the right with the help of Lemma 4.11 b). To this end, note that n , Q_n , and P_k , for $0 \leq k \leq n-1$, depend only on L_g , so there exists a constant $c = c(L_g) > 0$ depending only on L_g such that, for $\gamma|\lambda| > \|g\|_{\mathcal{B}}$, we have that

$$|[T_{g_r} Q_n(T_{g_r}) f_{\gamma,\lambda}](\lambda)| \leq c \frac{\|g\|_{\mathcal{B}}}{|\lambda|^\gamma (1-|\lambda|^2)^\gamma},$$

and

$$|[P_k(T_{g_r}) f_{\gamma,\lambda}](\lambda)| \leq c \left(\frac{|\lambda|}{(1-|\lambda|^2)^\gamma} + \frac{\|g\|_{\mathcal{B}}}{|\lambda|^\gamma (1-|\lambda|^2)^\gamma} \right) \quad (0 \leq k \leq n-1).$$

Using these inequalities in (4.14) we obtain

$$(4.15) \quad |(g_r^{n+1})'(\lambda)| \leq |(g_r^{n+1})'(\lambda)| \frac{c \|g\|_{\mathcal{B}}}{|\lambda|^\gamma} + \frac{c_\alpha c_{\alpha,\gamma} \|L_g\|_{\alpha,p}}{|\lambda| (1-|\lambda|^2)} \\ + c \sum_{k=0}^{n-1} \frac{|(g_r^{k+1})'(\lambda)|}{k+1} \left(1 + \frac{\|g\|_{\mathcal{B}}}{|\lambda|^\gamma} \right),$$

when $\gamma|\lambda| > \|g\|_{\mathcal{B}}$. Now if γ satisfies $\gamma > 8(c+1)\|g\|_{\mathcal{B}}$, (4.15) gives for $|\lambda| > \frac{1}{2}$

$$\frac{1}{2} |(g_r^{n+1})'(\lambda)| \leq \frac{2c_\alpha c_{\alpha,\gamma} \|L_g\|_{\alpha,p}}{(1-|\lambda|^2)} + \frac{3c}{2} \sum_{k=0}^{n-1} \frac{|(g_r^{k+1})'(\lambda)|}{k+1}.$$

Thus, we either have $\|g_r^{n+1}\|_{\mathcal{B}} = \sup_{|\lambda| \leq \frac{1}{2}} (1-|\lambda|^2) |(g_r^{n+1})'(\lambda)|$, or, by Proposition 2.1 and the last inequality,

$$\|g_r^{n+1}\|_{\mathcal{B}} \leq 4c_\alpha c_{\alpha,\gamma} \|L_g\|_{\alpha,p} + c' \sum_{k=0}^{n-1} \frac{1}{k+1} \|g_r^{n+1}\|_{\mathcal{B}}^{\frac{k+1}{n+1}}.$$

This shows that $\|g_r^{n+1}\|_{\mathcal{B}}$ stays bounded when $r \rightarrow 1^-$, i.e. $g^{n+1} \in \mathcal{B}$. \square

4.3. Compositions of two analytic paraproducs. Corollary 1.4 together with the identities

$$\begin{aligned} M_g^2 &= S_g^2 + 2S_gT_g + g^2(0)\delta_0 \\ M_gT_g &= S_gT_g + T_g^2 \\ S_gM_g &= S_gT_g + S_g^2 \\ T_gM_g &= S_gT_g \\ T_gS_g &= S_gT_g - T_g^2 - g(0)(g - g(0))\delta_0 \\ M_gS_g &= S_gT_g - T_g^2 + S_g^2 - g(0)(g - g(0))\delta_0 \end{aligned}$$

yield a complete characterization of compositions of two analytic paraproducs. A summary for $\alpha > -1$ is provided in the following table. The analogue for the H^p -case can be obtained replacing \mathcal{B} by $BMOA$.

Boundedness of composition of analytic paraproducs on A_α^p , $\alpha > -1$			
	T_g	S_g	M_g
T_g	$T_g^2 \in \mathcal{B}(A_\alpha^p) \Leftrightarrow T_g \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g \in \mathcal{B}$	$S_gT_g \in \mathcal{B}(A_\alpha^p) \Leftrightarrow T_{g^2} \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g^2 \in \mathcal{B}$	$M_gT_g \in \mathcal{B}(A_\alpha^p) \Leftrightarrow T_{g^2} \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g^2 \in \mathcal{B}$
S_g	$T_gS_g \in \mathcal{B}(A_\alpha^p) \Leftrightarrow T_{g^2} \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g^2 \in \mathcal{B}$	$S_g^2 \in \mathcal{B}(A_\alpha^p) \Leftrightarrow S_g \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g \in H^\infty$	$M_gS_g \in \mathcal{B}(A_\alpha^p) \Leftrightarrow S_g \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g \in H^\infty$
M_g	$T_gM_g \in \mathcal{B}(A_\alpha^p) \Leftrightarrow T_{g^2} \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g^2 \in \mathcal{B}$	$S_gM_g \in \mathcal{B}(A_\alpha^p) \Leftrightarrow S_g \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g \in H^\infty$	$M_g^2 \in \mathcal{B}(A_\alpha^p) \Leftrightarrow M_g \in \mathcal{B}(A_\alpha^p)$ $\Leftrightarrow g \in H^\infty$

5. PROOF OF THEOREM 1.5

The following proposition is strongly used in the proof of Theorem 1.5.

Proposition 5.1. *Let $g \in \mathcal{H}(\mathbb{D})$. Assume that g is bounded away from zero, that is, $\inf_{z \in \mathbb{D}} |g(z)| > 0$. Let h be a branch of the logarithm of g , and, for any $\beta \in \mathbb{R}$, define the β -power of g as $g^\beta := e^{\beta h}$. Then:*

- If $g \in BMOA$ ($g \in VMOA$), then $g^\beta \in BMOA$ ($g^\beta \in VMOA$, resp.), for any $\beta < 1$.*
- If $g \in BMOA$ ($g \in VMOA$), then $S_{g^\beta}T_{g^\beta}^2 \in \mathcal{B}(A_\alpha^p)$ ($S_{g^\beta}T_{g^\beta}^2 \in \mathcal{K}(A_\alpha^p)$, resp.), for any $\alpha \geq -1$, $\beta \in (0, \frac{2}{3})$, and $p > 0$.*

A key tool in the proof of Proposition 5.1 is the following simple computational lemma.

Lemma 5.2. *Let $g \in \mathcal{H}(\mathbb{D})$ be a zero free function, and, for any $\beta \in \mathbb{R}$, let g^β be as in the statement of the preceding proposition. Then*

$$(5.1) \quad S_{g^\beta}T_{g^\beta}^2 = \frac{(2\beta-1)\beta}{1-\varepsilon} T_gT_{g^{1-\varepsilon}}M_{g^{2\beta-2+\varepsilon}}T_{g^\beta} + \frac{\beta^2}{1-\varepsilon} T_gT_{g^{1-\varepsilon}}M_{g^{3\beta-2+\varepsilon}},$$

for every $\beta \in \mathbb{R}$ and $\varepsilon \in \mathbb{R} \setminus \{1\}$.

Proof. The fact that $(g^\beta)^2 = g^{2\beta}$ gives that $L := S_{g^\beta}T_{g^\beta}^2 = \frac{1}{2}T_{g^{2\beta}}T_{g^\beta}$. Thus, for any $f \in \mathcal{H}(\mathbb{D})$, we have that

$$(Lf)' = \frac{1}{2}(g^{2\beta})'T_{g^\beta}f = \beta g'F, \quad \text{where } F = g^{2\beta-1}T_{g^\beta}f.$$

Since $Lf(0) = 0$, it follows that $Lf = \beta T_g F$. Now $F(0) = 0$ and

$$\begin{aligned} F' &= (2\beta - 1) g^{2\beta-2} g' T_{g^\beta} f + \beta g^{3\beta-2} g' f \\ &= (g^{1-\varepsilon})' \left(\frac{2\beta-1}{1-\varepsilon} g^{2\beta-2+\varepsilon} T_{g^\beta} f + \frac{\beta}{1-\varepsilon} g^{3\beta-2+\varepsilon} f \right), \end{aligned}$$

for any $\varepsilon \in \mathbb{R} \setminus \{1\}$. Therefore

$$F = \frac{2\beta-1}{1-\varepsilon} T_{g^{1-\varepsilon}} M_{g^{2\beta-2+\varepsilon}} T_{g^\beta} f + \frac{\beta}{1-\varepsilon} T_{g^{1-\varepsilon}} M_{g^{3\beta-2+\varepsilon}} f,$$

and hence (5.1) holds. \square

Proof of Proposition 5.1.

a) Just observe that, since g is bounded away from zero, $g^{\beta-1}$ is bounded for $\beta < 1$, and so we have the estimate $(1 - |z|^2)|(g^\beta)'|^2 \lesssim (1 - |z|^2)|g'(z)|^2$.

b) Let $g \in BMOA$ ($g \in VMOA$), $\alpha \geq -1$, $\beta \in (0, \frac{2}{3})$, and $p > 0$. Since $\beta < \frac{2}{3}$, there is $\varepsilon \in \mathbb{R}$ such that $0 < \varepsilon < \min(2 - 3\beta, 1)$. Taking into account that $\beta > 0$, it follows that $\varepsilon \in (0, 1)$ and $2\beta - 2 + \varepsilon < 3\beta - 2 + \varepsilon < 0$. As a consequence, we have that:

- $T_g, T_{g^{1-\varepsilon}}, T_{g^\beta} \in \mathcal{B}(A_\alpha^p)$ ($T_g, T_{g^{1-\varepsilon}}, T_{g^\beta} \in \mathcal{K}(A_\alpha^p)$, resp.), by a).
- $M_{g^{2\beta-2+\varepsilon}}, M_{g^{3\beta-2+\varepsilon}} \in \mathcal{B}(A_\alpha^p)$, since $g^{2\beta-2+\varepsilon}, g^{3\beta-2+\varepsilon} \in H^\infty$, because g is bounded away from zero.

Moreover, since $\varepsilon < 1$, (5.1) holds, and we conclude that $S_{g^\beta} T_{g^\beta}^2 \in \mathcal{B}(A_\alpha)$ ($S_{g^\beta} T_{g^\beta}^2 \in \mathcal{K}(A_\alpha)$, resp.). Hence the proof is complete. \square

We also need the following auxiliary result.

Lemma 5.3. *Let $f \in C(\overline{\mathbb{D}} \setminus \{1\})$ such that*

$$(5.2) \quad \lim_{\substack{z \rightarrow 1 \\ z \in \mathbb{D}}} (1 - z) f(z) = 0.$$

Then $d\mu(z) = (1 - |z|^2)|f(z)|^2 dA(z)$ is a vanishing Carleson measure for H^p , $0 < p < \infty$.

Proof. Let $\Omega_\delta = \mathbb{D} \cap D(1, \delta)$, for every $0 < \delta < 1$. Let $a \in \mathbb{D}$ and $0 < \delta < 1$. Then

$$\int_{\mathbb{D}} \frac{1 - |a|^2}{|1 - \bar{a}z|^2} d\mu(z) = \left\{ \int_{\mathbb{D} \setminus \Omega_\delta} + \int_{\Omega_\delta} \right\} (1 - |\phi_a(z)|^2) |f(z)|^2 dA(z) = I_\delta + J_\delta.$$

Now, by [11, Proposition 1.4.10], we have that

$$(5.3) \quad \begin{aligned} I_\delta &\leq \left(\sup_{z \in \mathbb{D} \setminus \Omega_\delta} |f(z)|^2 \right) \int_{\mathbb{D}} (1 - |\phi_a(z)|^2) dA(z) \\ &\leq C_1 (1 - |a|^2) \sup_{z \in \mathbb{D} \setminus \Omega_\delta} |f(z)|^2, \end{aligned}$$

where $C_1 > 0$ is an absolute constant. Next recall that, since $\log(1 - z)$ is a function in $BMOA$ (where \log denotes the principal branch of the logarithm), $d\mu_1(z) = \frac{1 - |z|^2}{|1 - z|^2} dA(z)$ is a Carleson measure for the Hardy

spaces, and so

$$(5.4) \quad \begin{aligned} J_\delta &\leq \left(\sup_{z \in \Omega_\delta} |1 - z| |f(z)| \right)^2 \int_{\mathbb{D}} \frac{1 - |a|^2}{|1 - \bar{a}z|^2} d\mu_1(z) \\ &\leq C_2 \left(\sup_{z \in \Omega_\delta} |1 - z| |f(z)| \right)^2, \end{aligned}$$

where $C_2 > 0$ is an absolute constant. Since $f \in C(\overline{\mathbb{D}} \setminus \{1\})$, it is clear that (5.2), (5.3), and (5.4) imply that μ is a vanishing Carleson measure for the Hardy spaces, *i.e.*

$$\lim_{|a| \rightarrow 1^-} \int_{\mathbb{D}} \frac{1 - |a|^2}{|1 - \bar{a}z|^2} d\mu(z) = 0. \quad \square$$

Proof of Theorem 1.5.

a) Assume the contrary, *i.e.* $S_g T_g^2 \in \mathcal{B}(A_\alpha^p)$, for some $\alpha \geq -1$, $p > 0$. Then a standard estimate yields for every $f \in A_\alpha^p$,

$$(5.5) \quad |(S_g T_g^2 f)'(r)| \lesssim (1 - r)^{-1 - \frac{\alpha+2}{p}} \|S_g T_g^2\|_{\alpha,p} \|f\|_{\alpha,p} \quad (r \in [0, 1)).$$

The usual test functions $f_{r,k}(z) = (1 - rz)^{-k}$, for $z \in \mathbb{D}$, with $r \in (0, 1)$, $kp > \alpha + 2$, satisfy $\|f_{r,k}\|_{\alpha,p} \simeq (1 - r)^{-k + \frac{\alpha+2}{p}}$, and

$$\begin{aligned} |(S_g T_g^2 f_{r,k})'(r)| &= \frac{1}{1 - r} \log \frac{e}{1 - r} \int_0^r \frac{ds}{(1 - s)(1 - rs)^k} \\ &\geq \frac{1}{1 - r} \log \frac{e}{1 - r} \int_0^r \frac{r ds}{(1 - rs)^{k+1}} \\ &= \frac{1}{k(1 - r)} \log \frac{e}{1 - r} \left(\frac{1}{(1 - r^2)^k} - 1 \right), \end{aligned}$$

which contradicts (5.5) when $r \rightarrow 1^-$.

b) If $f \in \mathcal{B}$, then

$$|f(z)| \lesssim \log \left(\frac{e}{1 - |z|} \right) \quad (z \in \mathbb{D}),$$

but

$$\lim_{r \rightarrow 1^-} \frac{|g^{2\beta}(r)|}{\log \left(\frac{e}{1 - r} \right)} = \lim_{r \rightarrow 1^-} \left[\log \left(\frac{e}{1 - r} \right) \right]^{2\beta - 1} = \infty \quad (\beta > \frac{1}{2}),$$

and so $g^{2\beta} \notin \mathcal{B}$, for any $\beta > \frac{1}{2}$.

Now let us prove that $S_{g^\beta} T_{g^\beta}^2 \in \mathcal{K}(A_\alpha^p)$, for any $\alpha \geq -1$ and $p > 0$. We know that $g \in BMOA$. Moreover, since $z \mapsto \frac{e}{1 - z}$ maps the half-disc

$$D^- = \{z \in \mathbb{C} : |z - 1| < 1 + \frac{e}{2}, \operatorname{Re} z < 1\}$$

onto the domain $\{z \in \mathbb{C} : |z| > \frac{2e}{2+e}, \operatorname{Re} z > 0\}$, it follows that g is bounded away from zero and g^β extends analytically to D^- . In particular, $(g^\beta)' \in C(\overline{\mathbb{D}} \setminus \{1\})$ and satisfies

$$\lim_{\substack{z \rightarrow 1 \\ z \in \mathbb{D}}} (1 - z)(g^\beta)'(z) = \beta \lim_{\substack{z \rightarrow 1 \\ z \in \mathbb{D}}} g^{\beta-1}(z) = 0 \quad (\beta < 1).$$

Then Lemma 5.3 gives that $g^\beta \in VMOA$, for every $\beta < 1$, and Proposition 5.1 shows that $S_{g^\beta} T_{g^\beta}^2 \in \mathcal{K}(A_\alpha^p)$, for any $\alpha \geq -1$, $\beta \in (0, \frac{2}{3})$, and $p > 0$. \square

REFERENCES

- [1] W.W. Adams, P. Loustau, *An introduction to Gröbner bases*. Graduate Studies in Mathematics, 3. American Mathematical Society, Providence, RI, 1994.
- [2] A. Aleman, J. A. Cima, *An integral operator on H^p and Hardy's inequality*, J. Anal. Math. **85** (2001), 157–176.
- [3] A. Aleman and O. Constantin, *Spectra of integration operators on weighted Bergman spaces*, J. Anal. Math. **109** (2009), 199–231.
- [4] A. Aleman and A. Siskakis, *An integral operator on H^p* , Complex Variables Theory Appl. **28** (1995), no. 2, 149–158.
- [5] A. Aleman and A. Siskakis, *Integration operators on Bergman spaces*, Indiana Univ. Math. J. **46** (1997), 337–356.
- [6] S. Axler, *The Bergman space, the Bloch space, and commutators of multiplication operators*, Duke Math. J. **53** (1986), no. 2, 315–332.
- [7] P. L. Duren, B. W. Romberg and A. L. Shields, *Linear functionals on H^p spaces with $0 < p < 1$* , J. Reine Angew. Math. **238** (1969), 32–60.
- [8] D. Girela, *Analytic functions of bounded mean oscillation*, Complex function spaces (Mekrijärvi, 1999), 61–170, Univ. Joensuu Dept. Math. Rep. Ser., 4, Univ. Joensuu, Joensuu, 2001.
- [9] S. G. Krantz, *Harmonic and complex analysis in several variables*, Springer Monographs in Mathematics. Springer, Cham, 2017.
- [10] Ch. Pommerenke, *Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation*, Comment. Math. Helv. **52** (1977), no. 4, 591–602.
- [11] W. Rudin, *Function theory in the unit ball of \mathbb{C}^n* . Grundlehren der Mathematischen Wissenschaften, 241. Springer-Verlag, New York-Berlin, 1980.
- [12] M. Tjani, *Compact composition operators on Besov spaces*, Trans. Amer. Math. Soc. **355** (2003), no. 11, 4683–4698.
- [13] V. Ufnarovski, *Introduction to noncommutative Gröbner bases theory*. Gröbner bases and applications (Linz, 1998), London Math. Soc. Lecture Note Ser., 251, Cambridge Univ. Press, Cambridge, 1998, 259–280.
- [14] S. A. Vinogradov, *Multiplication and division in the space of analytic functions with area integrable derivative, and in some related spaces* (in Russian), Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **222** (1995), Issled. po Linein. Oper. i Teor. Funktsii **23**, 45–77, 308; translation in J. Math. Sci. (New York) **87**, no. 5 (1997), 3806–3827.
- [15] K. H. Zhu, *Operator Theory in Function Spaces*. Second Edition. Math. Surveys and Monographs, 138, American Mathematical Society, Providence, Rhode Island, 2007.

A. ALEMAN: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LUND, P.O. BOX 118, SE-221 00, LUND, SWEDEN

Email address: alexandru.aleman@math.lu.se

C. CASCANTE: DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA 585, 08071 BARCELONA, SPAIN

Email address: cascante@ub.edu

J. FÀBREGA: DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA 585, 08071 BARCELONA, SPAIN

Email address: joan-fabrega@ub.edu

D. PASCUAS: DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA 585, 08071 BARCELONA, SPAIN

Email address: daniel_pascuas@ub.edu

J. A. PELÁEZ: DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, 29071 MÁLAGA, SPAIN

Email address: japelaez@uma.es