

Sleep in youth with Autism Spectrum Disorders: systematic review and meta-analysis of subjective and objective studies

Amparo Díaz-Román ^{1*}, Junhua Zhang ^{2,3*}, Richard Delorme ^{4,5}, Anita Beggiato ^{4,5}, Samuele Cortese ^{3,6-9}

¹ Mind, Brain and Behavior Research Center, University of Granada, Granada, Spain

² School of Education, Jiangsu Key Laboratory for Big Data of Psychology and Cognitive Science, Yancheng Teachers University, 224002, Yancheng, China

³ Center for Innovation in Mental Health, Academic Unit of Psychology, University of Southampton, UK, SO17 1BJ

⁴ Unité de Génétique Humaine et Fonctions Cognitives, Département de Neurosciences, Institut Pasteur, Paris, France

⁵ Département de Psychiatrie de l'Enfant et de l'Adolescent, Hôpital Robert Debré, L'Assistance Publique-Hôpitaux de Paris, France

⁶ Clinical and Experimental Sciences (CNS and Psychiatry), Faculty of Medicine, University of Southampton, UK, SO17 1BJ

⁷ Solent NHS Trust, Southampton, UK, SO19 8BR

⁸ New York University Child Study Center, New York, NY, USA, 10016

⁹ Division of Psychiatry and Applied Psychology, School of Medicine, University of Nottingham, Nottingham, UK, NG72UH

*These authors contributed equally.

Word count (main text): 2740 words

Address correspondence to:

Dr. Amparo Díaz-Román, Sleep and Health Promotion Laboratory, Mind, Brain and Behavior Research Center, University of Granada, Granada, Spain.

E-mail: adiazroman@ugr.es

ABSTRACT

BACKGROUND: Sleep problems are common and impairing in individuals with autism spectrum disorders (ASD). Evidence synthesis including both subjective (i.e., measured with questionnaires) and objective (i.e., quantified with neurophysiologic tools) sleep alterations in youth with ASD is currently lacking. **OBJECTIVE:** To conduct a systematic review and meta-analysis of subjective and objective studies sleep studies in youth with ASD.

METHODS: We searched the following electronic databases with no language, date, or type of document restriction, up to May 23rd, 2018: Pubmed, PsycInfo, Embase+Embase Classic, Ovid Medline, and Web of Knowledge. Random-effects models were used. Heterogeneity was assessed with Cochran's Q and I² statistics. Publication (small studies) bias was assessed with funnel plots and the Egger's test. Study quality was evaluated with the Newcastle Ottawa Scale. Analyses were conducted using *Review Manager* and *Comprehensive meta-analysis*.

FINDINGS: From a pool of 3,359 non-duplicate potentially relevant references, 47 datasets were included in the meta-analyses. Subjective and objective sleep outcome measures were extracted from 37 and 15 studies, respectively. Only five studies were based on comorbidity free, medication-naïve participants. Compared to typically developing controls, youth with ASD significantly differed in 10/14 subjective parameters and in 7/14 objective sleep parameters. The average quality score in the Newcastle-Ottawa scale was 5.9/9.

CONCLUSIONS: A number of subjective and, to a less extent, objective sleep alterations might characterise youth with ASD, but future studies should assess the impact of pharmacological treatment and psychiatric comorbidities.

Summary box

What is already known about this subject?

- Autism spectrum disorders (ASD) represent common and impairing neurodevelopmental conditions.
- Sleep problems are commonly reported in individuals with ASD and have a negative impact on their daily functioning.
- Although a previous meta-analysis explored objective (i.e., measured with physiological tools) sleep problems in children with ASD, updated evidence synthesis including both subjective (i.e., assessed via questionnaires) and objective sleep problem is lacking.

What are the new findings?

- Our meta-analysis of 47 datasets showed that, compared to typically developing controls, youth with ASD significantly differed in 10/14 subjective parameters and in 7/14 objective sleep parameters.
- Comorbid psychiatric conditions and the pharmacological treatment may contribute to sleep impairment in ASD.
- Our findings allow a fine-grain characterisation of sleep alterations in ASD which is helpful for their daily clinical management.

How might it impact on clinical practice in the foreseeable future?

- Presence of sleep alterations in youths with ASD should be systematically screened in the clinical practice in order to reduce their impact on daytime functioning.

BACKGROUND

Autism spectrum disorders encompass a wide range of neurodevelopmental conditions characterized by a deficit in social communication, together with restricted, repetitive and stereotyped behaviours, interests, or activities.¹ Although not formally part of the diagnostic criteria,^{1,2} sleep problems are frequently reported in individuals with ASD (e.g.,³⁻⁵) and contribute to their functional impairment. Sleep difficulties are associated with a significant amount of distress for the patients and their families,⁶ but also negatively impact on cognitive abilities and self-regulation of disruptive behaviours during the daytime.⁷⁻⁹

In order to appropriately manage them, it is necessary to characterise the profile of sleep problems in children and adolescents with ASD. Whilst a number of individual studies have been conducted, we are aware of only one meta-analysis that summarised the available body of evidence.¹⁰ However, this meta-analysis was limited to objective sleep studies, i.e., studies relying on actigraphic or polysomnographic measures. Whilst these (in particular, polysomnography) are considered rigorous measures of sleep, it is important to also consider sleep measures subjectively reported by patient and/or their parents via questionnaires, as they are arguably more “ecological” and they reflect the subjective perception, which is important in the management process of the disorder. Furthermore, the meta-analysis by Elrod and Hood¹⁰ was published in 2015 and, as such, an update is warranted.

OBJECTIVE

To conduct a systematic review and meta-analysis of subjective and objective studies of sleep in children and adolescents with ASD compared to typically developing controls.

METHODS

We followed the recommendations of the Meta-Analysis of Observational Studies in Epidemiology group ¹¹ and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. ¹² The protocol of this systematic review was registered in PROSPERO (CRD42018100016).

Type of studies

We included case-control studies comparing children with ASD to typically developing individuals on subjective and/or objective sleep parameters.

Type of participants

We included studies on children/youth (≤ 20 years) diagnosed with ASD according to DSM III to 5 criteria or ICD-9 to 10 criteria, or according to a clinical diagnosis of ASD, compared to typically developing participants. Definition of ASD based on cut-off on questionnaires targeting ASD symptoms was not considered rigorous and as such was exclusionary. Psychiatric comorbidities were not an exclusionary criterion.

Outcomes

Any subjective sleep parameters from any sleep questionnaire and/or any objective sleep parameters measured using polysomnography (PSG), actigraphy or multiple sleep latency test (MSLT), which were presented in at least two studies, were meta-analysed. We selected the following subjective parameters: bedtime resistance, sleep onset delay, sleep duration, sleep anxiety, night awakenings, parasomnias, sleep-disordered breathing, daytime sleepiness, general sleep problems, sleep quality, sleep efficiency, sleep onset latency (min), sleep duration (min), and restorative value of sleep (*i.e.*, feeling well-rested after waking up).

For PSG, we considered total sleep time, sleep onset latency, time spent in each sleep stage, REM latency, sleep efficiency, and wake time after sleep onset. As for actigraphic parameters, we selected: sleep onset latency, true sleep, assumed sleep time, actual wake time, and sleep efficiency. For MST, we considered latency to falling asleep.

Search strategy/syntax

We searched the following electronic databases: Pubmed (Medline), OVID databases (PsycInfo, EMBASE+EMBASE classic, OVID Medline), and WEB OF KNOWLEDGE Databases (Web of science (Science Citation Index Expanded), Biological abstracts, Biosis, Food science and technology abstracts), up to May 23rd, 2018 with no language/date/type of document restrictions. Further details on the search strategy/syntax, including search terms for each database, are reported in the Supplemental Material 1. References of included studies and of reviews conducted on this topic were also hand-searched to find potential pertinent studies undetected with the electronic search strategy.

Screening and data extraction

Screening

Title and abstracts of all non-duplicated papers were independently screened by two of the authors (JZ, AD). Potential pertinent papers were retained and assessed for eligibility by screening the full-text. A third senior author (SC) acted as arbitrator when disagreement in any screening stage. If needed, corresponding authors of retained studies were also contacted to request further information.

Data extraction

Data extraction was independently performed by two of the authors (JZ, AD), and any discrepancy between them was resolved by consensus. The following data were extracted from each study: first author and publication year, country where the study was conducted, study participants' details (number, percentage of males, mean age and SD, ASD diagnostic criteria, medication status, and comorbidities), mean and SD for each outcome measure (subjective and/or objective sleep parameters), and nights recorded for sleep assessment.

Risk of bias assessment

Two authors (JZ, AD) independently assessed the methodological quality or risk of bias of included studies using the Newcastle-Ottawa Scale for case-control studies (http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp). This scale includes the following domains: case definition, representativeness of the cases, selection of controls, definition of controls, comparability of cases and controls on the basis of the design or analysis, ascertainment of exposure, and non-response rate. Disagreements between both authors were resolved by consensus.

Statistical analysis

Analyses were performed with Review Manager 5.3 (<http://community.cochrane.org/tools/reviewproduction-tools/revman-5>) and Comprehensive Meta-Analysis, CMA (<http://www.meta-analysis.com/index.php>). Random-effects models were used to compute standardized mean difference (SMD) for each sleep parameter, with 95% confidence interval and the Hedges' correction¹³ to avoid sample size bias. The inverse variance method and the Z statistic were used to calculate the pooled SMD and assess its statistical significance. Heterogeneity degree between studies was measured with Cochran's Q and I² statistics.¹⁴ Publication bias were explored using the Egger's test and the funnel

plots.¹⁵ We also conducted a post hoc analysis including only studies based on comorbidity free, medication-naïve participants.

FINDINGS

From a pool of 3,359 non-duplicate potentially relevant references, 47 datasets (reported in 48 references) were included in our meta-analysis^{5,6,22–31,7,32–41,8,42–51,16,52–56,17–21} (Figure 1). The list of excluded reports (with reasons for exclusions) and included studies are provided in the Supplemental Materials 2 and 3, respectively. Table 1 shows the main characteristics of the studies included in the meta-analysis. All studies were cross-sectional and the average quality score in the Newcastle-Ottawa scale was 5.9/9 (scores ranged from 3 to 8; Supplemental Material 4).

Subjective outcome measures were extracted from 37 studies,^{5–8,16–18,21,22,24,25,27–37,39,40,42–47,49–54,56} whilst objective outcome measures were obtained from 15 studies (eight studies using PSG,^{20,25,26,29,32,37,44,53} six using actigraphy,^{18,19,27,47,50,52} and one using both⁴¹). Overall, the number of participants ranged from 75 to 5,430 for studies reporting subjective sleep parameters, and from 144 to 312 for sleep objective studies. Two studies^{35,38} reported sleep data of two different samples and we included both samples in the meta-analysis independently.

Subjective measures of sleep difficulties

Compared to control individuals, participants with ASD, showed significantly higher bedtime resistance (SMD = 1.00, 95% CI: 0.67 to 1.33), sleep onset delay (0.98, 0.66 to 1.29), sleep anxiety (0.96, 0.61 to 1.32), night awakenings (0.72, 0.44 to 1.01), parasomnias (0.88, 0.60 to 1.15), sleep-disordered breathing (0.48, 0.28 to 0.67), daytime sleepiness (0.34, 0.16 to 0.52), sleep onset latency (in min) (0.81, 0.59 to 1.02), restorative value of sleep

(0.81, 0.59 to 1.02), and general sleep problems (0.93, 0.67 to 1.20). They also showed lower sleep duration (-0.88, -1.18 to -0.57). In contrast, children with ASD did not significantly differ from control individuals in sleep quality, sleep efficiency, or sleep duration in min (Table 2 and the Supplemental Material 5). As shown in Table 2, the heterogeneity between studies was statistically significant for almost all subjective sleep parameters (I^2 ranged from 81% to 95%), except for sleep efficiency and sleep onset latency (in min). There was also evidence for publication bias for 5 out of 14 subjective sleep parameters: sleep duration ($t = 2.19$, $p = 0.040$), sleep anxiety ($t = 2.69$, $p = 0.014$), parasomnias ($t = 3.30$, $p = 0.003$), daytime sleepiness ($t = 2.26$, $p = 0.032$) and general sleep problems ($t = 2.31$, $p = 0.028$). The results of the Egger's test and the funnel plots are reported in Table 2 and the Supplemental Material 6, respectively.

Objective parameters of sleep alterations

As reported in Table 3, children with ASD significantly differed from control individuals in several objective parameters measuring sleep patterns using PSG. Specifically, children with ASD showed lower total sleep time (-0.90, -1.51 to -0.30), longer sleep onset latency (0.53, 0.21 to 0.86), higher time spent in stage 1 sleep (0.48, 0.06 to 0.90), lower time of REM sleep (-0.88, -1.56 to -0.21), lower sleep efficiency (-1.20, -1.98 to -0.41), and higher time awake after sleep onset (0.49, 0.11 to 0.87). However, no significant differences were observed between children with ASD and control individuals in stage 2 sleep, slow wave sleep, and REM latency (Table 3 and the Supplemental Material 5). In relation to actigraphy, we found differences between both groups only in sleep onset latency (Table 3 and the Supplemental Material 5). Children with ASD displayed significantly longer sleep onset latency than control individuals (0.80, 0.55 to 1.05). Evidence of heterogeneity was found for almost all polysomnographic sleep parameters (I^2 ranged from 55% to 85%), with the

exception of sleep onset latency and wake time, but only for a single actigraphic sleep parameter (sleep efficiency, $I^2 = 62\%$). No evidence of publication bias was detected in the Egger's test (Table 3) and the funnel plots (Supplemental Material 6).

Post hoc analysis

The post hoc analysis based on studies including only comorbidity-free and medication naïve participants was limited to PSG studies as only two studies for subjective measures and one study for actigraphic measures, respectively, provided usable data. As shown in Table 4 the post hoc analysis of PSG studies replicated the results of the main analysis (excepted for the parameter duration of sleep stage 1, that was not more significant between participants with ASD and controls).

DISCUSSION

To our knowledge, this is the first meta-analysis including both subjective and objective measures of sleep in children with ASD. We found that, compared to typically developing children, those with ASD presented with a number of significant sleep impairments, quantified both by subjective and objective parameters.

Our results were in accordance with the findings of a previous meta-analysis in children with ASD,¹⁰ in which these children also showed significantly lower total sleep time, increased sleep onset latency, and worse sleep efficiency compared to typically developing children. However, these differences between groups observed during PSG were not consistent with actigraphy-defined measures since only sleep time or sleep efficiency statistically differed.

It should be noted that, although the previous meta-analysis by Elrod and Hood¹⁰ pooled both PSG and actigraphy sleep outcomes together, their analyses of moderating

factors revealed a significant impact of sleep assessment method on sleep efficiency. Specifically, their results suggested no difference in children with ASD and controls in actigraphic sleep efficiency, being this consistent with our results to a greater extent.

Our work adds meta-analytic evidence to the Elrod and Hood study,¹⁰ extending to subjective measures of sleep disturbances in ASD. Our results stress further that children with ASD displayed a considerable burden of sleep problems. These children seemed to experience greater bedtime resistance, sleep anxiety, sleep-disordered breathing, and parasomnias, as well as longer sleep onset latency and higher daytime sleepiness. However, there was less consistency about total sleep time, depending on how it was estimated (i.e., with a score in a sleep questionnaire or a length in min). Finally, despite children with ASD showed significantly higher scores in general sleep problems than typically developing children, they did not differ in terms of subjectively reported sleep quality and sleep efficiency, although the limited number of included subjective studies reporting sleep quality (n = 3) and sleep efficiency (n = 2) suggest that this conclusion should be considered with caution.

Our results based on subjective measures were generally not consistent with those obtained with objective parameters. For instance, children with ASD and control individuals did not significantly differ in terms of sleep duration based on parents' report. By contrast, children with ASD showed a significantly lower total sleep time compared to typically developing children according to PSG measures. This is not surprising and reflects the well-known mismatch between subjective and objective measures.⁵⁷ Indeed, discrepancies between subjective and objective sleep measures have been reported in earlier studies in both children with ASD (e.g.,^{18,41}) and children with other neurodevelopmental disorders (e.g.,^{58,59}). For example, objective measures were usually taken on one or two nights, while subjective measures reflected the perception of the parent over several nights. Taking into

account the advantages and limitations of both subjective and objective sleep measures, rather than considering these two types of measures as exclusionary, we would suggest they should be seen as providing complementary information. Additionally, even within objective studies, there were some discrepancies among apparently similar parameters. Of note, sleep efficiency was significantly different between participants with and without ASD when measured with PSG but not when assessed via actigraphy. The different degree of ecological validity of PSG (usually implemented in a lab) and actigraphy (in the home environment) may contribute to explain these discrepancies.

Our findings could have been impacted by the presence of psychiatric comorbidities and the drug intake of the subjects included in our meta-analysis. For example, psychiatric comorbidities, including epilepsy and Attention-Deficit/Hyperactive Disorder (AHDH) were reported in at least 19% (7/37) of the studies included in our meta-analysis. Drug intake, including stimulants and melatonin, was mentioned in 35% (13/37) of the studies included. In fact, the impact of psychiatric comorbidities on sleep has been consistently reported in research.^{60,61} The effects of drug on sleep patterns were also well-documented.⁶¹ Thus, the high heterogeneity of the results among studies we observed (based on the results of the Egger's test and the funnel plot) may be related to participants' comorbid conditions or medication intake. Unfortunately, most of the studied taking account in our analysis did not provide this information which could have been useful to perform a meta-regression to assess the impact of these possible confounders. Additionally, our post hoc analysis based on studies including only comorbidity free and medication naïve participants could only confirm the results of the main meta-analysis of PSG studies, as there were not enough studies for the analysis of subjective and actigraphic parameters.

In addition to the possible role of psychiatric comorbidities and medications, the causes of sleep impairments in children with ASD are likely to be complex and not mutually

exclusive. Behavioral factors such as dysfunctional bedtime routines, exacerbated by comorbid anxiety or ADHD, may disrupt sleep, especially sleep onset delay. There is also an increasing body of evidences suggesting the contributing role of biological clock factors (mainly endocrine and genetic) which could be involved in dysregulation of day-night rhythm and sleep patterns in subjects with ASD (see review of⁶²). For instance, ASD was associated with decreased urinary or blood melatonin level,⁶³ probably due to genetic and epigenetic abnormalities affecting the enzymes of the melatonin synthesis and degradation pathways.⁶⁴ Similarly, several studies suggested that BMAL1 or additional clock genes involved in the synchronisation of biological rhythms, may be impaired in ASD. This may affect the ability of patients with ASD to anticipate and adapt their behaviours (including their sleep patterns) to environmental changes.⁶⁵

The results of our systematic meta-analysis should be considered in the light of its strengths and limitations. As for the strengths, we pre-registered the protocol in a publicly available repository (PROSPERO), reducing the risk of reporting bias. Furthermore, we endeavoured to perform a comprehensive and systematic search in several databases, with no restrictions in terms of language or document type, and we gathered unpublished data from study authors. Additionally, we used a state-of-the-art tool, the Newcastle-Ottawa scale, to assess the quality of the retained studies.

There were also a number of limitations that should be taken into account. First, statistical heterogeneity was significant for the majority of the included measures. Although this did not invalidate the results, it indicated that the pooled effect sizes could not appropriately summarise the results from all datasets. Second, whilst we endeavoured to perform a comprehensive search, there was evidence of publication bias for a number of measures, which suggested that a more transparent report of research findings in the field is needed. Third, as for the quality of individual studies, most of them were rated at overall

medium quality or risk of bias using the Newcastle-Ottawa scale, and main concerns were noted in relation to the comparability of groups and exposure-related items. The latter, along with the relatively sparse evidences available for some sleep parameters, calls for more studies assessing sleep impairments in children with ASD and by controlling the mentioned concerns.

Despite these caveats, we deem that our study provides meta-analytic evidence of objective and subjective sleep difficulties in patients with ASD. Clinicians managing children with ASD should systematically query about sleep alterations, during the first assessment and all along the follow-up. Subjective questionnaires such as the scale by Bruni et al.⁶⁶ or Owens et al.⁶⁷ can be used to screen sleep difficulties at the first assessment and at each follow-up visit with children with ASD. The extent to which these alterations are accounted for by comorbid disorders and/or the effect of pharmacotherapy should be better explored in future studies recruiting only medication naïve and comorbidity free participants, although our post hoc analysis suggest that objective differences inn sleep parameters are detected regardless the effect of comorbidities and medications. Additionally, further research needs to be performed to dissect the dysfunction of biological regulators in ASD. This may offer new promising avenues for early detection and therapeutic intervention in ASD.

Table 1. Descriptive table of the studies included in the meta-analysis.

First author (year)	Country	ASD group			Control group			Co-occurrent drugs	Comorbidities	Main sleep measures	Nights recorded (PSG/ACT)
		Diagnosis	n (% male)	Age (years)	Type	n (% male)	Age				
Aathira (2017)	India	DSM-IV	71 (80.28)	5.3±1.8	TD	65 (61.54)	5.7±1.6	22 psychotropic medication, 11 antiepileptics	11 children with ASD had epilepsy	CSHQ	NA
Al-Farsi (2018)	Oman	DSM-V-TR	122 (82.8)	3-14	TD	90 (64)	3-14	NS	NS	CSHQ	NA
Allik (2006) ^a	Sweden	ICD-10	32 (87.5)	10.80±1.25	TD	32 (87.5)	10.9 ± 1.3	0	0	ACT, Other subjective	7
Baker (2013)	Australia	Clinical diagnosis	27 (81.48)	15.5 ± 1.3	TD	27 (81.48)	15.5 ± 1.1	12 ASD (including anti-depressants, psycho-stimulants, sleeping agents, anti-acne treatments, the contraceptive pill, antipsychotics and anti-inflammatories) and 2 TD (anti-acne medication and thyroxine)	NS	ACT, Other subjective	7
Bruni (2007)	Italy	ICD-10 and DSM-IV	18 (88.89)	12.26±2.50	TD	12 (58.33)	12.6±3.7	0	0	PSG	2
Chou (2012)	China	DSM-IV	110 (87.3)	7.6 ± 2.4	TD	110 (87.3)	7.9 ± 2.0	NS	NS	Other subjective	NA
Cotton (2006) ^a	Australia	Clinical diagnosis	37 (75.7)	7.09 ± 2.49	TD	55 (58.2)	8.88 ± 3.95	NS	NS	Other subjective	NA
Cotton (2010) ^a	Australia	Clinical diagnosis	34 (73.53)	7.2 ± 2.5	TD	33 (69.70)	8.2 ± 3.2	NS	NS	Other subjective	NA
Couturier (2005)	Canada	DSM-IV	23 (95.7)	9.4 ± 2.0	TD	23 (95.7)	9.5± 2.0	75% ASD (56% stimulants, antipsychotics, and antidepressants, and 12.9% others)	NS	CSHQ	NA
Elia (1991)	Italy	DSM-III-R	4 (100)	13.3 (10.5-15)	TD	5 (100)	13.0 (9-17)	NS	0	PSG	3

Elia (2000)	Italy	DSM-IV	17 (100)	10.36 ± 3.79	TD	5 (100)	9.22 ± 2.02	0	0	PSG	2
Fletcher (2017)	Australia	DSM-IV-TR	21 (81.0)	106.67 ± 26.82 months	TD	29 (48.3)	102.10 ± 17.07 months	ASD group: 3 melatonin, 2 stimulant, 2 SSRI, 1 atomoxetine	NS	ACT, CSHQ	14
Giannotti (2008) ^a	Italy	DSM-IV-TR	104 (86.54)	2.3-7.10	TD	162 (62)	2.2-7.11	0	19.44% epilepsy (ASD group)	CSHQ	NA
Giannotti (2011) ^a	Italy	DSM-IV-TR	40 (77.5)	5.32±3.12	TD	12 (75)	5.8 ± 2.4	0	0	PSG	2
Guler (2016)	Turkey	DSM-5	60 (73.3)	7.10 ± 1.50	TD	60 (65)	6.93 ± 1.59	NS	NS	CSHQ	NA
Han (2017)	China	DSM-5	212 (85.4)	6.0 ± 2.7	TD	334 (81.4)	5.9 ± 2.6	NS	0	CSHQ	NA
Harder (2016) ^a	USA	Clinical diagnosis	21 (100)	7.8 ± 1.8	TD	23 (78.26)	8.0 ± 1.9	0	0	PSG	1
Henderson (2011)	USA	Clinical diagnosis	58 (86.2)	9.0 ± 2.09	TD	57 (52.6)	8.25 ± 1.98	NS	NS	Other subjective	NA
Hirata (2016)	Japan	DSM-5	193 (80.83)	4.45 ± 1.24	Community group	965 (80.83)	4.51 ± 1.15	NS	0	Other subjective	NA
Hodge (2014) ^a	USA	DSM-IV-TR (questionnaire)	108 (83.33)	7.33 ± 3.18	TD	108 (83.33)	7.71 ± 3.13	NS	NS	CSHQ	NA
Hoffman (2006) ^a	USA	DSM-IV-TR	106 (84)	8.2 ± 2.69	TD	168 (55)	8.62 ± 3.28	NS	ASD group: 14 seizure disorders, 15 ADHD, 6 cerebral palsy	CSHQ	NA
Inamuna (1984) (1) ^b	Japan	ICD-9 (modified version)	11 (-)		TD	16 (-)	NS	NS	NS	Other subjective	NA
Inamuna (1984) (2) ^b	Japan	ICD-9 (modified version)	19/20 (-)		TD	17/18 (-)	NS	NS	NS	Other subjective	NA
Kelmanson (2018)	Russia	DSM-5 (questionnaire)	18 (100)	5	TD	54 (100)	5	NS	NS	CSHQ	NA
Kheirouri (2016)	Iran	DSM-IV-TR	35 (68.6)	8.1 ± 4.0	TD	31 (58.1)	7.3 ± 2.6	97.1 ASD	0	CSHQ	NA

Lambert (2016) ^a	Canada	DSM-IV	11 (NS)	10.27 ± 2.24	TD	13 (NS)	10.23 ± 2.01	1 ASD (methylphenidate)	0	PSG, CSHQ, Other subjective	2
Levin (2016)	Israel	DSM-IV	34 (73.5)	3.28±0.43	TD	31 (48.4)	3.02±0.48	NS	0	CSHQ	NA
Li (2012) (1) ^b	China	DSM-IV	49 (-)	4.7 ± 0.7	TD	49 (-)	4.5 ± 0.9	NS	NS	CSHQ	NA
Li (2012) (2) ^b	China	DSM-IV	35 (-)	9.0 ± 2.0	TD	42 (-)	8.3 ± 1.8	NS	NS	CSHQ	NA
Lopez-Wagner (2008) ^a	USA	DSM-IV-TR	106 (84)	8.2 ± 2.69	TD	168 (55)	8.62 ± 3.28	NS	ASD group: 14 seizure disorders, 15 ADHD, 6 cerebral palsy	CSHQ	NA
Malow (2009) ^a	USA	Clinical diagnosis	93 (90)	5.7 ± 2.1	TD	64 (59)	6.8 ± 2.2	13% ASD and 5% TD (including atomoxetine, benadryl, citralopram, clonidine, dexmethylphenidate, divalproex sodium, fluoxetine, guanfacine, melatonin, methylphenidate, oxycarbamazapine, risperidone, and sertraline)	0	CSHQ	NA
Maski (2015)	USA	Clinical diagnosis	22 (86)	11.3 ± 2.1	TD	20 (90)	12.3 ± 2.1	31.8 % ASD (3 stimulants, 2 antidepressants, 1 guanfacine ± antidepressant, 1 mood stabilizer ± antidepressant)	0	PSG, ACT, CSHQ	1 PSG/7 ACT
Matsuoka (2014)	Japan	DSM-IV-TR	31 (93.55)	6-12	TD	372 (48.9)	9.4 ± 4.5	Anti-allergics, antipsychotics, anti-epileptics, methylphenidate (exact percentage for	2 TD had epilepsy, ASD no reported	CSHQ	NA

								ASD group is not reported)			
May (2015)	Australia	DSM-IV-TR	46 (52.17)	9.84±1.89	TD	38 (63.16)	9.04±1.62	7 ASD (including serotonin-specific reuptake inhibitors, stimulants, risperidone, and melatonin)	NS	CSHQ	NA
Miano (2007)	Italy	DSM-IV	16 (100)	9.4 ± 2.33	TD	18 (50)	10.2 ± 2.93	0	0	PSG	2
Mutluer (2016)	Turkey	DSM-5	64 (79.69)	11.66 ± 3.8	TD	53 (75.47)	11.75 ± 0.85	0	0	Other subjective	NA
Paavonen (2008)	Finland	DSM-IV and ICD-10	52 (76.9)	10.1 ± 3.4	TD	61 (47.5)	10.0 ± 1.9	0	NS	Other subjective	NA
Pace (2016)	France	DSM-5	19 (NS)	10.7 ± 1.2	TD	19 (NS)	9.9 ± 1.6	0	0	ACT	7
Park (2012)	Korea	DSM-IV-R	166 (87.3)	7.49 ± 3.05	TD (siblings)	111 (47.7)	7.94 ± 3.50	NS	0	Other subjective	NA
Patzold (1998) ^a	Australia	DSM-III and DSM-III-R	38 (81.58)	7.79±2.63	Some with developmental disabilities	36 (80.56)	8.42±2.58	45% ASD (antipsychotics and over-the-counter medications), 20% TD (anticonvulsants and asthma medications)	NS	Other subjective	NA
Phung (2017)	USA	Clinical diagnosis	19 (84.2)	16.88 ± 2.50	Neurotypical adolescents	10 (60)	15.73 ± 2.00	NS	NS	ACT, Other subjective	7
Richdale (1995)	Australia	DSM-III and DSM-III-R	12 (50)	9.12±4.99	Non-ASD	35 (57.14)	7.33±2.61	NS	2 ASD had epilepsy, 4 TD were asthmatic, 1 had allergy, 1 both	Other subjective	NA
Souders (2009)	USA	DSM-IV-TR (checklist)	59 (81.4)	7.53 ± 1.92	TD	40 (65)	7.09 ± 2.09	56.7% ASD (15 melatonin, 3 catapres, 4 risperidone, 2 aripipazole, 1 hydroxyzine, 1 fluoxetine)	0	ACT, CSHQ	10

Tessier (2015) ^a	Canada	DSM-IV-TR	13 (100)	10.23 ± 2.08	TD	13 (100)	10.23 ± 2.0	0	0	PSG	2
Tzischinsky (2018)	Israel	DSM-5	69 (81.16)	4.94 ± 1.23	TD	62 (66.13)	4.82 ± 1.15	22 ASD (including melatonin, risperdal, ritalin, and neuleptil)	NS	CSHQ	NA
van der Heijden (2018)	Netherlands	DSM-IV	68 (89.7)	9.6 ± 1.9	TD	243 (51.9)	8.7 ± 2.1	NS	ASD: 16 ADHD, 7 other	Other subjective	NA
Yang (2018)	China	DSM-IV	169 (85.80)	5.23 ± 2.0	TD	172 (84.88)	5.29 ± 1.58	NS	NS	CSHQ	NA

PSG = polysomnography; ACT = actigraphy; TD = typically development children; NA = not applicable; NS = not specified.

^aThe following studies presented some overlapped participants: 1) Allik 2006, Allik 2006, Allik 2008; 2) Cotton 2006, Cotton 2010, Patzold 1998; 3) Giannotti 2008, Giannotti 2011; 4) Harder 2016, Malow 2009; 5) Hodge 2014, Hodge 2013; 6) Hoffman 2006, Lopez-Wagner 2008; and 7) Lambert 2016, Tessier 2015. Studies reported in this table are those studies from which data for meta-analysis were extracted.

^bNumbers 1 and 2 within parenthesis denote two different samples were reported in studies.

Table 2. Summary of the results of the meta-analysis with subjective sleep parameters.

Sleep parameter	k	N	SMD (95% CI)	Z	p	Heterogeneity			Egger's test	
						Q	p	I ²	t	p
Bedtime resistance	21	3589	1.00 [0.67, 1.33]	5.98	<0.00001	362.85	<0.00001	94	2.00	0.059
Sleep onset delay	22	3636	0.98 [0.66, 1.29]	6.03	<0.00001	362.72	<0.00001	94	2.09	0.049
Sleep duration	22	3376	-0.88 [-1.18, -0.57]	5.69	<0.00001	304.14	<0.00001	93	2.19	0.040
Sleep anxiety	20	3312	0.96 [0.61, 1.32]	5.30	<0.00001	375.97	<0.00001	95	2.69	0.014
Night awakenings ^a	20	3312	0.74 [0.44, 1.04]	4.81	<0.00001	274.60	<0.00001	93	1.26	0.222
Parasomnias	22	4747	0.88 [0.60, 1.15]	6.24	<0.00001	330.97	<0.00001	94	3.30	0.003
Sleep-disordered breathing	24	4129	0.48 [0.28, 0.67]	4.80	<0.00001	181.14	<0.00001	87	1.06	0.299
Daytime sleepiness	28	5430	0.34 [0.16, 0.52]	3.62	0.0003	226.25	<0.00001	88	2.26	0.032
General sleep problems	27	5291	0.93 [0.67, 1.20]	6.97	<0.00001	409.23	<0.00001	94	2.31	0.028
Sleep quality	3	155	0.24 [-1.05, 1.52]	0.36	0.72	27.38	<0.00001	93	0.41	0.752
Sleep efficiency	2	75	-0.28 [-1.07, 0.51]	0.68	0.49	2.63	0.11	62		
Sleep onset latency (min)	6	787	0.81 [0.59, 1.02]	7.33	<0.00001	7.82	0.17	36	0.94	0.198
Sleep duration (min)	6	766	-0.32 [-0.74, 0.11]	1.47	0.14	31.78	<0.00001	84	0.08	0.939
Restorative value of sleep	2	91	0.13 [-0.96, 1.23]	0.24	0.81	5.23	0.02	81		

^aIn order to reduce the heterogeneity between effect sizes, data already reported in similar measures were not included. This decision led to the exclusion of the data provided in two studies ^{23,51} for subjective night awakenings, as well as to the division of sleep duration and sleep onset delay in two distinct variables: sleep duration and sleep onset delay reported as a score in questionnaire or as a length (in min).

Table 3. Summary of the results of the meta-analysis with objective sleep parameters.

Sleep parameter	k	N	SMD (95% CI)	Z	p	Heterogeneity			Egger's test	
						Q	p	I ²	t	p
<i>Polysomnography</i>										
Total sleep time	8	247	-0.90 [-1.51, -0.30]	2.93	0.003	29.76	0.0001	76	0.35	0.734
Sleep onset latency	7	211	0.53 [0.21, 0.86]	3.26	0.001	7.06	0.32	15	0.24	0.818
Stage 1 sleep	8	247	0.48 [0.06, 0.90]	2.25	0.02	15.67	0.03	55	0.37	0.723
Stage 2 sleep	8	247	0.12 [-0.50, 0.73]	0.37	0.71	33.06	<0.0001	79	1.15	0.292
Slow wave sleep	8	247	-0.15 [-0.83, 0.53]	0.43	0.66	41.08	<0.00001	83	0.56	0.596
REM latency	7	211	-0.03 [-0.48, 0.42]	0.13	0.90	13.79	0.03	56	0.42	0.689
REM	9	273	-0.88 [-1.56, -0.21]	2.56	0.01	46.07	<0.00001	83	1.47	0.182
Sleep efficiency	7	238	-1.20 [-1.98, -0.41]	2.99	0.003	40.31	<0.00001	85	1.76	0.138
Wake time	7	211	0.49 [0.11, 0.87]	2.53	0.01	9.67	0.14	38	0.75	0.485
<i>Actigraphy</i>										
Sleep onset latency	5	276	0.80 [0.55, 1.05]	6.23	<0.00001	2.49	0.65	0	0.36	0.745
True sleep	6	301	-0.04 [-0.37, 0.29]	0.24	0.81	9.65	0.09	48	1.74	0.156
Assumed sleep time	2	144	-0.14 [-0.47, 0.20]	0.80	0.42	0.05	0.83	0		
Actual wake time	4	237	0.12 [-0.14, 0.38]	0.92	0.36	2.03	0.57	0	0.45	0.697
Sleep efficiency	6	312	-0.16 [-0.54, 0.22]	0.82	0.41	13.20	0.02	62	0.91	0.416

Table 4. Summary of the results of the post hoc analysis with polysomnographic parameters.

Sleep parameter	k	N	SMD (95% CI)	Z	p	Heterogeneity			Egger's test	
						Q	p	I ²	t	p
Total sleep time	5	178	-1.12 [-1.57, -0.66]	4.76	<0.00001	7.10	0.13	44	1.32	0.278
Sleep onset latency	5	178	0.56 [0.21, 0.90]	3.12	0.002	4.68	0.32	15	0.95	0.413
Stage 1 sleep	5	178	0.20 [-0.11, 0.52]	1.28	0.20	2.25	0.69	0	4.02	0.028
Stage 2 sleep	5	178	0.03 [-0.68, 0.73]	0.07	0.94	18.55	0.001	78	0.35	0.747
Slow wave sleep	5	178	-0.17 [-1.14, 0.80]	0.35	0.73	33.61	<0.00001	88	0.40	0.714
REM latency	5	178	0.01 [-0.57, 0.58]	0.02	0.98	12.42	0.01	68	0.68	0.545
REM	6	204	-0.58 [-1.13, -0.02]	2.04	0.04	16.49	0.006	70	0.04	0.972
Sleep efficiency	5	178	-0.90 [-1.38, -0.42]	3.67	0.0002	8.14	0.09	51	0.44	0.689
Wake time	5	178	0.57 [0.10, 1.04]	2.38	0.02	8.30	0.08	52	0.59	0.597

FIGURES CAPTIONS

Figure 1. PRISMA flow chart.

Footnote : *reasons for exclusion for each paper are reported in the Supplemental material 2

Acknowledgements

We are very grateful to the authors of five of the studies included in the meta-analysis (Aathira et al., 2017; Bruni et al., 2007; Kelmanson 2018; Lopez-Wagner et al., 2008; Miano et al., 2007) for providing additional information. We would also acknowledge Dr Yuta Aoki for his valuable help with translation of paper in Japanese.

References

1. Association AP. *DSM 5.*; 2013. doi:10.1176/appi.books.9780890425596.744053
2. WHO. *International Classification of Diseases*. 11 ط. WHO; 2018.
3. Elrod MG, Nylund CM, Susi AL, وآخ. Prevalence of Diagnosed Sleep Disorders and Related Diagnostic and Surgical Procedures in Children with Autism Spectrum Disorders. *J Dev Behav Pediatr*. 2016;37(5):377–384. doi:10.1097/DBP.0000000000000248
4. Fadini CC, Lamã´nica DA, Fett-Conte AC, وآخ. Influence of sleep disorders on the behavior of individuals with autism spectrum disorder. *Front Hum Neurosci*. 2015;9(June):1–8. doi:10.3389/fnhum.2015.00347
5. Hodge D, Carollo TM, Lewin M, Hoffman CD, Sweeney DP. Sleep patterns in children with and without autism spectrum disorders: Developmental comparisons. *Res Dev Disabil*. 2014;35(7):1631–1638. doi:10.1016/j.ridd.2014.03.037
6. Levin A, Scher A. Sleep Problems in Young Children with Autism Spectrum Disorders: A Study of Parenting Stress, Mothers’ Sleep-Related Cognitions, and Bedtime Behaviors. *CNS Neurosci Ther*. 2016;22(11):921–927. doi:10.1111/cns.12651
7. Hirata I, Mohri I, Kato-Nishimura K, وآخ. Sleep problems are more frequent and associated with problematic behaviors in preschoolers with autism spectrum disorder. *Res Dev Disabil*. 2016;49–50:86–99. doi:10.1016/j.ridd.2015.11.002
8. Kelmanson IA. Sleep disturbances and their associations with emotional / behavioural problems in 5-year-old boys with autism spectrum disorders. *Early Child Dev Care*. 2018;0(0):1–16. doi:10.1080/03004430.2018.1464622
9. Taylor MA, Schreck KA, Mulick JA. Sleep disruption as a correlate to cognitive and adaptive behavior problems in autism spectrum disorders. *Res Dev Disabil*. 2012;33(5):1408–1417. doi:10.1016/j.ridd.2012.03.013
10. Elrod MG, Hood BS. Sleep Differences Among Children With Autism Spectrum Disorders and Typically Developing Peers. *J Dev Behav Pediatr*. 2015;36(3):166–177. doi:10.1097/DBP.0000000000000140
11. Stroup DF, Berlin JA, Morton SC, وآخ. *Meta-analysis of Observational Studies*. 2000.
12. Liberati A, Altman DG, Tetzlaff J, وآخ. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009;339. doi:10.1136/bmj.b2700
13. Hedges L V. Distribution Theory for Glass’s Estimator of Effect size and Related Estimators. *J Educ Behav Stat*. 1981;6(2):107–128. doi:10.3102/10769986006002107
14. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539–1558. doi:10.1002/sim.1186
15. Stuck AE, Rubenstein LZ, Wieland D, وآخ. Bias in meta-analysis detected by a simple,

- graphical. *Bmj*. 1998;316(7129):469–469. doi:10.1136/bmj.316.7129.469
16. Aathira R, Gulati S, Tripathi M, وآخ. Prevalence of Sleep Abnormalities in Indian Children With Autism Spectrum Disorder: A Cross-Sectional Study. *Pediatr Neurol*. 2017;74:62–67. doi:10.1016/j.pediatrneurol.2017.05.019
 17. Al-Farsi OA, Al-Farsi YM, Al-Sharbati MM, Al-Adawi S. Sleep habits and sleep disorders among children with autism spectrum disorders, intellectual disabilities and typically developing children in Oman: a case-control study. *Early Child Dev Care*. 2018;0(0):1–11. doi:10.1080/03004430.2018.1455671
 18. Allik H, Larsson J-O, Smedje H. Sleep Patterns of School-Age Children with Asperger Syndrome or High-Functioning Autism. *J Autism Dev Disord*. 2006;36(5):585–595. doi:10.1007/s10803-006-0099-9
 19. Baker E, Richdale A, Short M, Gradisar M. An investigation of sleep patterns in adolescents with high-functioning autism spectrum disorder compared with typically developing adolescents. *Dev Neurorehabil*. 2013;16(3):155–165. doi:10.3109/17518423.2013.765518
 20. Bruni O, Ferri R, Vittori E, وآخ. Sleep architecture and NREM alterations in children and adolescents with Asperger syndrome. *Sleep*. 2007;30(11):1577–1585. doi:10.1093/sleep/30.11.1577
 21. Chou MC, Chou WJ, Chiang HL, وآخ. Sleep problems among Taiwanese children with autism, their siblings and typically developing children. *Res Autism Spectr Disord*. 2012;6(2):665–672. doi:10.1016/j.rasd.2011.09.010
 22. Cotton S, Richdale A. Brief report: Parental descriptions of sleep problems in children with autism, Down syndrome, and Prader-Willi syndrome. *Res Dev Disabil*. 2006;27(2):151–161. doi:10.1016/j.ridd.2004.12.003
 23. Cotton SM, Richdale AL. Sleep patterns and behaviour in typically developing children and children with autism, Down syndrome, Prader-Willi syndrome and intellectual disability. *Res Autism Spectr Disord*. 2010;4(3):490–500. doi:10.1016/j.rasd.2009.11.006
 24. Couturier JL, Speechley KN, Steele M, Norman R, Stringer B, Nicolson R. Parental perception of sleep problems in children of normal intelligence with pervasive developmental disorders: Prevalence, severity, and pattern. *J Am Acad Child Adolesc Psychiatry*. 2005;44(8):815–822. doi:10.1097/01.chi.0000166377.22651.87
 25. Elia M, Ferri R, Musumeci SA, Bergonzi P. Rapid eye movement modulation during night sleep in autistic subjects. *Brain Dysfunction*. 1991.
 26. Elia M, Ferri R, Musumeci SA, وآخ. Sleep in subjects with autistic disorder: A neurophysiological and psychological study. *Brain Dev*. 2000;22(2):88–92. doi:10.1016/S0387-7604(99)00119-9
 27. Fletcher FE, Foster-Owens MD, Conduit R, Rinehart NJ, Riby DM, Cornish KM. The developmental trajectory of parent-report and objective sleep profiles in autism spectrum disorder: Associations with anxiety and bedtime routines. *Autism*.

2017;21(4):493–503. doi:10.1177/1362361316653365

28. Giannotti F, Cortesi F, Cerquiglini A, وآخ. An investigation of sleep characteristics, EEG abnormalities and epilepsy in developmentally regressed and non-regressed children with autism. *J Autism Dev Disord*. 2008;38(10):1888–1897. doi:10.1007/s10803-008-0584-4
29. Giannotti F, Cortesi F, Cerquiglini A, Vagnoni C, Valente D. Sleep in children with autism with and without autistic regression. *J Sleep Res*. 2011;20(2):338–347. doi:10.1111/j.1365-2869.2010.00882.x
30. Guler S, Yesil G, Ozdil M, Ekici B, Onal H. Sleep disturbances and serum vitamin D levels in children with autism spectrum disorder. 2016;9(7):14691–14697.
31. Han PP, Zou MY, Yang XL, وآخ. [Sleep problems and the association with the levels of 6-sulfatoxymelatonin in children with autism spectrum disorder]. *Zhonghua er ke za zhi = Chinese J Pediatr*. 2017;55(12):911–915.
32. Harder R, Malow BA, Goodpaster RL, وآخ. Heart rate variability during sleep in children with autism spectrum disorder. *Clin Auton Res*. 2016;26(6):423–432. doi:10.1007/s10286-016-0375-5
33. Henderson JA, Barry TD, Bader SH, Jordan SS. The relation among sleep, routines, and externalizing behavior in children with an autism spectrum disorder. *Res Autism Spectr Disord*. 2011;5(2):758–767. doi:10.1016/j.rasd.2010.09.003
34. Hoffman CD, Sweeney DP, Gilliam JE, Lopez-Wagner MC. Sleep Problems in Children With Autism and in Typically Developing Children. *Focus Autism Other Dev Disabl*. 2006;21(3):146–152. doi:10.1177/10883576060210030301
35. Inanuma K. Sleep-wake patterns in autistic children. *Japanese J Child Adolesc Psychiatry*. 1984;25(4):205–217.
36. Kheirouri S, Kalejahi P, Noorazar SG. Plasma levels of serotonin, gastrointestinal symptoms, and sleep problems in children with autism. *Turkish J Med Sci*. 2016;46(6):1765–1772. doi:10.3906/sag-1507-68
37. Lambert A, Tessier S, Rochette AC, Scherzer P, Mottron L, Godbout R. Poor sleep affects daytime functioning in typically developing and autistic children not complaining of sleep problems: A questionnaire-based and polysomnographic study. *Res Autism Spectr Disord*. 2016;23:94–106. doi:10.1016/j.rasd.2015.11.010
38. Li S-Y, Jin Y, Jing J et al. Sleep behaviors in children with autism spectrum disorders. *Chinese Ment Heal J*. 2012;26(6):471–475.
39. Lopez-Wagner MC, Hoffman CD, Sweeney DP, Hodge D, Gilliam JE. Sleep problems of parents of typically developing children and parents of children with autism. *J Genet Psychol*. 2008;169(3):245–259. doi:10.3200/GNTP.169.3.245-260
40. Malow BA, Crowe C, Henderson L, وآخ. A sleep habits questionnaire for children with autism spectrum disorders. *J Child Neurol*. 2009;24(1):19–24. doi:10.1177/0883073808321044

41. Maski K, Holbrook H, Manoach D, Hanson E, Kapur K, Stickgold R. Sleep Dependent Memory Consolidation in Children with Autism Spectrum Disorder. *Sleep*. 2015;38(12):1955–1963. doi:10.5665/sleep.5248
42. Matsuoka M, Nagamitsu S, Iwasaki M, وآخ. High incidence of sleep problems in children with developmental disorders: Results of a questionnaire survey in a Japanese elementary school. *Brain Dev*. 2014;36(1):35–44. doi:10.1016/j.braindev.2012.12.004
43. May T, Cornish K, Conduit R, Rajaratnam SMW, Rinehart NJ. Sleep in High-Functioning Children With Autism: Longitudinal Developmental Change and Associations With Behavior Problems. *Behav Sleep Med*. 2015;13(1):2–18. doi:10.1080/15402002.2013.829064
44. Miano S, Bruni O, Elia M, وآخ. Sleep in children with autistic spectrum disorder: A questionnaire and polysomnographic study. *Sleep Med*. 2007;9(1):64–70. doi:10.1016/j.sleep.2007.01.014
45. Mutluer T, Karakoc Demirkaya S, Abali O. Assessment of sleep problems and related risk factors observed in Turkish children with Autism spectrum disorders. *Autism Res*. 2016;9(5):536–542. doi:10.1002/aur.1542
46. Paavonen EJ, Vehkalahti K, Vanhala R, Von Wendt L, Nieminen-von Wendt T, Aronen ET. Sleep in children with Asperger syndrome. *J Autism Dev Disord*. 2008;38(1):41–51. doi:10.1007/s10803-007-0360-x
47. Pace M, Dumortier L, Favre-Juvin A, Guinot M, Bricout VA. Heart rate variability during sleep in children with autism spectrum disorders. *Physiol Behav*. 2016;167:309–312. doi:10.1016/j.physbeh.2016.09.027
48. Park S, Cho SC, Cho IH, وآخ. Sleep problems and their correlates and comorbid psychopathology of children with autism spectrum disorders. *Res Autism Spectr Disord*. 2012;6(3):1068–1072. doi:10.1016/j.rasd.2012.02.004
49. Patzold LM, Richdale AL, Tonge BJ. An investigation into sleep characteristics of children with autism and Asperger’s Disorder. *J Paediatr Child Health*. 1998;34(6):528–533. doi:10.1046/j.1440-1754.1998.00291.x
50. Phung JN, Goldberg WA. Poor sleep quality is associated with discordant peer relationships among adolescents with Autism Spectrum Disorder. *Res Autism Spectr Disord*. 2017;34:10–18. doi:10.1016/j.rasd.2016.11.008
51. Richdale AL, Prior MR. The sleep/wake rhythm in children with autism. *Eur Child Adolesc Psychiatry*. 1995;4(3):175–186. doi:10.1007/BF01980456
52. Souders MC, Mason TBA, Valladares O, وآخ. Sleep Behaviors and Sleep Quality in Children with Autism Spectrum Disorders. *Sleep*. 2009. doi:10.1093/sleep/32.12.1556
53. Tessier S, Lambert A, Scherzer P, Jemel B, Godbout R. REM sleep and emotional face memory in typically-developing children and children with autism. *Biol Psychol*. 2015;110:107–114. doi:10.1016/j.biopsycho.2015.07.012
54. Tzischinsky O, Meiri G, Manelis L, وآخ. Sleep disturbances are associated with

- specific sensory sensitivities in children with autism. *Mol Autism*. 2018;9(1):1–10. doi:10.1186/s13229-018-0206-8
55. van der Heijden KB, Stoffelsen RJ, Popma A, Swaab H. Sleep, chronotype, and sleep hygiene in children with attention-deficit/hyperactivity disorder, autism spectrum disorder, and controls. *Eur Child Adolesc Psychiatry*. 2018;27(1):99–111. doi:10.1007/s00787-017-1025-8
 56. Yang XL, Liang S, Zou MY, وآخ. Are gastrointestinal and sleep problems associated with behavioral symptoms of autism spectrum disorder? *Psychiatry Res*. 2018;259(January 2017):229–235. doi:10.1016/j.psychres.2017.10.040
 57. Perlis ML, Giles DE, Mendelson WB, Bootzin RR, Wyatt JK. Psychophysiological insomnia: The behavioural model and a neurocognitive perspective. *J Sleep Res*. 1997;6(3):179–188. doi:10.1046/j.1365-2869.1997.00045.x
 58. Choi J, Yoon I-Y, Kim H-W, Chung S, Yoo HJ. Differences between objective and subjective sleep measures in children with attention deficit hyperactivity disorder. *J Clin Sleep Med JCSM Off Publ Am Acad Sleep Med*. 2010;6(6):589–595. http://ezproxy.deakin.edu.au/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=mdc&AN=21206548&site=ehost-live&scope=site%5Cnhttp://www.ncbi.nlm.nih.gov/pmc/articles/PMC3014246/pdf/jcs_m.6.6.589.pdf.
 59. Cortese S, Faraone S V., Konofal E, Lecendreux M. Sleep in Children With Attention-Deficit/Hyperactivity Disorder: Meta-Analysis of Subjective and Objective Studies. *J Am Acad Child Adolesc Psychiatry*. 2009;48(9):894–908. doi:10.1097/CHI.0b013e3181ac09c9
 60. Chokroverty S, محرر. *Sleep Disorders Medicine. Basic Science, Technical Considerations and Clinical Aspects*. 3rd editio. Elsevier Inc.; 2009.
 61. Kryger, M., Roth, T., Dement WC, محرر. *Principles and Practice of Sleep Medicine*. 6th editio. Elsevier Inc.; 2017.
 62. Mazzone L, Postorino V, Siracusano M, Riccioni A, Curatolo P. The Relationship between Sleep Problems, Neurobiological Alterations, Core Symptoms of Autism Spectrum Disorder, and Psychiatric Comorbidities. *J Clin Med*. 2018;7(5):102. doi:10.3390/jcm7050102
 63. Tordjman S, Anderson GM, Pichard N, Charbuy H, Touitou Y. Nocturnal excretion of 6-sulphatoxymelatonin in children and adolescents with autistic disorder. *Biol Psychiatry*. 2005;57(2):134–138. doi:10.1016/j.biopsych.2004.11.003
 64. Pagan C, Goubran-Botros H, Delorme R, وآخ. Disruption of melatonin synthesis is associated with impaired 14-3-3 and miR-451 levels in patients with autism spectrum disorders. *Sci Rep*. 2017. doi:10.1038/s41598-017-02152-x
 65. Pagan C, Delorme R, Callebert J, وآخ. The serotonin-N-acetylserotonin-melatonin pathway as a biomarker for autism spectrum disorders. *Transl Psychiatry*. 2014. doi:10.1038/tp.2014.120

66. Bruni O, Ottaviano S, Guidetti V, وآخ. The Sleep Disturbance Scale for Children (SDSC) construction and validation of an instrument to evaluate sleep disturbances in childhood and adolescence. *J Sleep Res.* 1996. doi:10.1111/j.1365-2869.1996.00251.x
67. Owens JA, Spirito A, McGuinn M. The Children's Sleep Habits Questionnaire (CSHQ): psychometric properties of a survey instrument for school-aged children. *Sleep.* 2000. doi:10.1111/j.1469-8749.2001.tb00204.x