

Solid Phase Synthesis of Globomycin and SF-1902 A5

| | |
|-------------------------------|---|
| Journal: | <i>The Journal of Organic Chemistry</i> |
| Manuscript ID: | jo-2010-025145.R2 |
| Manuscript Type: | Article |
| Date Submitted by the Author: | 09-Feb-2011 |
| Complete List of Authors: | Sarabia, Francisco; University of Malaga, Organic Chemistry Chammaa, Samy; University of Malaga, Organic Chemistry García-Ruiz, Cristina; University of Malaga, Organic Chemistry |
| | |

SCHOLARONE™
Manuscripts


Solid Phase Synthesis of Globomycin and SF-1902 A₅

Francisco Sarabia,* Samy Chammaa and Cristina García-Ruiz

Department of Organic Chemistry, Faculty of Sciences.

University of Malaga. Campus de Teatinos s/n 29071. Malaga. SPAIN

frsarabia@uma.es




The syntheses of the natural lipocyclodepsipeptide-type antibiotics globomycin and SF-1902 A₅ are reported utilizing solid phase technology for the construction of the peptidic fragment and a new asymmetric methodology of epoxidation for the preparation of the lipidic chain. The linkage between both fragments was successfully achieved in solid phase to complete the syntheses via a macrolactonization reaction executed prior to the cleavage of the acyclic precursors from the solid support. These syntheses provide access to the rapid generation of a library of analogues via modification of the aminoacid residues as well as the lipidic chain, thus facilitating the identification of new antibiotics with interesting mechanisms of action based upon the inhibition of the enzyme signal peptidase II.

Introduction


The lipocyclodepsipeptides¹ represent a rich and intriguing class of natural products that possess a broad range of biological properties including antitumoral, antibiotic, antifungal and antiinflammatory activities.² Among them, globomycin (**1**)³ and its congeners **2-6**⁴ (Figure 1, part A), isolated from four different strains of actinomycetes in 1978, are representative and interesting examples, especially featured by their striking antibiotic activities against Gram-negative bacteria.⁵ Recently proven as specific inhibitors of signal peptidase II,⁶ an enzyme responsible for transforming acylated lipoproteins into apolipoproteins,⁷ these compounds trigger the accumulation of acylated forms of lipoproteins in the cytoplasmatic membrane resulting in cell death. Therefore, this enzyme represents an attractive biological target for the development of new antibiotics,⁸ which has not been fully exploited. In fact, despite the discovery of this class of natural products being reported more than thirty years ago, globomycin (**1**), the major component of this family, has been commonly used as a biochemical tool for the identification of new lipoproteins amenable to acylation⁹ and for biosynthetic studies of them.^{10,11} Comparatively, globomycin and SF-1902 A₅ display greater antibiotic activities when compared to other antibiotics,¹² such as ampicillin or streptomycin, against *E. coli*, *Salmonella enteritidis* and *Enterobacter cloacae*. On the other hand, SF-1902 A₅ is more active than globomycin (MIC against *E. coli* NIHJ JC-2: 6.25 µg/mL for **1**, 1.56 µg/mL for **2**), revealing that the lipidic chain plays an important role in the biological activity. More recently, it was discovered that globomycin exhibited antibacterial activity against *Mycobacterium tuberculosis*, showing that the biological action was not dependent on its inhibition of signal peptidase II, thus suggesting an alternative mechanism of action in this bacteria.¹³ All these biological properties render these cyclodepsipeptides as attractive synthetic endeavours in the search for new antibiotics. Chemically, the structure of globomycin, established by Haneishi *et al* in 1980,¹⁴ was recently confirmed with its first total

synthesis in 2000 by the Kogen group.¹⁵ This same group has investigated the initial chemistry in the area with the synthesis of SF-1902 A₅¹⁶ and the preparation of a series of analogues with modifications of various constituents of the molecule including the amino acids, lipidic chain, stereochemistry and heteroatoms.^{12,17} This impressive chemical study led to the generation of a set of fifteen analogues whose biological evaluation allowed the establishment of a preliminary structure-activity relationship describing the structural elements essentials for biological activity (Figure 1, part B).

FIGURE 1. Part A: Molecular Structures of Globomycin and its Congeners. **Part B:** SAR Studies of Analogues.




(B) Structure-Activity Relationship Studies by the Kogen's Group



Based on these chemical precedents and prompted by the potential that these compounds may represent in the field of antibiotics, as well as their unique mechanism of action, we decided to embark on a research program directed to the design of an efficient and

readily accessible route towards this class of natural products and analogues. With this purpose in mind, we initially planned the synthesis of the more relevant natural members, globomycin (**1**) and SF-1902 A₅ (**2**) utilizing solid phase synthesis as a suitable technology to reach our goals. To this aim, resins **7** and **8** would contain the acyclic precursors of **1** and **2**, which would be prepared via a macrolactonization reaction after the cleavage of such precursors from the resin. Functionalized resins **7** and **8** would be prepared via coupling of the acids **9** or **10** with the peptidic chain linked in a solid support in form of resin **11** (Scheme 1).

SCHEME 1. Synthetic Plan for Globomycin and SF-1902 A₅.



In the present article we describe the total synthesis of these natural products based on the delineated strategy to provide access to analogues with modification of the amino acids


1
2 and the lipidic chain in an efficient and rapid way, taking advantage of solid phase
3 methodology.
4
5
6
7

8 9 **Results and Discussion**

10
11
12
13
14 We began the synthesis of globomycin and SF-1902 A₅ with the construction of the
15 linear peptide on a 2-chlorotrityl chloride (CTC) resin¹⁸ using the Fmoc strategy.¹⁹ Thus,
16 Fmoc-Gly was linked onto the CTC resin by esterification with *N,N*-diisopropylethylamine
17 (DIPEA) to obtain resin **12**. After removing the Fmoc (9-fluorenylmethoxycarbonyl) group
18 (DIPEA) to obtain resin **12**. After removing the Fmoc (9-fluorenylmethoxycarbonyl) group
19 by treatment with 20% of piperidine in *N,N*-dimethylformamide (DMF), the following Fmoc
20 aminoacid derivative, Fmoc-*L-allyl*-Thr(TBS)-OH,²⁰ was loaded onto the resulting resin by the
21 action of *N,N'*-diisopropylcarbodiimide (DIC) in the presence of 1-hydroxybenzotriazole
22 (HOBt) in DMF. To check the loading of the amino acid and to assure the effectiveness of the
23 chosen coupling procedure, we decided to cleave the dipeptide by treatment of **13** with
24 CH₂Cl₂/AcOH/CF₃CH₂OH (TFE) (7/2/1), which gave pure dipeptide **14**. The synthesis was
25 continued from **13** repeating the procedure of coupling and Fmoc deprotection steps for each
26 amino acid being loaded in the following order: 1) Fmoc-*L*-Ser(Bzl)-OH, 2) Fmoc-*L-allyl*-Ile-
27 OH and 3) Fmoc-*N*-Me-*L*-Leu-OH to obtain resin **15**. At this stage, we again checked the
28 overall loading of the resin, thus **15** was treated with CH₂Cl₂/AcOH/TFE (7/2/1) to provide
29 pure Fmoc protected pentapeptide **16** in an amount that revealed a load of 0.6 mmol/g.
30 Finally, resin **15** was treated with piperidine to remove the Fmoc group and prepare the
31 polymer-bound pentapeptide **11** for the coupling with the fatty acid derivatives **9** or **10**
32 (Scheme 2).
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHEME 2. Synthesis of the Peptidic

Fragment in Solid Phase.



For the synthesis of the lipidic fragments, we planned to apply the recent methodology of asymmetric epoxidation developed in our laboratories based on the use of cyclic sulfonium salts derived from α -amino acids²¹ as a means of stereoselectively constructing an oxirane ring.²² In fact, our experience in this field has allowed us to demonstrate the utility and efficiency of the resulting epoxy amides in reactions with nucleophiles with complete C-2

1 regioselectivity.²³ Thus, treatment of commercially available aldehydes **18** and **19** with the
2 sulfur ylide, generated from its corresponding sulfonium salt **17**, provided epoxy amides **20**
3 and **21** in 80 and 73% yields respectively and excellent diastereoselectivities (>98% according
4 to NMR and GC-MS analyses). This excellent diastereoselectivity was confirmed via
5 reduction of epoxyamide **20** to epoxy alcohol **30** (See Part B of Scheme 3) by the action of
6 Super-H in 85% yield, displaying spectroscopic and physical properties, especially its optical
7 rotation ($[\alpha]_{\text{D}}^{25} = +39.5^{\circ}$ (c 1.0, CHCl_3); lit.²⁴ $[\alpha]_{\text{D}}^{20} = +38.7^{\circ}$ (c 0.03, CHCl_3)), that matched
8 with the described in the literature.²⁴


9
10
11
12
13
14
15
16
17
18
19
20
21
22
23 Therefore, we proceeded with the introduction of the methyl groups at the C-2
24 position. To this end, epoxy amides **20** and **21** were exposed to the action of lithium
25 dimethylcuprate²⁵ to obtain hydroxy amides **22** and **23** in excellent yields (82 and 96%
26 respectively). At this stage, protection of the hydroxyl groups of both hydroxy amides was
27 required prior to the reduction of the amide functionality to the corresponding alcohol. Thus,
28 **22** and **23** were transformed into their corresponding silyl ethers **24** and **25** and then subjected
29 to the action of lithium triethylborohydride (Super-H).²⁶ Unexpectedly, these reactions did not
30 proceed as desired, recovering starting material in both cases. The lack of reactivity of
31 compounds **24** and **25** towards Super-H was ascribed to steric factors, thus prompting us to
32 consider alternative methods to achieve the amide reduction. Among the various alternatives
33 described in the literature,²⁷ we considered the use of LiNH_2BH_3 , which has been proven to be
34 an efficient reducing agent for the conversion of amides to alcohols.²⁸ Thus, when **24** and **25**
35 were treated with this reagent at room temperature, a 1.5:1 mixture consisting of the desired
36 alcohols²⁹ together with the desilylated derivatives, diols **26** and **27**, were obtained in 75%
37 combined yields. In light of these promising results, we carried out this reaction under reflux
38 in THF finding that, after four hours, **24** and **25** were transformed into the corresponding diols
39 **26** and **27** in 85 and 92% yields respectively. Unable to preserve the silyl protecting group
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 during the reduction process, we decided to attempt this transformation directly from the
2 hydroxy amides **22** and **23** by use of a large excess of LiNH_2BH_3 in refluxing THF. To our
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


delight, these reactions afforded very good results, providing the corresponding diols **26** and **27** in very high yields (90 and 93% respectively). Finally, chemoselective oxidations of **26** and **27** to the targeted fatty hydroxy acids **9** and **10** were accomplished by the oxidative system 2,2,6,6-tetramethyl-1-piperidinyloxy free radical/bisacetoxyiodobenzene (TEMPO/BAIB) in the presence of water³⁰ (Scheme 3, part A). Comparatively, Sharpless asymmetric epoxidation³¹ was similarly exploited for allylic alcohol **28** to obtain epoxy alcohol **30**³² in 75% yield. However, the opening reaction of the oxirane ring with lithium dimethylcuprate furnished an inseparable mixture of 2-methyl and 3-methyl opened products³³ in a 3:1 ratio. The mixture was subjected to the action of sodium periodate and the desired diol **26** was isolated in 41% yield from epoxy alcohol **30** after purification by flash column chromatography. Oxidation of **26** provided the desired hydroxy acid **9** as described above. In order to circumvent the lack of regioselectivity observed during the opening process, we prepared epoxy alcohol **31**,³⁴ via Sharpless asymmetric epoxidation of allylic alcohol **29** in 70% yield, which was then smoothly oxidized to the corresponding epoxy aldehyde with $\text{DMSO}/\text{pyr}.\text{SO}_3$.³⁵ With this epoxy aldehyde in hand, we planned to use Bode's carbene-catalyzed epoxide-opening reaction³⁶ utilizing the thiazolium salt **32**³⁷ which has proven to be very efficient in stereoselective oxidative openings of epoxy aldehydes. In our case, this reaction afforded the corresponding 2-methyl-3-hydroxy ester **33** with an *anti* relative configuration accompanied with its *syn* isomer in a 7:3 ratio and in 70% overall yield from **31**. Finally, hydrolysis of **33** provided hydroxy acid **9** contaminated with its C-2 epimer which was not separable by chromatographic methods (Scheme 3, part B). In summary, we concluded that our approach using sulfur ylides was more efficient than the Sharpless epoxidation approach in terms of chemical yields as well as in regio- and stereo-selectivities.

SCHEME 3. Synthesis of the Lipidic Chains of
Globomycin and SF-1902 A₅.

A) Via Sulfur Ylides




B) Via Sharpless Asymmetric Epoxidation



Having extensively explored the synthesis of the lipidic chain via epoxide chemistry, with concomitant manipulation of the amide function by reduction to the alcohol and reoxidation to the carboxylic acid, we did not wish to discard the possibility of a direct hydrolysis of the amide group despite its recognized robustness. To this aim, we initially

1 hydrolyzed the acetal function installed at the amino alcohol unit by treating hydroxy amide
2
3
4 **22** with 1 N aqueous HCl in dioxane³⁸ to provide compound **34** in 98% yield. Attempts at
5
6 direct amide hydrolysis of **22** or from **34** required harsh acidic conditions (H₂SO₄ in reflux)³⁹
7
8 and were unsuccessful due to decomposition of starting material for the first case and
9
10 formation of hydroxy acid **9** for the second, albeit in poor yield (30%). Thus, we proceeded to
11
12 investigate the hydrolysis of compound **34** under mild conditions. This task required
13
14 activating the amide group for subsequent hydrolysis. For this objective, we considered
15
16 different possibilities. One of them was the formation of an oxazolidinone ring, amenable to
17
18 hydrolysis under basic conditions.⁴⁰ For this purpose, dihydroxy amide **34** was transformed
19
20 into its monocarbonate **35** by reaction with 1.0 equivalent of ethyl chloroformate in good
21
22 yield (87%) which was subjected to basic conditions provided by different bases such as
23
24 LDA, LHMDs or NaH. Unfortunately in all attempted cases, we were not able to construct
25
26 the oxazolidinone ring, compound **38**, recovering starting material or obtaining **34** via
27
28 hydrolysis of the carbonate group. Considering the possibility of an interference of the free
29
30 hydroxyl group, we decided to prepare bis (silyl ether) **36** by treatment of **34** with excess of
31
32 *tert*-butyldimethylsilyl trifluoromethanesulfonate (TBSOTf), which was selectively
33
34 desilylated by the action of (1*S*)-(+)-10-camphorsulfonic acid (CSA) to obtain hydroxy amide
35
36 **37**. Reaction of **37** with ethyl chloroformate and subsequent basic treatment of the resulting
37
38 carbonate, or reaction of **37** with triphosgene⁴¹ or *N,N'*-carbonyldiimidazole (CDI)⁴² proved
39
40 similarly unsuccessful in the formation of the coveted oxazolidinone **39**. In an alternative way,
41
42 activation of the amide group by introduction of Boc,⁴³ Ac or Ms⁴⁴ groups via amide
43
44 deprotonation of **36** by the action of base (LDA, BuLi, LHMDs or NaH), followed by the
45
46 addition of the corresponding electrophilic agent, did not provide the expected products **40a-c**,
47
48 recovering starting bis (silyl ether) **36** instead (Scheme 4).
49
50
51
52
53
54
55
56
57
58
59
60

SCHEME 4. Studies on Hydrolysis of Hydroxy


Amide **22**.

In light of the lack of reactivity of **36** or related amides, as described before, we considered the introduction of a *N*-nitroso functional group as a way of activation of the amide group.⁴⁵ Interestingly, *N*-nitrosation of **36** under conventional conditions,⁴⁵ afforded the silyloxy acid derivative **42** in 75% yield, probably formed through *N*-nitroso derivative **41a**, which surprisingly was not detected. Silyloxy acid **42** was subjected to treatment with tetrabutylammonium fluoride (TBAF) to provide the hydroxy acid **9** in 95% yield (Scheme 5). This interesting result encouraged us to try direct *N*-nitrosation of **34** obtaining, in this case, acetate **43**, which, under basic conditions, afforded hydroxy acid **9** in a lower 31% overall yield. Then, we extended this synthetic path to amide **23** to get acid **46**, through compounds

44 and **45** obtained in very similar yields compared with the globomycin series, which was finally desilylated to acid **10** in 98% yield (Scheme 5).

SCHEME 5. N-Nitrosations of Amides 36 and

45. Synthesis of Hydroxy Acids 9 and 10.



With all this chemistry spread around hydroxy amides **34** and **44** and with both peptidic and lipidic key fragments prepared, we proceeded to completion of the synthesis of the natural cyclodepsipetides. For this purpose, we initially coupled hydroxy acids **9** and **10** with peptidic derivative **11** linked onto the resin by treatment with diethyl cyanophosphonate (DEPC).⁴⁶ This treatment was repeated once more in order to obtain a complete loading of acids **9** and **10** onto the solid support through an amide bond to give resins **7** and **8**. Once the


1
2 acyclic precursors were prepared in solid phase, we then achieved the corresponding
3
4 cleavages by the action of AcOH/CF₃CH₂OH in CH₂Cl₂ to obtain compounds **47** and **48** in
5
6 high purity as determined by their NMR spectra and HPLC analyses, verifying the efficiency
7
8 of the solid phase synthesis. The syntheses of globomycin (**1**) and SF-1902 A₅ (**2**) were then
9
10 completed following the procedure described by Kogen *et al* using a Yamaguchi
11
12 macrolactonization⁴⁷ and final deprotection of the silyl and benzyl ether groups via
13
14 compounds **49** and **51** for globomycin (**1**) and **50** and **52** for SF-1902 A₅ (**2**) (Scheme 6).
15
16
17
18
19
20

21 Conclusions

22
23
24
25

26 In conclusion, we have described new syntheses of the antibiotics globomycin (**1**) and
27
28 SF-1902 A₅ (**2**) that incorporate two important novel methodologies for the total syntheses of
29
30 these cyclodepsipeptides: 1) The use of solid phase for the assembly of the peptidic and
31
32 lipidic fragments that are found in these molecules, and 2) extension of our new asymmetric
33
34 methodology of epoxidation for the stereoselective synthesis of the lipidic chains. The
35
36 implementation of this new methodology of epoxidation offers the advantage of constructing
37
38 the lipidic chain through an oxirane ring, a versatile functional group which allows the
39
40 generation of 1,2-bifunctional groups. This in turn provides access to a broad variety of
41
42 modified-lipidic chains which appear to play an essential role in the antibiotic properties of
43
44 these compounds. This feature combined with the use of a solid phase-based synthesis allows
45
46 rapid and easy access to a wide array of globomycin analogues in the quest for new antibiotics
47
48 with novel mechanism of action. The generation of a broad library of globomycin analogues
49
50 as well as their biological evaluations represent our priorities in current and future
51
52 investigations.
53
54
55
56
57
58
59
60

SCHEME 6. Synthesis of Globomycin (**1**) and SF-1902 A₅ (**2**) via a Yamaguchi Macrolactonization.



Experimental⁴⁸

Fmoc-Gly-OH loaded 2-chlorotrityl resin 12. A 5 mL polypropylene syringe fitted with polyethylene porous disk charged with 2-chlorotrityl chloride resin (300 mg, L=1.3 mmol/g, 0.39 mmol, 1.0 equiv.), was loaded with a solution of Fmoc-Gly-OH (348 mg, 1.17 mmol, 3.0 equiv.) and DIPEA (235 μL , 1.37 mmol, 3.5 equiv.) in dry DMF (3 mL). The resulting suspension was shaken at 280 r.p.m. for 30 hours, then, the solution was unloaded

1
2 and the resin washed by shaking with dry DMF (5x3 mL). The resulting swelled resin was
3
4 used in the next step.
5
6
7

8
9 **2-Chlorotrityl-Gly-L-*allo*-Thr(TBS)-Fmoc resin 13.** The polypropylene syringe
10 loaded with the swelled resin **12** was treated with 20% piperidine in DMF (3 x 3 mL x 10
11 min.). After the last run, the resin was washed with dry DMF (5 x 3 mL) and loaded with a
12 solution of Fmoc-L-*allo*-Thr(TBS)-OH (355 mg, 0.78 mmol, 2.0 equiv.), HOBt (107 mg, 0.78
13 mmol, 2.0 equiv.) and DIC (155 μ L, 1.0 mmol, 2.5 equiv.) in dry DMF (3 mL). The resulting
14 suspension was shaken at 280 r.p.m. for 24 hours, and then, the solution was unloaded and the
15 resin washed with dry DMF (5 x 3 mL). The resulting swelled resin was used in the next step.
16
17
18
19
20
21
22
23
24
25

26
27
28 **2-Chlorotrityl-Gly-L-*allo*-Thr(TBS)-L-Ser(Bzl)-L-*allo*-Ile-N-Me-L-Leu-Fmoc**
29 **resin 15.** The polypropylene syringe loaded with the swelled resin **13** was treated with 20%
30 piperidine in DMF (3 x 3 mL x 10 min.), after the last run, the resin was washed with dry
31 DMF (5 x 3 mL) and loaded with a solution of Fmoc-L-Ser(Bzl)-OH (559 mg, 1.18 mmol, 3.0
32 equiv.), HOBt (161 mg, 1.17 mmol, 3.0 equiv.) and DIC (216 μ L, 1.37 mmol, 3.5 equiv.) in
33 dry DMF (3 mL). The resulting suspension was shaken at 280 r.p.m. for 24 hours, and then,
34 the solution was unloaded and the resin washed with dry DMF (5 x 3 mL). The resulting
35 swelled resin was used in the next step. This sequence was repeated with Fmoc-L-*allo*-Ile-OH
36 (276 mg, 0.78 mmol, 2.0 equiv.), HOBt (107 mg, 0.78 mmol, 2.0 equiv.) and DIC (155 μ L,
37 1.0 mmol, 2.5 equiv.) followed, after treatment with 20% piperidine in DMF, by loading of
38 Fmoc-N-Me-L-Leu-OH (358 mg, 0.98 mmol, 2.5 equiv.), HOBt (135 mg, 0.98 mmol, 2.5
39 equiv.) and DIC (195 μ L, 1.25 mmol, 3.2 equiv.). Finally, the resin was washed with DMF (5
40 x 3 mL), DCM (3 x 3 mL), MeOH (3 x 3 mL) and Et₂O (3 x 3 mL). The resulting resin was
41 dried under vacuum to recover 626 mg of polymer-bound N-Fmoc protected pentapeptide **15**.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fmoc-N-Me-L-Leu-L-*allo*-Ile-L-Ser(Bzl)-L-*allo*-Thr(TBS)-Gly-OH (16). Release of a small amount of peptide from resin **15** (17.5 mg) by treatment with CH₂Cl₂:AcOH:TFE (1.0 mL, 7:2:1) gave the *N*-Fmoc protected pentapeptide **16** (9.6 mg), which revealed a load of 0.6 mmol/g for resin **15**. [**16**]: ¹H NMR (400 MHz, DMSO-*d*₆, two rotamers in a 2.9:1 ratio) Major rotamer: δ = 0.03 (s, 3 H), 0.01 (s, 3 H), 0.71-0.89 (m, 9 H), 0.80 (s, 9 H), 0.88 (d, *J* = 6.6 Hz, 3 H), 0.95-1.07 (m, 1 H), 1.05 (d, *J* = 6.2 Hz, 3 H), 1.21-1.36 (m, 2 H), 1.47-1.61 (m, 2 H), 1.73-1.84 (m, 3/4 H), 2.68 (s, 9/4 H), 2.79 (s, 3/4 H), 3.52-3.59 (m, 3 H), 3.77 (dd, *J* = 17.5, 6.2 Hz, 1 H), 4.01 (q, *J* = 6.5 Hz, 1 H), 4.25-4.45 (m, 4 H), 4.38 (dd, *J* = 9.0, 7.2 Hz, 1 H), 4.48 (m, 2 H), 4.65-4.70 (m, 2 H), 7.24-7.33 (m, 7 H), 7.41 (t, *J* = 7.5 Hz, 2 H), 7.62-7.70 (m, 1 H), 7.63 (d, *J* = 7.1 Hz, 2 H), 7.89 (d, *J* = 7.5 Hz, 2 H), 7.99 (d, *J* = 9.0 Hz, 1 H), 8.05 (d, *J* = 7.6 Hz, 1 H), 8.16 (t, *J* = 5.0 Hz, 1 H); FAB HRMS (NBA) *m/e* 930.5043, M+H⁺ calcd for C₅₀H₇₁N₅O₁₀Si 930.5049.

Epoxy amide 20. To a solution of sulfonium salt **17** (3.66 g, 11.59 mmol, 1.2 equiv) in ^tBuOH (20.0 mL) was added a solution of NaOH 3.0 M in H₂O (3.86 mL, 11.59 mmol, 1.2 equiv). After 1 h at 25 °C, a solution of heptanal **18** (1.34 mL, 9.66 mmol, 1.0 equiv) in ^tBuOH (5.0 mL) was added and the reaction mixture was vigorously stirred overnight at 25°C. After this time, reaction mixture was diluted with Et₂O and H₂O, both phases separated and the aqueous layer extracted with Et₂O twice. Combined organic extracts were then washed with water and brine, filtered and concentrated. Purification by flash column chromatography (silica gel, 30% AcOEt in hexanes) provided epoxy amide **20** (2.55 g, 80%) as a colorless oil: *R*_f = 0.48 (silica gel, 50% AcOEt in hexanes); [α]_D²⁵ = +11.9° (*c* 1.8, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.86 (t, *J* = 6.9 Hz, 3 H), 1.26-1.30 (m, 4 H), 1.32-1.36 (m, 2 H), 1.40-1.55 (m, 2 H), 1.51 (s, 3 H), 1.62 (s, 3 H), 1.67-1.74 (m, 2 H), 1.77-1.81 (m, 1 H), 2.01-2.07 (m, 1 H), 2.10 (s, 3 H), 2.45 (ddd, *J* = 13.1, 8.6, 7.0 Hz, 1 H), 2.58 (ddd, *J* = 12.7, 7.3, 5.2 Hz, 1 H), 3.15 (ddd, *J* = 6.0, 4.3, 1.9 Hz, 1 H), 3.32 (d, *J* = 1.9 Hz, 1 H), 3.88 (d, *J* = 9.2 Hz, 1 H), 3.99

(ddd, $J = 9.2, 5.3, 1.3$ Hz, 1 H), 4.28 (ddd, $J = 10.3, 5.0, 3.2$ Hz, 1 H); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 14.0, 16.0, 22.5, 22.9, 25.8, 26.3, 29.0, 30.9, 31.5, 31.7, 34.5, 53.9, 55.8, 58.5, 67.0, 95.8, 164.2$; FAB HRMS (NBA) m/e 330.2093, $\text{M}+\text{H}^+$ calcd for $\text{C}_{17}\text{H}_{31}\text{NO}_3\text{S}$ 330.2103.

Epoxy amide 21. To a solution of sulfonium salt **17** (2.51 g, 7.95 mmol, 1.2 equiv) in $t\text{BuOH}$ (12.0 mL) was added a solution of NaOH 3.0 M in H_2O (2.65 mL, 7.95 mmol, 1.2 equiv). After 1 h at 25°C , a solution of nonanal **19** (1.14 mL, 6.63 mmol) in $t\text{BuOH}$ (5.0 mL) was added and the reaction mixture was vigorously stirred overnight at 25°C . After this time, reaction mixture was diluted with Et_2O and H_2O , both phases separated and the aqueous layer extracted with Et_2O twice. Combined organic extracts were then washed with water and brine, filtered and concentrated. Purification by flash column chromatography (silica gel, 30% AcOEt in hexanes) provided epoxy amide **21** (1.72 g, 73%) as a colorless oil: $R_f = 0.51$ (silica gel, 50% AcOEt in hexanes); $[\alpha]_D^{25} = +19.4^\circ$ (c 0.8, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) $\delta = 0.88$ (t, $J = 7.0$ Hz, 3 H), 1.20-1.37 (m, 12 H), 1.43-1.49 (m, 1 H), 1.53 (s, 3 H), 1.64 (s, 3 H), 1.69-1.78 (m, 1 H), 1.79-1.84 (m, 1 H), 2.03-2.08 (m, 1 H), 2.12 (s, 3 H), 2.47 (ddd, $J = 13.1, 8.7, 7.0$ Hz, 1 H), 2.60 (ddd, $J = 12.8, 7.4, 5.2$ Hz, 1 H), 3.17 (ddd, $J = 6.5, 4.6, 2.0$ Hz, 1 H), 3.33 (d, $J = 2.0$ Hz, 1 H), 3.89 (d, $J = 9.2$ Hz, 1 H), 4.01 (ddd, $J = 9.2, 5.3, 1.5$ Hz, 1 H), 4.30 (ddd, $J = 10.3, 4.9, 3.1$ Hz, 1 H); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 14.1, 16.0, 22.6, 22.9, 25.9, 26.3, 29.2, 29.3, 29.4, 30.9, 31.5, 31.8, 34.6, 53.9, 55.8, 58.5, 67.0, 95.8, 164.2$; FAB HRMS (NBA) m/e 358.2386, $\text{M}+\text{H}^+$ calcd for $\text{C}_{19}\text{H}_{35}\text{NO}_3\text{S}$ 358.2416.

Hydroxy Amide 22. To a suspension of CuI (925 mg, 4.84 mmol, 4.0 equiv) in THF (20 mL) was added dropwise MeLi (1.6 M in Et_2O , 6.10 mL, 9.68 mmol, 8.0 equiv) at 0°C . Then, a solution of epoxy amide **20** (400 mg, 1.21 mmol, 1.0 equiv) in THF (5 mL) was added to the resulting colorless solution of Me_2CuLi at 0°C . The reaction mixture was stirred for 8 h at this temperature and quenched by careful addition of aqueous saturated NH_4Cl

1
2 solution, followed by dilution with Et₂O. After separation of both phases, the aqueous phase
3
4 was extracted with Et₂O and the combined organic layers were sequentially washed with
5
6 aqueous saturated NH₄Cl solution, water and brine. After treatment with MgSO₄, the solvents
7
8 were removed by reduced pressure to obtain crude 2-methyl-3-hydroxy amide which was
9
10 subjected to purification by flash column chromatography (silica gel, 50% AcOEt in hexanes)
11
12 to obtain **22** (341 mg, 82%) as a colorless oil: $R_f = 0.38$ (silica gel, 33% AcOEt in hexanes);
13
14 $[\alpha]_D^{25} = +19.7^\circ$ (*c* 0.9, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) $\delta = 0.88$ (t, $J = 6.9$ Hz, 3 H), 1.26
15
16 (d, $J = 7.0$ Hz, 3 H), 1.27-1.32 (m, 8 H), 1.37-1.47 (m, 2 H), 1.55 (s, 3 H), 1.64 (s, 3 H), 1.72-
17
18 1.78 (m, 1 H), 2.00-2.08 (m, 1 H), 2.12 (s, 3 H), 2.42 (ddd, $J = 13.3, 9.4, 6.5$ Hz, 1 H), 2.59
19
20 (ddd, $J = 13.3, 6.7, 4.8$ Hz, 1 H), 2.68 (dq, $J = 7.0, 5.1$ Hz, 1 H), 3.32 (d, $J = 7.7$ Hz, 1 H),
21
22 3.57-3.63 (m, 1 H), 3.84 (d, $J = 9.2$ Hz, 1 H), 3.98 (ddd, $J = 9.1, 5.2, 1.5$ Hz, 1 H), 4.14 (ddd,
23
24 $J = 10.6, 5.0, 2.6$ Hz, 1 H); ¹³C NMR (100 MHz, CDCl₃) $\delta = 14.1, 14.2, 15.8, 16.0, 22.6, 22.9,$
25
26 25.8, 26.5, 29.3, 30.9, 31.8, 33.5, 43.1, 57.0, 66.5, 74.8, 95.3, 173.6; FAB HRMS (NBA) *m/e*
27
28 346.2428, M+H⁺ calcd for C₁₈H₃₅NO₃S 346.2416.

29
30
31
32
33
34
35
36
37
38 **Hydroxy Amide 23.** To a suspension of CuI (2.13 g, 11.20 mmol, 4.0 equiv) in THF
39
40 (50 mL) was added dropwise MeLi (1.6 M in Et₂O, 14.0 mL, 22.40 mmol, 8.0 equiv) at 0°C.
41
42 Then, a solution of epoxy amide **21** (1.00 g, 2.80 mmol, 1.0 equiv) in THF (10 mL) was added
43
44 to the resulting colorless solution of Me₂CuLi at 0°C. The reaction mixture was stirred for 8 h
45
46 at this temperature and quenched by careful addition of aqueous saturated NH₄Cl solution,
47
48 followed by dilution with Et₂O. After separation of both phases, the aqueous phase was
49
50 extracted with Et₂O and the combined organic layers were sequentially washed with aqueous
51
52 saturated NH₄Cl solution, water and brine. After treatment with MgSO₄, the solvents were
53
54 removed by reduced pressure to obtain crude 2-methyl-3-hydroxy amide which was subjected
55
56 to purification by flash column chromatography (silica gel, 30% AcOEt in hexanes) to obtain
57
58
59 **23** (1.0 g, 96%) as a colorless oil: $R_f = 0.42$ (silica gel, 33% AcOEt in hexanes); $[\alpha]_D^{25} =$
60

1
2 +19.3° (*c* 1.9, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.88 (t, *J* = 7.1 Hz, 3 H), 1.26 (d, *J* =
3 6.7 Hz, 3 H), 1.23-1.32 (m, 12 H), 1.41-1.48 (m, 1 H), 1.56 (s, 3 H), 1.64 (s, 3 H), 1.72-1.77
4 (m, 1 H), 1.79-1.88 (m, 1 H), 2.00-2.08 (m, 1 H), 2.13 (s, 3 H), 2.42 (ddd, *J* = 15.7, 9.4, 6.5
5 Hz, 1 H), 2.56-2.64 (m, 1 H), 2.68 (dq, *J* = 6.8, 5.1 Hz, 1 H), 3.58-3.62 (m, 1 H), 3.83 (d, *J* =
6 9.2 Hz, 1 H), 3.98 (ddd, *J* = 9.2, 5.2, 1.6 Hz, 1 H), 4.14 (ddd, *J* = 11.2, 5.1, 2.6 Hz, 1 H); ¹³C
7 NMR (100 MHz, CDCl₃) δ = 14.0, 15.7, 15.9, 22.6, 22.8, 25.8, 26.5, 29.2, 29.5, 29.6, 30.9,
8 31.8, 33.5, 35.8, 43.1, 56.9, 66.5, 74.8, 95.3, 173.6; FAB HRMS (NBA) *m/e* 374.2735, M+H⁺
9 calcd for C₂₀H₃₉NO₃S 374.2729.
10
11
12
13
14
15
16
17
18
19
20
21
22

23 **Silylether 24.** A solution of hydroxy amide **22** (2.92 g, 8.44 mmol, 1.0 equiv) in
24 CH₂Cl₂ (20 mL) was treated with tertbutyldimethylsilyl trifluoromethanesulphonate
25 (TBSOTf) (3.9 mL, 16.88 mmol, 2.0 equiv) at 0°C in the presence of 2,6-lutidine (2.5 mL,
26 21.10 mmol, 2.5 equiv). After 0.5 h at 0°C, the reaction mixture was quenched by addition of
27 MeOH (2.0 mL), followed by addition of aqueous saturated NH₄Cl solution and dilution with
28 Et₂O (50 mL). After separation of both phases, the aqueous phase was extracted with Et₂O (2
29 x 30 mL), the combined organic layers were washed with brine and dried with MgSO₄. After
30 filtration, the solvents were removed by reduced pressure to obtain a crude product which was
31 purified by flash column chromatography (silica gel, 20% EtOAc in hexanes) to afford silyl
32 ether **24** (3.70 g, 95%) as a colorless oil: *R_f* = 0.40 (silica gel, 20% AcOEt in hexanes); [α]_D²⁵
33 = -14.3° (*c* 1.6, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = -0.05 (s, 3 H), -0.02 (s, 3 H), 0.80 (s,
34 3 H), 0.83 (t, *J* = 7.1 Hz, 3 H), 0.95 (d, *J* = 6.9 Hz, 3 H), 1.18-1.23 (m, 12 H), 1.36-1.50 (m, 2
35 H), 1.51 (s, 3 H), 1.53 (s, 3 H), 1.62-1.72 (m, 1 H), 1.94-2.02 (m, 1 H), 2.07 (s, 3 H), 2.30-
36 2.41 (m, 1 H), 2.48-2.57 (m, 1 H), 2.68 (dq, *J* = 13.7, 6.9 Hz, 1 H), 3.74 (d, *J* = 9.1 Hz, 1 H),
37 3.84 (ddd, *J* = 9.0, 4.8, 1.5 Hz, 1 H), 3.94 (dt, *J* = 9.0, 3.5 Hz, 1 H), 4.10 (ddd, *J* = 10.8, 4.8,
38 2.3 Hz, 1 H); ¹³C NMR (100 MHz, CDCl₃) δ = -4.9, -4.2, 13.9, 14.8, 17.8, 21.7, 22.5, 23.6,
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 25.8, 26.3, 29.6, 31.8, 32.9, 43.4, 56.4, 66.5, 73.7, 94.7, 172.6; FAB HRMS (NBA) m/e
3
4 460.3308, $M+H^+$ calcd for $C_{24}H_{49}NO_3SSi$ 460.3281.
5
6
7

8
9 **Silylether 25.** A solution of hydroxy amide **23** (350 mg, 0.94 mmol, 1.0 equiv) in
10 CH_2Cl_2 (10 mL) was treated with tertbutyldimethylsilyl trifluoromethanesulphonate
11 (TBSOTf) (0.43 mL, 1.88 mmol, 2.0 equiv) at 0°C in the presence of 2,6-lutidine (0.27 mL,
12 2.35 mmol, 2.5 equiv). After 0.5 h at 0°C, the reaction mixture was quenched by addition of
13 MeOH (0.5 mL), followed by addition of aqueous saturated NH_4Cl solution and dilution with
14 Et_2O (20 mL). After separation of both phases, the aqueous phase was extracted with Et_2O (2
15 x 10 mL), the combined organic layers were washed with brine and dried with $MgSO_4$. After
16 filtration, the solvents were removed by reduced pressure to obtain a crude product which was
17 purified by flash column chromatography (silica gel, 10% EtOAc in hexanes) to afford silyl
18 ether **25** (450 mg, 98%) as a colorless oil: $R_f = 0.28$ (silica gel, 10% AcOEt in hexanes); $[\alpha]_D^{25}$
19 = -9.6° (c 0.8, CH_2Cl_2); 1H NMR (400 MHz, $CDCl_3$) δ = 0.04 (s, 3 H), 0.05 (s, 3 H), 0.86 (s, 9
20 H), 0.87 (t, $J = 6.8$ Hz, 3 H), 1.00 (d, $J = 6.8$ Hz, 3 H), 1.21-1.32 (m, 12 H), 1.42-1.53 (m, 2
21 H), 1.57 (s, 3 H), 1.58 (s, 3 H), 1.70-1.78 (m, 1 H), 1.99-2.08 (m, 1 H), 2.12 (s, 3 H), 2.43
22 (ddd, $J = 13.3, 9.0, 6.9$ Hz, 1 H), 2.58 (ddd, $J = 12.3, 6.8, 5.0$ Hz, 1 H), 2.74 (dq, $J = 8.9, 6.8$
23 Hz, 1 H), 3.80 (d, $J = 9.0$ Hz, 1 H), 3.89 (ddd, $J = 9.0, 4.8, 1.3$ Hz, 1 H), 3.99 (dt, $J = 9.0, 3.3$
24 Hz, 1 H), 4.16 (ddd, $J = 10.7, 4.6, 2.2$ Hz, 1 H); ^{13}C NMR (100 MHz, $CDCl_3$) δ = -4.7, -4.1,
25 14.1, 14.9, 15.9, 17.9, 21.9, 22.7, 23.8, 25.9, 26.5, 29.3, 29.7, 30.1, 31.2, 31.9, 33.1, 33.3,
26 43.6, 56.6, 66.7, 73.9, 94.8, 172.8; FAB HRMS (NBA) m/e 488.3605, $M+H^+$ calcd for
27 $C_{26}H_{53}NO_3SSi$ 488.3594.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54
55
56 **Diol 26 from Hydroxy Amide 22.** A freshly prepared solution of LDA
57 [Diisopropylamine (0.41 mL, 2.90 mmol, 10.0 equiv) was added to a solution of n -BuLi (1.6
58 M solution in hexanes, 1.8 mL, 2.90 mmol, 10.0 equiv) in THF (5.0 mL) at 0°C] was treated
59
60

1
2 with borane-ammonia complex (90 mg, 2.90 mmol, 10.0 equiv) at 0°C. After 15 min at this
3
4 temperature, the resulting suspension was stirred at room temperature for additional 30 min
5
6 before cooling again at 0°C. Then, a solution of hydroxy amide **22** (100 mg, 0.29 mmol, 1.0
7
8 equiv) in THF (3.0 mL) was added. The reaction mixture was warmed to room temperature
9
10 and then heated under reflux for 8 h. After this time, reaction mixture was allowed to warm to
11
12 room temperature and hydride excess was quenched by careful addition of MeOH (2.0 mL).
13
14 The crude mixture was then diluted with Et₂O and treated with a saturated aqueous NH₄Cl
15
16 solution. The organic phase was separated, and the aqueous phase was extracted with Et₂O
17
18 (twice). The combined organic solution was sequentially washed with water and brine, dried
19
20 (MgSO₄), filtered and concentrated under reduced pressure. The resulting crude product was
21
22 purified by flash column chromatography (silica gel, 20% → 50% AcOEt in hexanes) to obtain
23
24 to obtain diol **26** (45 mg, 90%) as a colorless oil: *R_f* = 0.45 (silica gel, 50% AcOEt in
25
26 hexanes); [α]_D²⁵ = +16.1° (*c* 0.7, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.87 (t, *J* = 7.0 Hz,
27
28 3 H), 0.88 (d, *J* = 6.6 Hz, 3 H), 1.24-1.35 (m, 7 H), 1.39-1.58 (m, 3 H), 1.67 (dsext, *J* = 7.2, 3.7
29
30 Hz, 1 H), 3.29 (s, 2 H), 3.53 (dt, *J* = 7.8, 3.2 Hz, 1 H), 3.59 (dd, *J* = 10.8, 7.3 Hz, 1 H), 3.75
31
32 (dd, *J* = 10.8, 3.7 Hz, 1 H); ¹³C NMR (100 MHz, CDCl₃) δ = 13.9, 14.1, 22.6, 25.1, 29.4, 31.8,
33
34 35.3, 39.7, 67.6, 77.3; FAB HRMS (NBA) *m/e* 175.1708, M+H⁺ calcd for C₁₀H₂₂O₂ 175.1698.
35
36
37
38
39
40
41
42
43
44

45 **Diol 26 from Silyl Ether 24.** A freshly prepared solution of LDA [Diisopropylamine
46
47 (0.44 mL, 3.2 mmol, 4.0 equiv) was added to a solution of *n*-BuLi (1.6 M solution in hexanes,
48
49 2.0 mL, 3.2 mmol, 4.0 equiv) in THF (5.0 mL) at 0°C] was treated with borane-ammonia
50
51 complex (100 mg, 3.2 mmol, 4.0 equiv) at 0°C. After 15 min at this temperature, the resulting
52
53 suspension was stirred at room temperature for additional 30 min before cooling again at 0°C.
54
55 Then, a solution of the silylether **24** (367 mg, 0.80 mmol, 1.0 equiv) in THF (5.0 mL) was
56
57 added. The reaction mixture was warmed to room temperature and then heated under reflux
58
59 for 4 h. After this time, reaction mixture was allowed to warm to room temperature and
60

1
2 worked out as described in Procedure A to obtain a crude product that was purified by flash
3
4 column chromatography (silica gel, 50% AcOEt in hexanes) to obtain diol **26** (119 mg, 85%)
5
6 that exhibited identical physical and spectroscopic properties than the obtained from **22** as
7
8 described above.
9

10
11
12
13 **Diol 27 from Hydroxy Amide 23.** A freshly prepared solution of LDA
14 [Diisopropylamine (0.36 mL, 2.60 mmol, 10.0 equiv) was added to a solution of *n*-BuLi (1.6
15 M solution in hexanes, 1.6 mL, 2.60 mmol, 10.0 equiv) in THF (5.0 mL) at 0°C] was treated
16 with borane-ammonia complex (80 mg, 2.60 mmol, 10.0 equiv) at 0°C. After 15 min at this
17 temperature, the resulting suspension was stirred at room temperature for additional 30 min
18 before cooling again at 0°C. Then, a solution of hydroxy amide **23** (97 mg, 0.26 mmol, 1.0
19 equiv) in THF (3.0 mL) was added. The reaction mixture was warmed to room temperature
20 and then heated under reflux for 8 h. After this time, reaction mixture was allowed to warm to
21 room temperature and hydride excess was quenched by careful addition of MeOH (2.0 mL).
22 The crude mixture was then diluted with Et₂O and treated with a saturated aqueous NH₄Cl
23 solution. The organic phase was separated, and the aqueous phase was extracted with Et₂O
24 (twice). The combined organic solution was sequentially washed with water and brine, dried
25 (MgSO₄), filtered and concentrated under reduced pressure. The resulting crude product was
26 purified by flash column chromatography (silica gel, 20% → 50% AcOEt in hexanes) to obtain
27 to obtain diol **27** (49 mg, 93%) as a colorless oil: *R_f* = 0.38 (silica gel, 50% AcOEt in
28 hexanes); $[\alpha]_D^{25} = +22.7^\circ$ (*c* 1.1, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.88 (t, *J* = 7.1 Hz,
29 3 H), 0.89 (d, *J* = 7.0 Hz, 3 H), 1.22-1.39 (m, 12 H), 1.41-1.51 (m, 1 H), 1.53-1.58 (m, 1 H),
30 1.71 (dsext, *J* = 7.1, 3.8 Hz, 1 H), 2.59 (bs, 1 H), 2.83 (bs, 1 H), 3.55 (dt, *J* = 7.6, 3.2 Hz, 1 H),
31 3.62 (dd, *J* = 10.9, 7.1 Hz, 1 H), 3.76 (dd, *J* = 10.9, 3.7 Hz, 1 H); ¹³C NMR (100 MHz, CDCl₃)
32 δ = 13.9, 14.1, 22.6, 25.2, 29.3, 29.6, 29.7, 31.9, 35.4, 39.9, 67.6, 77.4; FAB HRMS (NBA)
33 *m/e* 203.2015, M+H⁺ calcd for C₁₂H₂₆O₂ 203.2011.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Diol 27 from Silyl Ether 25. A freshly prepared solution of LDA [Diisopropylamine (0.24 mL, 1.72 mmol, 4.0 equiv) was added to a solution of *n*-BuLi (1.6 M solution in hexanes, 1.1 mL, 1.72 mmol, 4.0 equiv) in THF (4.0 mL) at 0°C] was treated with borane-ammonia complex (53 mg, 1.72 mmol, 4.0 equiv) at 0°C. After 15 min at this temperature, the resulting suspension was stirred at room temperature for additional 30 min before cooling again at 0°C. Then, a solution of the silylether **25** (209 mg, 0.43 mmol, 1.0 equiv) in THF (4.0 mL) was added. The reaction mixture was warmed to room temperature and then heated under reflux for 4 h. After this time, reaction mixture was allowed to warm to room temperature and worked out as described in Procedure A to obtain a crude product that was purified by flash column chromatography (silica gel, 50% AcOEt in hexanes) to obtain diol **27** (80 mg, 92%) that exhibited identical physical and spectroscopic properties than the obtained from **23** as described above.

Hydroxy Acid 9. Diol **26** (100 mg, 0.57 mmol) was dissolved in a mixture of CH₃CN/H₂O (8.0 mL, 1/1) and the resulting solution treated with BAIB (918 mg, 2.85 mmol, 5.0 equiv) followed by TEMPO (44 mg, 0.29 mmol, 0.5 equiv) at 25°C. After 6 h, the crude mixture was diluted with AcOEt, quenched by the addition of a saturated aqueous Na₂S₂O₃ solution and, after separation of both layers, the aqueous phase was then extracted with AcOEt. The combined organic solution was washed with saturated aqueous Na₂S₂O₃ solution again, dried over anhydrous MgSO₄ and the solvent evaporated under reduced pressure. Purification of the crude acid by flash column chromatography (silica gel, 3%→5% MeOH in CH₂Cl₂) provided hydroxy acid **9** (77 mg, 72%) as a colorless oil: *R*_f = 0.25 (silica gel, 5% MeOH in CH₂Cl₂); [α]_D²⁵ = +5.0° (*c* 0.54, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.86 (t, *J* = 6.8 Hz, 3 H), 1.20-1.37 (m, 7 H), 1.23 (d, *J* = 7.2 Hz, 3 H), 1.40-1.57 (m, 3 H), 2.54 (quint, *J* = 6.9 Hz, 1 H), 3.64-3.70 (m, 1 H); ¹³C NMR (100 MHz, CDCl₃) δ = 14.0, 14.2, 22.6, 25.4,

1
2 29.2, 31.8, 34.6, 45.2, 67.6, 77.3, 180.8; FAB HRMS (NBA) m/e 189.1504, $M+H^+$ calcd for
3
4 $C_{10}H_{20}O_3$ 189.1491.
5
6
7

8
9 **Hydroxy Acid 10.** Diol **27** (70 mg, 0.37 mmol) was dissolved in a mixture of
10
11 CH_3CN/H_2O (6.0 mL, 1/1) and the resulting solution treated with BAIB (602 mg, 1.84 mmol,
12
13 5.0 equiv) followed by TEMPO (30 mg, 0.18 mmol, 0.5 equiv) at 25°C. After 6 h, the crude
14
15 mixture was diluted with AcOEt, quenched by the addition of a saturated aqueous $Na_2S_2O_3$
16
17 solution and, after separation of both layers, the aqueous phase was then extracted with
18
19 AcOEt. The combined organic solution was washed with saturated aqueous $Na_2S_2O_3$ solution
20
21 again, dried over anhydrous $MgSO_4$ and the solvent evaporated under reduced pressure.
22
23 Purification of the crude acid by flash column chromatography (silica gel, 3% → 5% MeOH in
24
25 CH_2Cl_2) provided hydroxy acid **10** (49 mg, 65%) as a colorless oil: $R_f = 0.29$ (silica gel, 5%
26
27 MeOH in CH_2Cl_2); $[\alpha]_D^{25} = +1.9^\circ$ (c 2.5, CH_2Cl_2); 1H NMR (400 MHz, $CDCl_3$) $\delta = 0.88$ (t, $J =$
28
29 6.9 Hz, 3 H), 1.24 (d, $J = 7.2$ Hz, 3 H), 1.25-1.39 (m, 11 H), 1.42-1.57 (m, 3 H), 2.56 (quint, J
30
31 = 6.9 Hz, 1 H), 3.67-3.72 (m, 1 H), 6.22 (bs, 1 H); ^{13}C NMR (100 MHz, $CDCl_3$) $\delta = 14.1,$
32
33 14.2, 22.6, 25.4, 29.2, 29.5, 31.8, 34.6, 45.2, 77.3, 180.9; FAB HRMS (NBA) m/e 217.1824,
34
35 $M+H^+$ calcd for $C_{12}H_{24}O_3$ 217.1804.
36
37
38
39
40
41
42
43

44 **Epoxy Alcohols 30 and 31. General Procedure.** To a solution of titanium
45
46 tetraisopropoxide (0.20 equiv) and 4Å molecular sieves in CH_2Cl_2 (0.1 M) was added (-)-DET
47
48 (0.25 equiv) at -23°C. After 15 min at this temperature, a solution of allylic alcohols **28** or **29**
49
50 (1.0 equiv) in CH_2Cl_2 (1.0 M) was added dropwise, followed by the addition, after additional
51
52 30 min, of TBHP (5.5 M solution in decane, 1.5 equiv) at -23°C. After 8 h at this temperature,
53
54 the reaction mixture was filtered and the filtrate was diluted with EtOAc and washed with a
55
56 saturated aqueous solution of sodium sulphate. After decantation, the aqueous phase was
57
58 extracted with EtOAc (3 times), and the combined organic layer was dried ($MgSO_4$), filtered
59
60

1
2 and concentrated. The resulting crude products were purified by flash column chromatography
3
4 (silica gel, 20% AcOEt in hexanes for both cases) to obtain epoxy alcohols **30** and **31** (75 and
5
6 70% yields respectively). [**30**]: Described in the literature²⁴; white solid; m. p. 45 °C; $R_f = 0.39$
7
8 (silica gel, 30% AcOEt in hexanes); $[\alpha]_D^{25} = +38.6^\circ$ (c 1.1, CH_2Cl_2); ^1H NMR (400 MHz,
9
10 CDCl_3) $\delta = 0.88$ (t, $J = 7.0$ Hz, 3 H), 1.25-1.37 (m, 6 H), 1.41-1.46 (m, 2 H), 1.54-1.59 (m, 2
11
12 H), 1.89 (dd, $J = 6.5, 6.1$ Hz, 1 H), 2.92 (dt, $J = 4.9, 2.5$ Hz, 1 H), 2.95 (dt, $J = 5.7, 2.5$ Hz, 1
13
14 H), 3.61 (ddd, $J = 11.5, 7.0, 4.4$ Hz, 1 H), 3.90 (ddd, $J = 12.6, 5.3, 2.6$ Hz, 1 H); ^{13}C NMR
15
16 (100 MHz, CDCl_3) $\delta = 13.8, 22.4, 25.8, 28.9, 31.5, 31.6, 56.0, 58.7, 61.9$. [**31**]: colorless oil;
17
18 $R_f = 0.45$ (silica gel, 20% AcOEt in hexanes); $[\alpha]_D^{25} = +16.2^\circ$ (c 2.3, CH_2Cl_2); ^1H NMR (400
19
20 MHz, CDCl_3) $\delta = 0.80$ (t, $J = 6.8$ Hz, 3 H), 1.19-1.21 (m, 11 H), 1.41-1.51 (m, 2 H), 2.27 (bs,
21
22 1 H), 2.93 (t, $J = 6.1$ Hz, 1 H), 3.46 (dd, $J = 12.1, 6.7$ Hz, 1 H), 3.59 (d, $J = 11.8$ Hz, 1 H); ^{13}C
23
24 NMR (100 MHz, CDCl_3) $\delta = 14.0, 14.1, 22.5, 26.3, 28.1, 29.0, 31.7, 60.3, 61.0, 65.4$; FAB
25
26 HRMS (NBA) m/e 173.1557, $\text{M}+\text{H}^+$ calcd for $\text{C}_{10}\text{H}_{20}\text{O}_2$ 173.1542.
27
28
29
30
31
32
33
34

35 **Epoxy Alcohol 30 from epoxyamide 20.** To a solution of epoxyamide **20** (85 mg,
36
37 0.26 mmol, 1.0 equiv) in THF (4.0 mL) was added lithium triethylborohydride (Super-H) (1.0
38
39 M solution in THF, 0.78 mL, 0.78 mmol, 3.0 equiv) at 0°C. After 15 min at this temperature,
40
41 the excess of Super-H was carefully quenched by addition of MeOH (0.5 mL), the resulting
42
43 mixture diluted with Et_2O (5 mL) and washed with a saturated aqueous NH_4Cl solution (8
44
45 mL). After decantation, the aqueous phase was extracted with Et_2O (2 x 10 mL), and the
46
47 combined organic extracts were washed with brine (10 mL), dried (MgSO_4) and filtered.
48
49 Concentration under reduced pressure provided a crude product that was purified by flash
50
51 column chromatography (silica gel, 20% AcOEt in hexanes) to afford epoxy alcohol **30** (35
52
53 mg, 85% yield) whose physical and spectroscopic data were identical to those described in the
54
55 literature and completely matched with the one obtained from Sharpless asymmetric
56
57 epoxidation.
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Diol 26 from Epoxy Alcohol 30. To a suspension of CuI (4.57 g, 24.0 mmol, 3.0 equiv) in THF (50 mL) was added dropwise MeLi (1.6 M in Et₂O, 30.0 mL, 48.0 mmol, 6.0 equiv) at 0°C. Then, a solution of epoxy alcohol **30** (1.26 g, 8.0 mmol, 1.0 equiv) in THF (10 mL) was added to the resulting colorless solution of Me₂CuLi at 0°C. The reaction mixture was stirred for 8 h at this temperature and quenched by careful addition of aqueous saturated NH₄Cl solution, followed by dilution with Et₂O. After separation of both phases, the aqueous phase was extracted with Et₂O and the combined organic layers were sequentially washed with aqueous saturated NH₄Cl solution, water and brine. After treatment with MgSO₄, the solvents were removed by reduced pressure to obtain a crude consisting of a mixture of 2- and 3-Methyl opening products in a 3:1 proportion which was subjected to the next step without purification. Thus, crude diol (1.39 g, 8.0 mmol, 1.0 equiv) in THF (30 mL) and water (30 mL) was treated with NaIO₄ (855 mg, 4.0 mmol, 0.5 equiv) at room temperature. After vigorous stirring for 18 h, the reaction mixture was diluted with Et₂O and water. After separation of both phases, the aqueous phase was extracted with Et₂O (3 x 15 mL) and the combined organic layers were washed with brine. After treatment with MgSO₄, the solvents were removed by reduced pressure and the crude product purified by flash column chromatography (silica gel, 50% AcOEt in hexanes) to obtain the desired 1,3-diol **26** (571 mg, 41%) as a colorless oil and with identical physical and spectroscopic properties to those reported above.

51
52
53
54
55
56
57
58
59
60

Hydroxy ester 33. To a solution of epoxy alcohol **31** (1.17 g, 6.80 mmol) in CH₂Cl₂ (20 mL) was sequentially added DMSO (8.0 mL), TEA (4.7 mL, 34.0 mmol, 5.0 equiv) and pyr•SO₃ complex (2.2 g, 13.60 mmol, 2.0 equiv) at 0°C. After stirring for 2 h, the reaction mixture was diluted with Et₂O, and a saturated aqueous NH₄Cl solution added. The organic phase was separated, and the aqueous phase was extracted with Et₂O (twice). The combined

1
2 organic solution was sequentially washed with water and brine, dried (MgSO₄), filtered and
3
4 concentrated under reduced pressure. The resulting aldehyde was used in the next step without
5
6 further purification. To a solution of catalyst **32** (163 mg, 0.68 mmol, 0.1 equiv) in CH₂Cl₂
7
8 (10 mL) was added crude aldehyde (~ 6.80 mmol), followed by EtOH (1.2 mL, 20.4 mmol,
9
10 3.0 equiv) and DIPEA (0.1 mL, 0.54 mmol, 0.08 equiv). The resulting yellow solution was
11
12 stirred at 25°C overnight after which, TLC analysis showed depletion of starting material. The
13
14 reaction mixture was then treated with a saturated aqueous NH₄Cl solution and diluted with
15
16 Et₂O. The organic phase was separated, and the aqueous phase was extracted with Et₂O
17
18 (twice). The combined organic solution was sequentially washed with water and brine, dried
19
20 (MgSO₄), filtered and concentrated under reduced pressure. The resulting crude product was
21
22 purified by flash column chromatography (silica gel, 20% AcOEt in hexanes) to obtain an
23
24 inseparable mixture of hydroxy esters **33** and its C-2 epimer (1.0 g, 70% combined overall
25
26 yield from epoxy alcohol **31**, dr 7:3 *anti:syn*) as a colorless oil: ¹H NMR (400 MHz, CDCl₃)
27
28 (major isomer) δ = 0.80 (t, *J* = 7.0 Hz, 3 H), 1.12 (d, *J* = 7.1 Hz, 3 H), 1.19 (t, *J* = 7.3 Hz, 3
29
30 H), 1.17-1.24 (m, 8 H), 1.32-1.42 (m, 2 H), 2.42 (quint, *J* = 7.1 Hz, 1 H), 3.57 (dt, *J* = 6.7, 3.3
31
32 Hz, 1 H), 4.08 (q, *J* = 7.3 Hz, 2 H); ¹³C NMR (100 MHz, CDCl₃) (major isomer) δ = 14.0,
33
34 14.1, 14.3, 22.5, 25.4, 29.2, 31.7, 34.7, 45.1, 60.5, 73.3, 176.1; FAB HRMS (NBA) *m/e*
35
36 217.1815, M+H⁺ calcd for C₁₂H₂₄O₃ 217.1804.
37
38
39
40
41
42
43
44
45
46

47 **Acid 9 from Hydroxy Ester 33.** A solution of hydroxy ester **33** (450 mg, 2.1 mmol)
48
49 in THF (10 mL) was treated with a 1.0 M aqueous LiOH solution (4.2 mL, 4.2 mmol, 2.0
50
51 equiv) at 0°C. After stirring for 8 h at 25°C, the reaction mixture was treated with a 2.0 M
52
53 aqueous HCl solution until pH 2. Then, the crude mixture was diluted with EtOAc and water,
54
55 both phases separated and the aqueous phase extracted with AcOEt (3 x 15 mL). The
56
57 combined organic extracts were sequentially washed with water and brine, dried over MgSO₄,
58
59 filtered and concentrated by reduced pressure. The crude product was purified by flash
60

1
2 column chromatography (silica gel, 3%→5% MeOH in CH₂Cl₂) to provide hydroxy acid **9**
3
4 contaminated with its 2-epimer (365 mg, 92% combined yield in a 7:3 proportion) as a
5
6 colorless oil whose major isomer exhibited identical spectroscopic properties to those reported
7
8
9 above.

10
11
12
13
14 **Dihydroxy Amide 34.** A solution of hydroxy amide **22** (1.86 g, 5.39 mmol) in
15
16 dioxane (30 mL) was treated with a 2.7 M aqueous HCl solution (15 mL) at 25°C. After
17
18 stirring for 8 h at this temperature, a white solid was formed in the reaction mixture, which
19
20 was filtered and washed with cold water (three times) and Et₂O (three times). After drying
21
22 under high vacuum, it is obtained a white solid that corresponded to the titled compound **34**
23
24 (1.60 g, 98%) which did not require further purification: $R_f = 0.35$ (silica gel, 5% MeOH in
25
26 CH₂Cl₂); m. p. 140-143°C; $[\alpha]_D^{25} = -25.3^\circ$ (*c* 0.9, DMSO); ¹H NMR (400 MHz, DMSO-*d*₆) $\delta =$
27
28 0.86 (t, *J* = 6.8 Hz, 3 H), 0.96 (d, *J* = 7.0 Hz, 3 H), 1.21-1.27 (m, 8 H), 1.35-1.40 (m, 2 H),
29
30 1.53 (ddt, *J* = 14.6, 9.7, 5.0 Hz, 1 H), 1.76-1.84 (m, 1 H), 2.02 (s, 3 H), 2.25 (quint, *J* = 6.9
31
32 Hz, 1 H), 2.37 (ddd, *J* = 12.9, 9.7, 6.5 Hz, 1 H), 2.48 (ddd, *J* = 12.9, 10.0, 5.0 Hz, 1 H), 3.23
33
34 (dt, *J* = 10.7, 6.2 Hz, 1 H), 3.35 (dt, *J* = 10.5, 5.2 Hz, 1 H), 3.44 (bq, *J* = 6.7 Hz, 1 H), 3.74-
35
36 3.83 (m, 1 H), 4.45 (d, *J* = 6.6 Hz, 1 H), 4.65 (t, *J* = 5.6 Hz, 1 H), 7.47 (d, *J* = 8.6 Hz, 1 H);
37
38 ¹³C NMR (100 MHz, DMSO-*d*₆) $\delta = 13.9, 14.1, 14.6, 22.0, 25.0, 28.8, 29.9, 30.8, 31.3, 34.1,$
39
40 45.5, 49.6, 63.2, 71.9, 174.7; FAB HRMS (NBA) *m/e* 306.2095, M+H⁺ calcd for C₁₅H₃₁NO₃S
41
42 306.2103.
43
44
45
46
47
48
49
50
51

52 **Bis(silyl ether) 36.** A solution of dihydroxy amide **34** (562 mg, 1.84 mmol, 1.0 equiv)
53
54 in CH₂Cl₂ (10 mL) was treated with tertbutyldimethylsilyl trifluoromethanesulfonate
55
56 (TBSOTf) (1.05 mL, 4.60 mmol, 2.5 equiv) at 0°C in the presence of 2,6-lutidine (0.64 mL,
57
58 5.52 mmol, 3.0 equiv). After 1.0 h at 0°C, the reaction mixture was quenched by addition of
59
60 MeOH (1.0 mL), followed by addition of aqueous saturated NH₄Cl solution (15 mL) and

1
2 dilution with Et₂O (20 mL). After separation of both phases, the aqueous phase was extracted
3
4 with Et₂O (2 x 15 mL), and the combined organic layers were washed with brine and dried
5
6 with MgSO₄. After filtration, the solvents were removed under reduced pressure to obtain a
7
8 crude product which was purified by flash column chromatography (silica gel, 10% EtOAc in
9
10 hexanes) to afford bis(silyl ether) **36** (956 mg, 97%) as a colorless oil: *R_f* = 0.54 (silica gel,
11
12 10% AcOEt in hexanes); [α]_D²⁵ = -29.7° (*c* 0.9, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.04
13
14 (s, 3 H), 0.05 (s, 3 H), 0.08 (s, 3 H), 0.09 (s, 3 H), 0.87 (t, *J* = 7.0 Hz, 3 H), 0.89 (s, 9 H), 0.92
15
16 (s, 9 H), 1.21 (d, *J* = 7.3 Hz, 3 H), 1.24-1.27 (m, 8 H), 1.48-1.62 (m, 2 H), 1.66-1.75 (m, 1 H),
17
18 1.83-1.92 (m, 1 H), 2.09 (s, 3 H), 2.45 (dq, *J* = 7.3, 3.4 Hz, 1 H), 2.47-2.51 (m, 2 H), 3.61 (d,
19
20 *J* = 4.1 Hz, 2 H), 3.70 (ddd, *J* = 8.0, 5.4, 3.4 Hz, 1 H), 4.02-4.08 (m, 1 H), 6.75 (d, *J* = 8.9 Hz,
21
22 1 H); ¹³C NMR (100 MHz, CDCl₃) δ = -5.4, -4.8, -4.2, 14.0, 15.5, 16.5, 18.0, 18.3, 22.6, 25.7,
23
24 25.9, 26.0, 29.4, 30.9, 31.3, 31.7, 35.6, 45.9, 49.7, 64.9, 74.7, 174.7; FAB HRMS (NBA) *m/e*
25
26 534.3845, M+H⁺ calcd for C₂₇H₅₉NO₃SSi₂ 534.3832.
27
28
29
30
31
32
33
34

35 **Acid 42.** To a stirred solution of bis(silyl ether) **36** (52 mg, 0.097 mmol, 1.0 equiv) in
36
37 a mixture of Ac₂O (4 mL) and AcOH (2 mL) was portionwise added NaNO₂ (140 mg, 2.02
38
39 mmol, 20.0 equiv) at 0°C. The reaction mixture was stirred for 4 h at this temperature, after
40
41 which, was diluted with cold water (5 mL) and extracted with Et₂O (3 x 10 mL). Then, the
42
43 combined organic solution was washed with a 5% aqueous NaHCO₃ solution several times
44
45 until removal of acetic acid was complete (~ 10 x 10 mL). Finally, the ethereal solution was
46
47 washed with brine (20 mL), dried with MgSO₄, filtered and concentrated under reduced
48
49 pressure. Purification by flash column chromatography (silica gel, CH₂Cl₂ → 1% MeOH in
50
51 CH₂Cl₂) furnished acid derivative **42** (22 mg, 75%) as a yellow oil: *R_f* = 0.43 (silica gel, 1%
52
53 MeOH in CH₂Cl₂); [α]_D²⁵ = -2.7° (*c* 0.7, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.10 (s, 3
54
55 H), 0.11 (s, 3 H), 0.88 (t, *J* = 7.0 Hz, 3 H), 0.91 (s, 9 H), 1.21 (d, *J* = 7.2 Hz, 3 H), 1.24-1.33
56
57 (m, 8 H), 1.45-1.59 (m, 2 H), 2.67 (dq, *J* = 7.3, 4.3 Hz, 1 H), 3.85 (ddd, *J* = 5.4, 4.3 Hz, 1 H);
58
59
60

¹³C NMR (100 MHz, CDCl₃) δ = -4.9, -4.4, 14.1, 18.0, 22.6, 24.8, 25.7, 29.3, 31.7, 34.7, 44.7, 74.4, 177.7; FAB HRMS (NBA) *m/e* 303.2352, M+H⁺ calcd for C₁₆H₃₄O₃Si 303.2356.

Acid 9 from Silyloxy Acid 42. A solution of silyloxy acid **42** (13 mg, 0.043 mmol) in THF (3 mL) was treated with TBAF (1.0 M in THF, 65 μ L, 0.065 mmol, 1.5 equiv) at 0°C. After 2.0 h at 0°C, the reaction mixture was diluted with EtOAc (5 mL) and washed with a saturated aqueous NH₄Cl solution (5 mL). After separation of both phases, the aqueous phase was extracted with AcOEt (3 x 5 mL), and the combined organic layers were washed with brine and dried with MgSO₄. After filtration, the solvents were removed under reduced pressure to obtain a crude product which was purified by flash column chromatography (silica gel, 3%→5% MeOH in CH₂Cl₂) to provide hydroxy acid **9** (7.8 mg, 95%) which exhibited identical spectroscopic properties to those reported above.

Dihydroxy Amide 44. A solution of hydroxy amide **23** (44 mg, 0.118 mmol) in dioxane (5 mL) was treated with a 2.7 M aqueous HCl solution (2.5 mL) at 25°C. After stirring for 8 h at this temperature, a white solid was formed in the reaction mixture, which was filtered and washed with cold water (three times) and Et₂O (three times). After drying under high vacuum, it is obtained a white solid that corresponded to the titled compound **44** (36 mg, 91%) which did not require further purification: *R_f* = 0.37 (silica gel, 5% MeOH in CH₂Cl₂); m. p. 158-160°C; ¹H NMR (400 MHz, CDCl₃) δ = 0.88 (t, *J* = 6.7 Hz, 3 H), 1.25 (d, *J* = 7.1 Hz, 3 H), 1.26-1.32 (m, 12 H), 1.45-1.50 (m, 2 H), 1.81-1.96 (m, 4 H), 2.12 (s, 3 H), 2.29 (dq, *J* = 7.1, 5.4 Hz, 1 H), 2.48-2.59 (m, 2 H), 3.59-3.64 (m, 1 H), 3.65 (dd, *J* = 11.2, 5.2 Hz, 1 H), 3.71 (dd, *J* = 11.2, 3.6 Hz, 1 H), 4.02-4.09 (m, 1 H), 6.15 (d, *J* = 7.0 Hz, 1 H); ¹³C NMR (100 MHz, CDCl₃) δ = 14.1, 15.6, 15.7, 22.7, 25.8, 29.3, 29.5, 29.6, 30.4, 30.8, 31.9, 35.5, 46.5, 51.1, 65.1, 74.0, 176.6; FAB HRMS (NBA) *m/e* 334.2425, M+H⁺ calcd for C₁₇H₃₅NO₃S 334.2416.

Bis(silyl ether) 45. A solution of dihydroxy amide **44** (33 mg, 0.099 mmol, 1.0 equiv) in CH₂Cl₂ (3 mL) was treated with tertbutyldimethylsilyl trifluoromethanesulfonate (TBSOTf) (57 μL, 0.247 mmol, 2.5 equiv) at 0°C in the presence of 2,6-lutidine (35 μL, 0.30 mmol, 3.0 equiv). After 1.0 h at 0°C, the reaction mixture was quenched by addition of MeOH (0.2 mL), followed by addition of aqueous saturated NH₄Cl solution (10 mL) and dilution with Et₂O (10 mL). After separation of both phases, the aqueous phase was extracted with Et₂O (2 x 5 mL), and the combined organic layers were washed with brine and dried with MgSO₄. After filtration, the solvents were removed under reduced pressure to obtain a crude product which was purified by flash column chromatography (silica gel, 10% EtOAc in hexanes) to afford bis(silyl ether) **45** (55 mg, 98%) as a colorless oil: *R*_f = 0.59 (silica gel, 10% AcOEt in hexanes); [α]_D²⁵ = -30.4° (*c* 2.6, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.036 (s, 3 H), 0.042 (s, 3 H), 0.085 (s, 3 H), 0.089 (s, 3 H), 0.85 (t, *J* = 7.0 Hz, 3 H), 0.87 (s, 9 H), 0.92 (s, 9 H), 1.21 (d, *J* = 7.2 Hz, 3 H), 1.22-1.27 (m, 12 H), 1.46-1.61 (m, 2 H), 1.66-1.75 (m, 1 H), 1.83-1.92 (m, 1 H), 2.08 (s, 3 H), 2.45 (dq, *J* = 7.3, 3.3 Hz, 1 H), 2.46-2.51 (m, 2 H), 3.60 (d, *J* = 4.1 Hz, 2 H), 3.69 (ddd, *J* = 8.3, 5.3, 3.3 Hz, 1 H), 4.02-4.08 (m, 1 H), 6.76 (d, *J* = 8.9 Hz, 1 H); ¹³C NMR (100 MHz, CDCl₃) δ = -5.4, -4.8, -4.2, 14.07, 14.09, 15.5, 16.5, 18.0, 18.3, 22.6, 25.7, 25.9, 26.0, 29.2, 29.5, 29.7, 30.9, 31.3, 31.6, 31.8, 35.6, 45.9, 49.7, 64.9, 74.7, 174.8; FAB HRMS (NBA) *m/e* 562.4149, M+H⁺ calcd for C₂₉H₆₃NO₃SSi₂ 562.4145.

Acid 46. To a stirred solution of bis(silyl ether) **45** (51 mg, 0.091 mmol, 1.0 equiv) in a mixture of Ac₂O (4 mL) and AcOH (2 mL) was portionwise added NaNO₂ (125 mg, 1.81 mmol, 20.0 equiv) at 0°C. The reaction mixture was stirred for 4 h at this temperature, after which, was diluted with cold water (5 mL) and extracted with Et₂O (3 x 10 mL). Then, the combined organic solution was washed with a 5% aqueous NaHCO₃ solution several times until removal of acetic acid was complete (~ 10 x 10 mL). Finally, the ethereal solution was washed with brine (20 mL), dried with MgSO₄, filtered and concentrated under reduced

1
2 pressure. Purification by flash column chromatography (silica gel, CH₂Cl₂ → 1% MeOH in
3 CH₂Cl₂) furnished acid derivative **46** (20 mg, 65%) as a yellow oil: *R_f* = 0.47 (silica gel, 1%
4 CH₂Cl₂); [α]_D²⁵ = -3.5° (*c* 0.4, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃) δ = 0.11 (s, 3
5 H), 0.12 (s, 3 H), 0.88 (t, *J* = 7.2 Hz, 3 H), 0.91 (s, 9 H), 1.21 (d, *J* = 7.2 Hz, 3 H), 1.25-1.28
6 (m, 12 H), 1.47-1.59 (m, 2 H), 2.67 (dq, *J* = 7.1, 4.2 Hz, 1 H), 3.84 (ddd, *J* = 5.6, 4.4 Hz, 1 H);
7 ¹³C NMR (100 MHz, CDCl₃) δ = -4.9, -4.4, 14.1, 18.0, 22.6, 24.8, 25.7, 29.2, 29.4, 29.6, 31.8,
8 34.8, 44.5, 74.4, 177.3; FAB HRMS (NBA) *m/e* 331.2674, M+H⁺ calcd for C₁₈H₃₇O₃Si
9 331.2668.
10
11
12
13
14
15
16
17
18
19
20
21
22

23 **Acid 10 from Silyloxy Acid 46.** A solution of silyloxy acid **46** (18 mg, 0.054 mmol)
24 in THF (3 mL) was treated with TBAF (1.0 M in THF, 82 μL, 0.082 mmol, 1.5 equiv) at 0°C.
25 After 2.0 h at 0°C, the reaction mixture was diluted with EtOAc (5 mL) and washed with a
26 saturated aqueous NH₄Cl solution (5 mL). After separation of both phases, the aqueous phase
27 was extracted with AcOEt (3 x 5 mL), and the combined organic layers were washed with
28 brine and dried with MgSO₄. After filtration, the solvents were removed under reduced
29 pressure to obtain a crude product which was purified by flash column chromatography (silica
30 gel, 3%→5% MeOH in CH₂Cl₂) to provide hydroxy acid **10** (11.5 mg, 98%) which exhibited
31 identical spectroscopic properties to those reported above.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 **Polystyrene-bound pentapeptide-NMe-nonanoyl derivative 7.** To a 10 mL
48 polypropylene syringe fitted with polyethylene porous disk and loaded with the polymer-
49 bound Fmoc protected pentapeptide **15** (152 mg, L=0.6 mmol/g, 0.091 mmol, 1.0 equiv.), was
50 added a solution of 20% piperidine in DMF (3 x 6 mL x 10 min.) and shaken at 280 r.p.m..
51 After the last run, the resin was washed with dry DMF (5 x 6 mL) and treated with a solution
52 of the hydroxyacid **9** (52 mg, 0.273 mmol, 3.0 equiv.), TEA (39 μL, 0.273 mmol, 3.0 equiv.)
53 and DEPC (45 μL, 0.273 mmol, 3.0 equiv.) in dry DMF (4 mL) for two times, 8 hours each
54
55
56
57
58
59
60

1
2 time. Once the solution was unloaded, the resin was washed with DMF (4 x 4 mL) and DCM
3
4 (4 x 4 mL). The resulting swelled resin **7** was used in the next step.
5
6
7

8
9 **Seco-acid 47.** The resin **7** was treated with a solution of DCM:AcOH:TFE (7:2:1, 3
10 mL) for 30 minutes. After that, the solution was collected and the resin washed with DCM (2
11 x 3 mL). All the collected organic solvents were evaporated under reduced pressure and the
12 resulting seco acid **47**¹⁶ (52 mg, 65%) was obtained as a white solid and not requiring further
13 purification: $[\alpha]_D^{25} = -13.1$ (*c* 0.16, CH₃OH); ¹H NMR (400 MHz, DMSO-*d*₆, two rotamers in
14 a 4:1 ratio) $\delta = -0.03$ (s, 3 H), 0.02 (s, 3 H), 0.69 (d, *J* = 6.8 Hz, 12/5 H), 0.74 (d, *J* = 6.9 Hz,
15 3/5 H), 0.79-0.91 (m, 12 H), 0.81 (s, 9 H), 0.96 (d, *J* = 6.8 Hz, 3 H), 1.05 (d, *J* = 6.2 Hz, 3 H),
16 1.19-1.32 (m, 10 H), 1.33-1.55 (m, 4 H), 1.57-1.67 (m, 1 H), 1.74-1.82 (m, 1 H), 2.69 (s, 3/5
17 H), 2.78-2.89 (m, 1 H), 2.86 (s, 12/5 H), 3.49-3.61 (m, 4 H), 3.78 (dd, *J* = 17.5, 6.1 Hz, 1 H),
18 3.91 (dt, *J* = 15.0, 7.1 Hz, 1/5 H), 4.01 (q, *J* = 6.5 Hz, 4/5 H), 4.36-4.40 (m, 2 H), 4.48 (s, 2
19 H), 4.64-4.73 (m, 6/5 H), 5.11 (dd, *J* = 10.7, 4.8 Hz, 4/5 H), 7.25-7.34 (m, 6 H), 7.91 (d, *J* =
20 8.9 Hz, 1/5 H), 7.98 (d, *J* = 9.3 Hz, 1/5 H), 8.02 (d, *J* = 9.1 Hz, 3/5 H), 8.09 (d, *J* = 7.9 Hz, 1/5
21 H), 8.18 (d, *J* = 8.2 Hz, 4/5 H), 8.21 (t, *J* = 5.8 Hz, 4/5 H); ¹³C NMR (100 MHz, DMSO-*d*₆,
22 both rotamers) $\delta = -5.1, -4.7, 11.6, 13.2, 14.0, 14.1, 17.7, 19.5, 21.5, 22.1, 23.3, 24.1, 25.1,$
23 $25.6, 25.8, 28.8, 30.5, 31.4, 33.4, 36.1, 37.2, 40.6, 41.4, 52.3, 53.2, 55.3, 58.2, 68.7, 69.9,$
24 $71.8, 72.1, 127.4, 127.5, 127.6, 128.2, 138.1, 169.2, 169.7, 170.3, 170.78, 170.83, 176.1.$
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **Cyclodepsipeptide 49.** To a suspension of the seco-acid **47** (50 mg, 0.057 mmol, 1.0
51 equiv.) in dry THF (2 mL) was added dry TEA (25 μ L, 0.171 mmol, 3.0 equiv.) and 2,4,6-
52 trichlorobenzoyl chloride (12 μ L, 0.068 mmol, 1.2 equiv.) at room temperature. After 24 h,
53 the suspension was diluted with dry toluene (27 mL) and added over 5 hours via a syringe
54 pump into a refluxed solution of 4-DMAP (140 mg, 1.14 mmol, 20.0 equiv.) in dry toluene
55 (29 mL). The mixture was refluxed for additional 2 h. Once cooled, the solvents were
56
57
58
59
60

1
2 evaporated under reduced pressure, diluted with EtOAc and washed sequentially with 5%
3
4 HCl, saturated NaHCO₃ and brine aqueous solutions. The organic layer was dried with
5
6 anhydrous MgSO₄, filtered and the solvent evaporated under reduced pressure. The resulted
7
8 brown syrupe was purified by preparative HPLC (column: Phenomenex-luna C8(2), (10 mm x
9
10 250 mm), refraction index detector, flow rate: 4.7 mL/min with 90% MeOH) to give cyclic
11
12 depsipeptide **49**¹⁶ (12 mg, 24%) as a white solid: $[\alpha]_D^{25} = +16.0$ (*c* 0.45, CH₃OH); ¹H NMR
13
14 (400 MHz, CDCl₃, two rotamers in a 7:3 proportion) $\delta = 0.05$ (s, 21/10 H), 0.06 (s, 9/10 H),
15
16 0.07 (s, 21/10 H), 0.08 (s, 9/10H), 0.84-0.99 (m, 15 H), 0.84 (s, 63/10 H), 0.85 (s, 27/10 H),
17
18 1.07-1.16 (m, 6 H), 1.20-1.40 (m, 11 H), 1.41-1.57 (m, 3 H), 1.63-1.75 (m, 27/10 H), 1.88-
19
20 1.94 (m, 1 H), 1.98-2.03 (m, 3/10H), 2.09-2.14 (m, 3/10H), 2.16-2.24 (m, 7/10 H), 2.77 (s,
21
22 9/10 H), 3.01 (dq, *J* = 9.6, 7.0 Hz, 7/10 H), 3.11-3.16 (m, 3/10 H), 3.16 (s, 21/10 H), 3.54 (dd,
23
24 *J* = 17.1, 3.6 Hz, 7/10 H), 3.71 (bs, 7/10 H), 3.77 (dd, *J* = 10.1, 6.3 Hz, 7/10 H), 3.85 (dd, *J* =
25
26 10.1, 4.8 Hz, 7/10 H), 3.89 (dd, *J* = 10.4, 5.5 Hz, 3/10 H), 3.93-4.01 (m, 1 H), 4.20-4.23 (m,
27
28 7/10 H), 4.30 (q, *J* = 4.6 Hz, 3/10 H), 4.38 (dd, *J* = 17.2, 8.8 Hz, 7/10 H), 4.42-4.49 (m, 18/10
29
30 H), 4.51 (d, *J* = 11.8 Hz, 7/10 H), 4.55 (dd, *J* = 7.9, 4.6 Hz, 3/10 H), 4.61 (d, *J* = 11.8 Hz, 7/10
31
32 H), 4.61-4.65 (m, 14/10 H), 4.76 (dd, *J* = 9.4, 3.9 Hz, 3/10 H), 4.83 (d, *J* = 10.7 Hz, 3/10 H),
33
34 5.28 (ddd, *J* = 9.9, 7.5, 2.9 Hz, 7/10 H), 6.39 (d, *J* = 7.5 Hz, 7/10 H), 6.42 (d, *J* = 2.5 Hz, 7/10
35
36 H), 6.69-6.71 (m, 6/10 H), 6.88 (d, *J* = 9.6 Hz, 3/10 H), 7.04 (t, *J* = 5.6 Hz, 3/10 H), 7.28-7.40
37
38 (m, 5 H), 7.71 (dd, *J* = 8.6, 3.6 Hz, 7/10 H), 8.21 (bs, 7/10 H); ¹³C NMR (100 MHz, CDCl₃,
39
40 both rotamers) $\delta = -4.8, -4.7, -4.6, 11.9, 14.1, 14.2, 15.1, 15.2, 15.3, 18.1, 18.4, 19.0, 21.8,$
41
42 22.6, 22.7, 23.2, 23.3, 24.1, 25.0, 25.3, 25.9, 26.5, 27.1, 27.2, 29.1, 29.2, 29.4, 29.5, 31.6,
43
44 31.8, 31.9, 37.5, 37.6, 38.2, 38.5, 39.6, 40.2, 40.6, 56.1, 56.2, 56.3, 56.7, 59.1, 59.3, 59.4,
45
46 66.8, 67.1, 68.03, 68.12, 73.6, 73.9, 76.4, 77.4, 78.4, 128.2, 128.3, 128.5, 128.6, 128.9, 136.8,
47
48 137.0, 168.9, 169.4, 169.5, 169.7, 169.9, 170.1, 171.8.
49
50
51
52
53
54
55
56
57
58
59
60

1
2 **Cyclodepsipeptide 51.** To a solution of cyclic compound **49** (8 mg, 9.3 μ mol, 1.0
3
4 equiv.) in dry THF (0.5 mL) was added at room temperature AcOH (40 μ L, 0.698 mmol, 75.0
5
6 equiv.) and TBAF (0.5 mL, 1 M solution in THF, 0.5 mmol, 54.0 equiv.). After 26 h, the
7
8 mixture was diluted with EtOAc and washed with a saturated aqueous NaHCO₃ solution and
9
10 brine. Once dried over anhydrous MgSO₄ and the solvents evaporated under reduced pressure,
11
12 the crude was purified by preparative HPLC (column: Phenomenex-luna 5 μ m C8(2), (10 mm
13
14 x 250 mm), refraction index detector, flow rate: 4.7 mL/min with 90% MeOH) to give
15
16 cyclodepsipeptide **51**¹⁶ (6 mg, 87%) as a white solid: $[\alpha]_D^{25} = +26.0$ (*c* 0.25, CH₃OH); ¹H
17
18 NMR (400 MHz, CDCl₃, two rotamers in a 3.5/1 proportion) $\delta = 0.85$ -0.98 (m, 15 H), 1.10
19
20 (d, *J* = 6.9 Hz, 12/5 H), 1.16 (d, *J* = 7.0 Hz, 3/5 H), 1.07-1.16 (m, 6 H), 1.20-1.40 (m, 11 H),
21
22 1.19-1.69 (m, 17 H), 1.74 (ddd, *J* = 14.0, 8.6, 5.6 Hz, 1 H), 1.99-2.06 (m, 1 H), 2.08-2.16 (m,
23
24 1 H), 2.78 (s, 3/5 H), 3.07-3.15 (m, 1 H), 3.18 (s, 12/5 H), 3.62 (dd, *J* = 17.2, 3.9 Hz, 4/5 H),
25
26 3.69 (bs, 4/5 H), 3.79 (dd, *J* = 18.5, 3.7 Hz, 1/5 H), 3.84 (dd, *J* = 10.0, 4.8 Hz, 1 H), 3.90 (dd,
27
28 *J* = 10.0, 5.7 Hz, 1 H), 3.95 (dd, *J* = 10.1, 4.2 Hz, 1/5 H), 4.00 (dd, *J* = 11.3, 4.8 Hz, 1/5 H),
29
30 4.11-4.21 (m, 9/5 H), 4.29 (q, *J* = 5.0 Hz, 4/5 H), 4.34-4.47 (m, 2 H), 4.54 (d, *J* = 11.8 Hz, 1
31
32 H), 4.57 (d, *J* = 11.8 Hz, 1 H), 4.77 (dd, *J* = 9.4, 3.9 Hz, 1/5 H), 4.84 (d, *J* = 11.1 Hz, 1/5 H),
33
34 5.08-5.14 (m, 4/5 H), 6.71 (d, *J* = 4.1 Hz, 1/5 H), 6.80 (bs, 1 H), 6.89 (d, *J* = 7.7 Hz, 4/5 H),
35
36 7.19 (d, *J* = 9.2 Hz, 1/5 H), 7.29-7.39 (m, 5 H), 7.59 (dd, *J* = 8.6, 3.6 Hz, 4/5 H), 7.84 (d, *J* =
37
38 6.5 Hz, 4/5 H); ¹³C NMR (100 MHz, CDCl₃, both rotamers) $\delta = 11.69$, 11.75, 11.8, 14.0, 14.7,
39
40 14.8, 15.1, 19.9, 20.0, 21.8, 22.5, 22.6, 23.06, 23.14, 24.2, 24.8, 25.2, 26.3, 26.9, 27.1, 28.8,
41
42 29.0, 29.2, 29.4, 29.7, 31.3, 31.6, 31.7, 36.8, 37.5, 38.1, 38.4, 39.5, 40.5, 40.8, 41.2, 55.4,
43
44 55.6, 56.3, 56.9, 57.2, 58.0, 59.3, 67.7, 67.8, 68.0, 73.5, 73.9, 77.7, 78.2, 127.9, 128.2, 128.5,
45
46 128.65, 128.72, 137.1, 169.1, 169.2, 170.9, 171.0, 171.6, 172.3, 172.8, 173.4, 174.2, 176.7.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Globomycin (1). Cyclodepsipeptide **51** (4.0 mg, 5.4 μ mol) was dissolved in MeOH (1.5 mL) and to the mixture was added Pd(OH)₂ (2.5 mg, 20 wt%). Once the flask was purged

1
2 of air, the reaction was carried out under H₂ atmosphere for 6 h. Then, and once the flask was
3
4 purged of H₂, the mixture was diluted with MeOH (3 mL), and the catalyst removed by
5
6 filtration. The methanolic solution was then evaporated under reduced pressure, and the crude
7
8 product purified by preparative HPLC (column: Phenomenex-luna 5 μm C8(2), (10 mm x 250
9
10 mm), refraction index detector, flow rate: 4.7 mL/min with 90% MeOH) to give globomycin
11
12 (**1**) (3.0 mg, 86%): $[\alpha]_D^{25} = +21.9$ (*c* 0.13, CH₃OH) (Reported: *lit.*¹⁵ $[\alpha]_D^{25} = +23.8$ (*c* 0.5,
13
14 CH₃OH)); ¹H NMR (400 MHz, CDCl₃, 3.5 mM, two rotamers in a 5.9/1 proportion) $\delta = 0.83$ -
15
16 1.00 (m, 15 H), 1.09 (d, *J* = 6.9 Hz, 18/7 H), 1.17 (d, *J* = 6.9 Hz, 3/7 H), 1.19-1.44 (m, 13 H),
17
18 1.49-1.82 (m, 4 H), 1.98-2.06 (m, 2/7 H), 2.06-2.14 (m, 6/7 H), 2.19-2.28 (m, 6/7 H), 2.79 (s,
19
20 3/7 H), 2.95-3.01 (m, 1 H), 3.22 (s, 18/7 H), 3.60 (dd, *J* = 7.7, 5.6 Hz, 6/7 H), 3.81 (dd, *J* =
21
22 17.3, 4.6 Hz, 1 H), 3.91 (t, *J* = 7.2 Hz, 6/7 H), 4.00 (d, *J* = 4.4 Hz, 6/7 H), 4.05-4.07 (m, 3/7
23
24 H), 4.08-4.16 (m, 6/7 H), 4.28 (dd, *J* = 17.4, 7.5 Hz, 1 H), 4.36 (q, *J* = 4.9 Hz, 1 H), 4.41-4.46
25
26 (m, 9/7 H), 4.48-4.58 (m, 5/7 H), 4.78 (dd, *J* = 9.9, 4.0 Hz, 1/7 H), 4.85 (d, *J* = 10.6 Hz, 1/7
27
28 H), 4.92 (dt, *J* = 9.2, 3.0 Hz, 6/7 H), 6.70 (bs, 1/7 H), 7.00 (bs, 6/7 H), 7.08 (bs, 1/7 H), 7.16
29
30 (d, *J* = 8.9 Hz, 1/7 H), 7.24 (bs, 3/7 H), 7.40-7.49 (m, 1 H), 7.55-7.61 (m, 1 H), 7.76-7.79 (m,
31
32 2 H); ¹³C NMR (100 MHz, CDCl₃, both rotamers) $\delta = 11.7, 13.5, 14.1, 14.6, 15.0, 19.3, 20.2,$
33
34 22.0, 22.6, 22.7, 23.1, 24.6, 25.1, 25.4, 27.1, 29.0, 29.7, 31.2, 31.7, 31.9, 35.9, 38.0, 40.1,
35
36 40.6, 41.0, 56.2, 56.9, 59.0, 61.6, 66.8, 169.1, 170.6, 171.1, 172.6, 174.1, 176.4.
37
38
39
40
41
42
43
44
45
46
47

48 **Seco Acid 48.** Compound **48** was obtained following the same procedure described for
49
50 the synthesis of compound **47**, using the polymer-bound Fmoc protected pentapeptide **15** (118
51
52 mg, L=0.6 mmol/g, 0.071 mmol, 1.0 equiv.), hydroxyacid **10** (44 mg, 0.213 mmol, 3.0
53
54 equiv.), TEA (30 μL, 0.213 mmol, 3.0 equiv.) and DEPC (35 μL, 0.213 mmol, 3.0 equiv.) for
55
56 two times, 18 hours each time. After cleavage, compound **48**¹⁶ was obtained (49 mg, 77%): ¹H
57
58 NMR (400 MHz, DMSO-*d*₆, two rotamers in a 3.4:1 proportion) $\delta = -0.03$ (s, 3 H), 0.02 (s, 3
59
60 H), 0.70 (d, *J* = 6.9 Hz, 12/5 H), 0.74 (d, *J* = 6.9 Hz, 3/5 H), 0.79-0.91 (m, 12 H), 0.81 (s, 9

1
2 H), 0.96 (d, $J = 6.8$ Hz, 3 H), 1.06 (d, $J = 6.2$ Hz, 3 H), 1.19-1.31 (m, 14 H), 1.32-1.54 (m, 4
3
4 H), 1.58-1.67 (m, 1 H), 1.74-1.82 (m, 1 H), 2.69 (s, 3/5 H), 2.78-2.84 (m, 1 H), 2.87 (s, 12/5
5
6 H), 3.48-3.55 (m, 8/5 H), 3.57-3.61 (m, 12/5 H), 3.78 (dd, $J = 17.5, 6.2$ Hz, 1 H), 4.02 (q, $J =$
7
8 6.6 Hz, 1 H), 4.36-4.41 (m, 2 H), 4.49 (s, 2 H), 4.64-4.72 (m, 2 H), 5.11 (dd, $J = 10.6, 5.2$ Hz,
9
10 1 H), 7.24-7.34 (m, 6 H), 7.90 (d, $J = 9.5$ Hz, 1/5 H), 7.94 (d, $J = 8.8$ Hz, 1/5 H), 7.98 (d, $J =$
11
12 9.1 Hz, 3/5 H), 8.06 (d, $J = 8.0$ Hz, 1/5 H), 8.18-8.21 (m, 9/5 H); ^{13}C NMR (100 MHz,
13
14 DMSO- d_6 , both rotamers) $\delta = -5.2, -4.8, 11.4, 13.1, 13.9, 14.1, 17.6, 19.5, 21.5, 22.1, 23.2,$
15
16 24.1, 25.6, 28.6, 28.9, 29.1, 30.4, 31.2, 33.4, 36.0, 37.1, 40.6, 41.3, 52.2, 53.2, 55.3, 58.3,
17
18 68.6, 69.8, 71.8, 72.1, 127.4, 127.5, 128.2, 138.0, 169.2, 169.6, 170.2, 170.6, 170.8, 176.1.
19
20
21
22
23
24
25

26 **Cyclodepsipeptide 50.** Yamaguchi macrolactonization of compound **48** was carried
27
28 out following the same procedure described for globomycin, using seco-acid **48** (46 mg, 50.8
29
30 μmol , 1.0 equiv.), TEA (22 μL , 152.0 μmol , 3.0 equiv.) and 2,4,6-trichlorobenzoyl chloride
31
32 (12 μL , 66.0 μmol , 1.3 equiv.). After purification by preparative HPLC (column:
33
34 Phenomenex-luna C8(2), (10 mmx250 mm), refraction index detector, flow rate: 4.7 mL/min
35
36 with 95% MeOH) the expected cyclic compound **50**¹⁶ (13.4 mg, 30%) was obtained as a white
37
38 solid: $[\alpha]_D^{25} = +10.0$ (c 0.35, CHCl_3); ^1H NMR (400 MHz, CDCl_3 , two rotamers in a 7:3
39
40 proportion) $\delta = 0.056$ (s, 21/10 H), 0.061 (s, 9/10 H), 0.067 (s, 21/10 H), 0.08 (s, 9/10H),
41
42 0.84-0.99 (m, 15 H), 0.84 (s, 63/10 H), 0.85 (s, 27/10 H), 1.06-1.19 (m, 6 H), 1.21-1.39 (m, 15
43
44 H), 1.41-1.56 (m, 3 H), 1.60-1.76 (m, 27/10 H), 1.88-1.95 (m, 1 H), 1.98-2.05 (m, 3/10H),
45
46 2.09-2.14 (m, 3/10H), 2.16-2.25 (m, 7/10 H), 2.77 (s, 9/10 H), 3.01 (dq, $J = 9.6, 6.9$ Hz, 7/10
47
48 H), 3.10-3.16 (m, 3/10 H), 3.16 (s, 21/10 H), 3.54 (dd, $J = 17.1, 3.7$ Hz, 7/10 H), 3.71 (bs,
49
50 7/10 H), 3.77 (dd, $J = 10.2, 6.3$ Hz, 7/10 H), 3.85 (dd, $J = 10.1, 4.8$ Hz, 7/10 H), 3.89 (dd, $J =$
51
52 10.2, 5.5 Hz, 3/10 H), 3.93-4.01 (m, 1 H), 4.20-4.23 (m, 7/10 H), 4.30 (q, $J = 4.4$ Hz, 3/10 H),
53
54 4.39 (dd, $J = 17.3, 8.8$ Hz, 7/10 H), 4.42-4.49 (m, 18/10 H), 4.51 (d, $J = 11.8$ Hz, 7/10 H),
55
56 4.54-4.57 (m, 3/10 H), 4.61 (d, $J = 11.8$ Hz, 7/10 H), 4.61-4.65 (m, 14/10 H), 4.77(dd, $J = 9.4,$
57
58
59
60

1
2 3.9 Hz, 3/10 H), 4.83 (d, $J = 10.9$ Hz, 3/10 H), 5.28 (ddd, $J = 9.9, 7.4, 2.9$ Hz, 7/10 H), 6.39
3
4 (d, $J = 7.4$ Hz, 7/10 H), 6.41 (d, $J = 2.6$ Hz, 7/10 H), 6.68-6.71 (m, 6/10 H), 6.88 (d, $J = 9.2$
5
6 Hz, 3/10 H), 7.04 (t, $J = 5.5$ Hz, 3/10 H), 7.28-7.40 (m, 5 H), 7.70 (dd, $J = 8.8, 3.7$ Hz, 7/10
7
8 H), 8.21 (bs, 7/10 H); ^{13}C NMR (100 MHz, CDCl_3 , both rotamers) $\delta = -4.9, -4.8, -4.7, 11.8,$
9
10 14.1, 14.9, 15.1, 18.0, 18.2, 18.9, 21.6, 22.5, 22.6, 23.1, 23.3, 24.1, 24.9, 25.2, 25.7, 26.5,
11
12 27.0, 27.1, 29.0, 29.2, 29.3, 29.4, 29.5, 29.6, 29.7, 31.5, 31.8, 31.9, 37.4, 37.5, 38.2, 38.5,
13
14 39.6, 40.1, 40.5, 56.1, 56.2, 56.3, 56.6, 59.1, 59.3, 59.4, 66.6, 66.9, 67.9, 68.0, 73.4, 73.9,
15
16 76.3, 77.2, 77.7, 78.2, 128.1, 128.4, 128.5, 128.7, 136.7 136.9, 168.8, 169.2, 169.3, 169.6,
17
18 169.7, 169.9, 171.7, 172.9, 173.1, 174.2, 174.5, 177.2.
19
20
21
22
23
24
25

26 **Cyclodepsipeptide 52.** The desilylation of **50** (6.0 mg, 6.8 mmol) was carried out
27
28 using the same procedure described for globomycin, by treatment with AcOH (40 μL) and
29
30 TBAF (0.5 mL, 1 M in THF). After purification by preparative HPLC (column: Phenomenex-
31
32 luna C8(2), (10 mmx250 mm), refraction index detector, flow rate: 4.7 mL/min with 95%
33
34 MeOH) cyclodepsipeptide **52**¹⁶ (4.5 mg, 87%) was obtained as a white solid: $[\alpha]_{\text{D}}^{25} = +10.0$ (c
35
36 0.2, CHCl_3); ^1H NMR (400 MHz, CDCl_3 , two rotamers in a 3.5/1 proportion) $\delta = 0.85$ -1.03
37
38 (m, 15 H), 1.10 (d, $J = 6.9$ Hz, 12/5 H), 1.16 (d, $J = 7.0$ Hz, 3/5 H), 1.26-1.54 (m, 20 H), 1.55-
39
40 1.67 (m, 1 H), 1.74 (ddd, $J = 14.1, 8.6, 5.7$ Hz, 1 H), 1.99-2.07 (m, 1 H), 2.08-2.17 (m, 1 H),
41
42 2.78 (s, 3/5 H), 3.07-3.15 (m, 1 H), 3.18 (s, 12/5 H), 3.62 (dd, $J = 17.2, 3.9$ Hz, 4/5 H), 3.68
43
44 (bs, 4/5 H), 3.80-3.94 (m, 3/5 H), 3.84 (dd, $J = 10.0, 5.7$ Hz, 4/5 H), 3.90 (dd, $J = 10.0, 4.8$
45
46 Hz, 4/5 H), 4.00-4.08 (m, 3/5 H), 4.13 (m, 4/5 H), 4.20 (dd, $J = 8.2, 6.3$ Hz, 4/5 H), 4.30 (q, J
47
48 = 5.0 Hz, 4/5 H), 4.34-4.42 (m, 8/5 H), 4.46 (t, $J = 7.1$ Hz, 1/5 H), 4.52 (d, $J = 11.8$ Hz, 1/5
49
50 H), 4.53 (d, $J = 11.8$ Hz, 4/5 H), 4.57 (d, $J = 11.8$ Hz, 4/5 H), 4.60 (d, $J = 11.8$ Hz, 1/5 H),
51
52 4.77 (dd, $J = 9.4, 3.6$ Hz, 1/5 H), 4.82 (d, $J = 10.9$ Hz, 1/5 H), 5.11 (ddd, $J = 9.3, 7.3, 4.3$ Hz,
53
54 4/5 H), 6.78-6.84 (d, 7/5 H), 6.90 (d, $J = 8.2$ Hz, 4/5 H), 7.22 (bs, 1/5 H), 7.28-7.39 (m, 5 H),
55
56 7.59 (dd, $J = 8.2, 3.3$ Hz, 4/5 H), 7.84 (d, $J = 6.6$ Hz, 4/5 H); ^{13}C NMR (100 MHz, CDCl_3 ,
57
58
59
60

1
2 both rotamers) $\delta = 11.7, 11.8, 13.7, 14.1, 14.7, 14.8, 15.1, 19.8, 21.9, 22.5, 22.7, 23.0, 23.2,$
3
4 $24.2, 24.3, 24.7, 25.2, 26.3, 26.9, 27.0, 28.7, 29.2, 29.4, 29.6, 29.7, 31.3, 31.9, 36.8, 37.5,$
5
6 $38.1, 38.5, 39.5, 40.4, 40.7, 41.3, 55.4, 55.7, 56.2, 56.9, 57.5, 58.0, 59.1, 59.3, 67.6, 67.8,$
7
8 $68.0, 73.4, 73.9, 77.2, 77.7, 78.1, 127.9, 128.2, 128.3, 128.4, 128.6, 136.7, 137.1, 169.1,$
9
10 $169.2, 169.8, 170.1, 170.8, 171.0, 171.7, 172.3, 172.8, 173.8, 174.2, 176.7.$
11
12
13
14
15

16 **SF 1902 A₅ (2).** Natural compound SF 1902 A₅ (2) was obtained following the same
17
18 procedure described for globomycin, by treatment of **52** (3.0 mg, 3.9 mmol) with Pd(OH)₂
19
20 (3mg, 20 wt%) under H₂ atmosphere. After purification by preparative HPLC (column:
21
22 Phenomenex-luna 5 μ m C8(2), (10 mm x 250 mm), refraction index detector, flow rate: 4.7
23
24 mL/min with 90% MeOH) SF 1902 A₅ (2)¹⁶ (2.2 mg, 81%) was obtained as a white solid:
25
26 $[\alpha]_D^{25} = +25.8$ (*c* 0.12, CH₃OH) ($[\alpha]_D^{25} = +20.8$ (*c* 1.04, CH₃OH); ¹H NMR (400 MHz, CDCl₃,
27
28 3.5 mM, two rotamers in a 5.7/1 proportion) $\delta = 0.83-1.05$ (m, 17 H), 1.09 (d, *J* = 6.9 Hz,
29
30 18/7 H), 1.16 (d, *J* = 6.9 Hz, 3/7 H), 1.19-1.42 (m, 15 H), 1.48-1.79 (m, 6 H), 2.05-2.15 (m, 1
31
32 H), 2.20-2.27 (m, 1 H), 2.79 (s, 3/7 H), 3.16-3.27 (m, 1 H), 3.22 (s, 18/7 H), 3.61 (dd, *J* = 8.7,
33
34 6.1 Hz, 6/7 H), 3.80 (dd, *J* = 17.3, 4.7 Hz, 6/7 H), 3.92 (t, *J* = 7.1 Hz, 6/7 H), 4.00 (d, *J* = 4.6
35
36 Hz, 12/7 H), 4.04-4.06 (m, 1/7 H), 4.09-4.11 (m, 1/7 H), 4.14-4.16 (m, 1/7 H), 4.28 (dd, *J* =
37
38 17.3, 7.9 Hz, 6/7 H), 4.35 (q, *J* = 5.1 Hz, 6/7 H), 4.39-4.46 (m, 16/7 H), 4.52 (t, *J* = 7.0 Hz,
39
40 1/7 H), 4.79 (dd, *J* = 9.4, 3.8 Hz, 1/7 H), 4.85 (d, *J* = 9.9 Hz, 1/7 H), 4.94 (td, *J* = 8.7, 2.9 Hz,
41
42 6/7 H), 6.81 (d, *J* = 8.9 Hz, 1/7 H), 6.99 (d, *J* = 9.5 Hz, 1/7 H), 7.05 (bs, 8/7 H), 7.15 (bs, 1/7
43
44 H), 7.43 (d, *J* = 8.2 Hz, 1/7 H), 7.49 (t, *J* = 5.9 Hz, 8/7 H), 7.56 (d, *J* = 6.1 Hz, 8/7 H); ¹³C
45
46 NMR (100 MHz, CDCl₃, 3.5 mM, both rotamers) $\delta = 11.7, 14.1, 14.7, 14.9, 19.6, 22.0, 22.7,$
47
48 $23.2, 24.8, 25.3, 27.1, 29.3, 29.4, 31.2, 31.9, 35.7, 37.9, 40.2, 41.1, 55.7, 57.0, 59.1, 61.7, 66.8,$
49
50 $77.7, 169.1, 170.4, 171.3, 172.3, 173.7, 176.1.$
51
52
53
54
55
56
57
58
59
60

Acknowledgments

This work was financially supported by Ministerio de Educación y Ciencia (ref. CTQ2007-66518) and Junta de Andalucía (FQM-03329). We are deeply indebted to Prof. Masatoshi Inukai from International University of Health and Welfare in Ohtawara (Japan) for a generous gift of a natural sample of globomycin. We also thank Dr. J. I. Trujillo from Pfizer (Groton, CT) for assistance in the preparation of this manuscript. Finally, we thank Unidad de Espectroscopía de Masas de la Universidad de Granada for exact mass spectroscopic assistance.

Supporting Information Available

Experimental procedures and spectroscopic data of all new compounds, as well as ^1H - and ^{13}C -NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes


1. (a) Sarabia, F.; Chammaa, S. *J. Org. Chem.* **2005**, *70*, 7846-7857. (b) Sarabia, F.; García-Castro, M.; Chammaa, S. *Tetrahedron Lett.* **2005**, *46*, 7695-7699. (c) Bisek, N.; Wetzel, S.; Arndt, H.-D.; Waldmann, H. *Chem. Eur. J.* **2008**, *14*, 8847-8860 and references therein.
2. Sarabia, F.; Chammaa, S.; Sánchez-Ruiz, A.; Martín-Ortiz, L.; López-Herrera, F. J. *Curr. Med. Chem.* **2004**, *11*, 1309-1332.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
3. (a) Inukai, M.; Enokita, R.; Torikata, A.; Ankara, M. Iwado, S.; Arai, M. *J. Antibiot.* **1978**, *31*, 410-420. (b) Inukai, M.; Nakajima, M.; Osawa, M.; Haneishi, T.; Arai, M. *J. Antibiot.* **1978**, *31*, 421-425. (c) Nakajima, M.; Inukai, M.; Haneishi, T.; Terahara, A.; Enokita, R.; Arai, M.; Kinoshita, T.; Tamura, C. *J. Antibiot.* **1978**, *31*, 426-432.
4. (a) Omoto, S.; Suzuki, H.; Inouye, S. *J. Antibiot.* **1979**, *32*, 83-86. (b) Omoto, S.; Ogino, H.; Inouye, S. *J. Antibiot.* **1981**, *34*, 1416-1423.
5. (a) Fehri, L. F.; Wróblewski, H.; Blanchard, A. *Antimicrob. Agents Chemother.* **2007**, *51*, 468-474. (b) Malléa, M.; Chevalier, J.; Eyraud, A.; Pagès, J.-M. *Biochem. Biophys. Res. Commun.* **2002**, *293*, 1370-1373. (c) Inukai, M.; Takeuchi, M.; Shimizu, K. *Agric. Biol. Chem.* **1984**, *48*, 513-518.
6. (a) Inukai, M.; Takeuchi, M.; Shimizu, K.; Arai, M. *J. Antibiot.* **1978**, *31*, 1203-1205. (b) Hussain, M.; Ichihara, S.; Mizushima, S. *J. Biol. Chem.* **1980**, *255*, 3707-3712. (c) Dev, I. K.; Harvey, R. J.; Ray, P. H. *J. Biol. Chem.* **1985**, *260*, 5891-5894. (d) Geukens, N.; De Buck, E.; Meyen, E.; Maes, L; Vranckx, L.; Van Mellaert, L.; Anné, J.; Lammertyn, E. *Res. Microbiol.* **2006**, *157*, 836-841.
7. (a) Sankaran, K. in *Handbook of Proteolytic Enzymes*, Ed. Barrett, A. J.; Rawlings, N. D.; Woessner, J. F., 1, 201-204, 2004. (b) Paetzel, M.; Karla, A.; Strynadka, N. C. J.; Dalbey, R. E. *Chem. Rev.* **2002**, *102*, 4549-4579. (c) Pugsley, A. P. *Microbiol. Rev.* **1993**, *57*, 50-108.
8. (a) Walsh, C. in *Antibiotics: Actions, Origins, Resistance*, ASM Press (Washington), 2003. (b) Chu, D. T. W.; Plattner, J. J.; Katz, L. *J. Med. Chem.* **1996**, *39*, 3853-3874. (c) Süssmuth, R.D. *ChemBioChem* **2002**, *3*, 295-298. (d) Gao, Y. *Nat. Prod. Rep.* **2002**, *19*, 100-107. (e) Nicolaou, K. C.; Boddy, C.N.C.; Bräse, S.; Winssinger, N. *Angew. Chem. Int. Ed. Engl.* **1999**, *38*, 2096-2152.
9. (a) Huang, H.; Lin, M.; Wang, X.; Kikuchi, T.; Mottaz, H.; Norbeck, A.; Rikihisa, Y. *Infect. Immun.* **2008**, *76*, 3405-3414. (b) Gibbons, H. S.; Wolschendorf, F.; Abshire,

- 1
2 M.; Niederweis, M.; Braunstein, M. *J. Bacteriol.* **2007**, *189*, 5090-5100. (c) Lo, M.;
3
4 Boyce, J. D.; Wilkie, I. W.; Adler, B. *Microb. Infec.* **2004**, *6*, 58-67. (d) Jackowski, S.;
5
6 Rock, C. O. *J. Biol. Chem.* **1986**, *261*, 11328-11333. (e) Tokunaga, M.; Tokunaga, H.;
7
8 Wu, H. C. *Proc. Natl. Acad. Sci. USA* **1982**, *79*, 2255-2259.
9
10
11 10. Some relevant publications: (a) Ghrayeb, J.; Lunn, C. A.; Inouye, S.; Inouye, M. *J.*
12
13 *Biol. Chem.* **1985**, *260*, 10961-10965. (b) Inukai, M.; Ghrayeb, J.; Nakamura, K.;
14
15 Inouye, M. *J. Biol. Chem.* **1984**, *259*, 757-760. (c) Hussain, M.; Ichihara, S.;
16
17 Mizushima, S. *J. Biol. Chem.* **1982**, *257*, 5177-5182. (d) Ichihara, S.; Hussain, M.;
18
19 Mizushima, S. *J. Biol. Chem.* **1982**, *257*, 495-500. (e) Ichihara, S.; Hussain, M.;
20
21 Mizushima, S. *J. Biol. Chem.* **1981**, *256*, 3125-3129. (f) Ichihara, S.; Hussain, M.;
22
23 Mizushima, S. *J. Biol. Chem.* **1981**, *256*, 495-500. (g) Hussain, M.; Ichihara, S.;
24
25 Mizushima, S. *J. Biol. Chem.* **1980**, *255*, 3707-3712.
26
27
28
29
30 11. Recent publications in this area: (a) Banaei, N.; Kincaid, E. Z.; Lin, S.-Y. G.;
31
32 Desmond, E.; Jacobs Jr., W. R.; Ernst, J. D. *Antimicrob. Agents Chemother.* **2009**, *53*,
33
34 3799-3802. (b) Guglielmetti, S.; Tamagnini, I.; Mora, D.; Minuzzo, M.; Scarafoni, A.;
35
36 Arioli, S.; Hellman, J.; Karp, M.; Parini, C. *Appl. Environ. Microbiol.* **2008**, *74*, 4695-
37
38 4702. (c) Ashgar, S. S. A.; Oldfield, N. J.; Wooldridge, K. G.; Jones, M. A.; Irving, G.
39
40 J.; Turner, D. P. J.; Ala'Aldeen, D. A. A. *J. Bacteriol.* **2007**, *189*, 1856-1865. (d)
41
42 Valente, F. M. A.; Pereira, P. M.; Venceslau, S. S.; Regalla, M.; Coelho, A. V.;
43
44 Pereira, I. A. C. *FEBS Lett.* **2007**, *581*, 3341-3344. (e) Neff, L.; Daher, S.; Muzzin, P.;
45
46 Spenato, U.; Gülacar, F.; Gabay, C.; Bas, S. *J. Bacteriol.* **2007**, *189*, 4739-4748. (e)
47
48 Gonnet, P.; Rudd, K. E.; Lisacek, F. *Proteomics* **2004**, *4*, 1597-1613. (f) Venema, R.;
49
50 Tjalsma, H.; van Dijl, J. M.; de Jong, A.; Leenhouts, K.; Buist, G.; Venema, G. *J. Biol.*
51
52 *Chem.* **2003**, *278*, 14739-14746. (g) Harrington, D. J.; Greated, J. S.; Chanter, N.;
53
54 Sutcliffe, I. C. *Infect. Immun.* **2000**, *68*, 6048-6051.
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
12. Kiho, T.; Nakayama, M.; Yasuda, K.; Miyakoshi, S.; Inukai, M.; Kogen, H. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2315-2318.
 13. Banaiee, N.; Jacobs, W. R.; Ernst, J. D. *J. Antimicrob. Chemother.* **2007**, *60*, 414-416.
 14. Haneishi, T.; Nakajima, M.; Inukai, M.; Terahara, A.; Arai, M.; Kinoshita, T.; Tamura, C. *Peptide Chem.* **1980**, *17*, 25-30.
 15. Kogen, H.; Kiho, T.; Nakayama, M.; Furukawa, Y.; Kinoshita, T.; Inukai, M. *J. Am. Chem. Soc.* **2000**, *122*, 10214-10215.
 16. Kiho, T.; Nakayama, M.; Kogen, H. *Tetrahedron* **2003**, *59*, 1685-1697.
 17. Kiho, T.; Nakayama, M.; Yasuda, K.; Miyakoshi, S.; Inukai, M.; Kogen, H. *Bioorg. Med. Chem.* **2004**, *12*, 337-361.
 18. Barlos, D.; Gatos, D.; Schäfer, W. *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 590-593.
 19. For recent examples of synthesis of peptidic chains in solid phase by a Fmoc strategy see: (a) Izzo, I.; Acosta, G. A.; Tulla-Puche, J.; Cupido, T.; Martín-López, M. J.; Cuevas, C.; Albericio, F. *Eur. J. Org. Chem.* **2010**, 2536-2543. (b) Tannert, R.; Milroy, L.-G.; Ellinger, B.; Hu, T.-S.; Arndt, H.-D.; Waldmann, H. *J. Am. Chem. Soc.* **2010**, *132*, 3063-3077.
 20. The synthesis of this amino acid derivative is described in Supporting Information.
 21. (a) Sarabia, F.; Chammaa, S.; García-Castro, M.; Martín-Gálvez, F. *Chem. Commun.* **2009**, 5763-5765. (b) Sarabia, F.; Martín-Gálvez, F.; Chammaa, S.; Martín-Ortiz, L.; Sánchez-Ruiz, A. *J. Org. Chem.* **2010**, *75*, 5526-5532.
 22. Yudin, A. K. (Ed.) in *Aziridines and Epoxides in Organic Synthesis*; Wiley-VCH (Weinheim), 2006.
 23. (a) Sarabia, F.; Martín-Ortiz, L.; López-Herrera, F. J. *Org. Lett.* **2003**, *5*, 3927-3930. (b) Sarabia, F.; Sánchez-Ruiz, A.; Chammaa, S. *Bioorg. Med. Chem.* **2005**, *13*, 1691-1705. (c) Sarabia, F.; Martín-Ortiz, L. *Tetrahedron* **2005**, *61*, 11850-11865.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
24. Kitching, W.; Lewis, J. A.; Perkins, M. V.; Drew, R.; Moore, C. J.; Schurig, V.; König, W. A.; Francke, W. *J. Org. Chem.* **1989**, *54*, 3893-3902.
25. (a) Sarabia, F.; Sánchez-Ruiz, A.; Martín-Ortiz, L.; García-Castro, M.; Chammaa, S. *Org. Lett.* **2007**, *9*, 5091-5094. (b) Sarabia, F.; Martín-Gálvez, F.; García-Castro, M.; Chammaa, S.; Sánchez-Ruiz, A.; Tejón-Blanco, J. F. *J. Org. Chem.* **2008**, *73*, 8979-8986.
26. Brown, H. C.; Kim, S. C.; Krishnamurthy, S. *J. Org. Chem.* **1980**, *45*, 1-12.
27. (a) Spletstoser, J. T.; White, J. M.; Tunoori, A. R.; Georg, G. I. *J. Am. Chem. Soc.* **2007**, *129*, 3408-3419. (b) Tessier, A.; Pytkowicz, J.; Brigaud, T. *Angew. Chem. Int. Ed.* **2006**, *45*, 3677-3681.
28. (a) Myers, A. G.; Yang, B. H.; Chen, H.; Mckinstry, L.; Kopecky, D. J.; Gleason, J. L. *J. Am. Chem. Soc.* **1997**, *119*, 6496-6511. (b) Smith III, A. B.; Bosanac, T.; Basu, K. *J. Am. Chem. Soc.* **2009**, *131*, 2348-2358.
29. See Supporting Information for the description of hydroxyl silylethers **SI-2** and **SI-3**.



- 35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
30. Epp, J. B.; Widlanski, T. S. *J. Org. Chem.* **1999**, *64*, 293-295.
31. (a) Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* **1980**, *102*, 5974-5976. (b) Johnson, R. A.; Sharpless, K. B. in *Catalytic Asymmetric Synthesis* (Ed.: I. Ojima), VCH: Weinheim, New York, **1993**, pp. 103-158.
32. Preparation of epoxy alcohol **30** and its enantiomer by Sharpless asymmetric epoxidation were reported by Kitching *et al* indicated in reference 24, however yields were not reported. Later, the enantiomer of **30** was described in 95% yield by Ortega, N.; Martín, T.; Martín, V. S. *Eur. J. Org. Chem.* **2009**, 554-563.
33. Chong, J. M.; Cyr, D. R.; Mar, E. K. *Tetrahedron Lett.* **1987**, *28*, 5009-5012.

- 1
2 34. As for **30**, the synthesis of the enantiomer of epoxy alcohol **31** via Sharpless
3
4 asymmetric epoxidation is reported by Tanaka, T.; Hiramatsu, K.; Kobayashi, Y.;
5
6 Ohno, H. *Tetrahedron* **2005**, *61*, 6726-6742, in 52% yield.
7
8
9 35. Parikh, J. R.; Doering, W. v. E. *J. Am. Chem. Soc.* **1967**, *89*, 5505-5507.
10
11 36. Chow, K. Y.-K.; Bode, J. W. *J. Am. Chem. Soc.* **2004**, *126*, 8126-8127.
12
13 37. Morrison, K. C.; Litz, J. P.; Scherpelz, K. P.; Dossa, P. D.; Vosburg, D. A. *Org. Lett.*
14
15 **2009**, *11*, 2217-2218.
16
17 38. Tiong, E. A.; Gleason, J. L. *Org. Lett.* **2009**, *11*, 1725-1728.
18
19 39. Recent examples of amides hydrolysis under acidic conditions: (a) Arpin, A.;
20
21 Manthorpe, J. M.; Gleason, J. L. *Org. Lett.* **2006**, *8*, 1359-1362. (b) Tannert, R.;
22
23 Milroy, L.-G.; Ellinger, B.; Hu, T.-S.; Arndt, H.-D.; Waldmann, H. *J. Am. Chem. Soc.*
24
25 **2010**, *132*, 3063-3077.
26
27 40. Evans, D. A.; Bartroli, J.; Shih, T. L. *J. Am. Chem. Soc.* **1981**, *103*, 2127-2129.
28
29 41. Sergeeva, N. N.; Golubev, A. S.; Hennig, L.; Burger, K. *Synthesis* **2002**, 2579-2584.
30
31 42. Deutsch, H. M.; Glinski, J. A.; Hernandez, M.; Haugwitz, R. D.; Narayanan, V. L.;
32
33 Suffness, M.; Zalkow, L. H. *J. Med. Chem.* **1989**, *32*, 788-792.
34
35 43. For recent examples of *N*-Boc activation of amides see: (a) Liu, W.-J.; Lv, B.-D.;
36
37 Gong, L.-Z. *Angew. Chem. Int. Ed.* **2009**, *48*, 6503-6506. (b) Suarez del Villar, I.;
38
39 Gradillas, A.; Domínguez, G.; Pérez-Castells, J. *Tetrahedron Lett.* **2010**, *51*, 3095-
40
41 3098.
42
43 44. For a recent example of *N*-Sulfonyl activation of amides see: Liau, B. B.; Shair, M. D.
44
45 *J. Am. Chem. Soc.* **2010**, *132*, 9594-9595.
46
47 45. Sarabia, F.; Martín-Ortiz, L.; López-Herrera, F. J. *Org. Biomol. Chem.* **2003**, *1*, 3716-
48
49 3725.
50
51 46. Shioiri, T.; Yokoyama, Y.; Kasai, Y.; Yamada, S. *Tetrahedron* **1976**, *32*, 2211-2217
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
47. Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, 52, 1989-1993.
48. See the Supporting Information for General Techniques.