


Journal Pre-proof

Prevalence and correlates of suicidality in Andalusia (Spain): Results of the epidemiological study PISMA-ep

Paloma Huertas Maestre , Berta Moreno-Küstner ,
Blanca Gutiérrez , Jorge A. Cervilla

PII: S0165-0327(19)32450-4
DOI: <https://doi.org/10.1016/j.jad.2020.01.115>
Reference: JAD 11564



To appear in: *Journal of Affective Disorders*

Received date: 23 September 2019
Revised date: 6 December 2019
Accepted date: 20 January 2020

Please cite this article as: Paloma Huertas Maestre , Berta Moreno-Küstner , Blanca Gutiérrez , Jorge A. Cervilla , Prevalence and correlates of suicidality in Andalusia (Spain): Results of the epidemiological study PISMA-ep, *Journal of Affective Disorders* (2020), doi: <https://doi.org/10.1016/j.jad.2020.01.115>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V.

Highlights

- This is the first epidemiological study in Andalusia on suicidality which the main strength is the high sample response of a relatively large sample.
- The non-lethal suicide behaviour is affecting over 1 in every 25 persons in Andalusia.
- This study offers important results of clinical interest for suicide prevention.

Journal Pre-proof

Prevalence and correlates of suicidality in Andalusia (Spain): Results of the epidemiological study PISMA-ep

Paloma Huertas Maestre¹, Berta Moreno-Küstner², Blanca Gutiérrez³, Jorge A. Cervilla^{3,4}

¹Departament of Personality, Assessment and Psychological Treatment, University of Málaga, Málaga, Spain. Grupo GAP.

²Departament of Personality, Assessment and Psychological Treatment, University of Málaga, Málaga, Spain. IBIMA. Group GAP

³Department of Psychiatry, University of Granada, Granada, Spain

⁴San Cecilio University Hospital, Granada, Spain

Corresponding author:

Berta Moreno- Küstner

Departament of Personality, Assessment and Psychological Treatment

Faculty of Psychology, Campus Teatinos, s/n 29071 Málaga (Spain)

Email: bertamk@uma.es

Declaration of interest: None

Abstract:

Background: Suicidality is an important public health problem. Hence, the aims of this study are to report prevalence rates and correlates of suicidality in Andalusia (Southern Spain).

Methods: This is a cross-sectional household survey conducted on a representative sample of adults living in Andalusia. 4507 subjects were interviewed using the Mini International Neuropsychiatric Interview (MINI) to assess suicidality and standardized instruments were employed to evaluate associated variables. A multivariate logistic regression analysis was used to explore independent associations with suicidality.

Results: Current prevalence of suicidality was 6.4 % (95% CI: 5.7-7.12). 4.4% showed death wish, 1.4% had ideas of self-harm, 2.4% had suicidal thoughts, 1.1% had a suicidal plan, 0.6% had attempted suicide during the month prior to the interview, and, lastly, 2.6% reported to have had any sort of suicide attempt during his/her previous life.

Independent factors associated with suicidality were being female, older age, not having a stable couple, lower levels of social support, having had physical childhood abuse experience, having experienced an increasing number of stressful life events, higher neuroticism scores, having a family history of mental disorder and nicotine or drugs dependence.

Limitations: The instrument employed to measure suicidality is a screening tool rather than a more in-depth diagnostic measure. We have not included all potential correlates of suicidality. This is a cross-sectional study which cannot establish causal relationships between exposures and outcomes.

Conclusions: This is the first epidemiological study in Andalusia on suicidality offering important results of clinical interest for suicide prevention.

BACKGROUND

Suicide is a very important public health problem. According to the World Health Organization (WHO) over 800,000 people die due to suicide every year. For each person who dies by suicide, more than 20 others attempt suicide. In fact, suicide attempts are an important risk factor for subsequent suicide (WHO, 2019). If relatives, friends and the community of the person who attempts suicide or commit suicide are taken into account, millions of people are affected worldwide by this phenomenon every year (Pitman *et al.*, 2014; Cerel *et al.*, 2019; WHO, 2019). A worldwide systematic review (Nock *et al.*, 2008b) showed that for adults, the lifetime prevalence of suicidal ideation ranged from 3.1-56%; suicide planning ranged between 0.9-19.5% and suicide attempts ranged between 0.4-5.1%. A recent meta-analysis of epidemiological studies about suicidal behavior of the general population in Europe developed by our research group concluded that for point prevalence, the rate for all suicidal behavior was 3.96% (2.37 – 5.56%), for death wishes 1.53% (0.95 – 2.12%), for suicidal ideation 5.28% (3.5 – 7.06%), and for suicidal attempt 0.63% (-0.09 – 1.36%) (Castillejos *et al.*, in press). The Table 1 details prevalence of suicidality and its different components in European studies.

Concerning risks factors for suicidality, data collected in epidemiological studies round the world indicate that suicidal behavior is associated with a variety of demographic, psychosocial and clinical factors (Nock *et al.*, 2008b; Simon, 2010). A well-established correlated factor for suicidal risk is the presence of mental disorders (Mann *et al.*, 2005; Bebbington *et al.*, 2009; Nock, 2009; Koyanagi *et al.*, 2015; Lara *et al.*, 2015). Substance abuse as a risk factor has been examined additionally in many studies (Miret *et al.*, 2013; Hardt *et al.*, 2015; Hiswals *et al.*, 2015;). Other factors, like age (Rancans *et al.*, 2016), gender (Gisle and Van Oyen, 2013), marital status (Forkman *et al.*, 2012), employment status (Economou *et al.*, 2013), adverse childhood experience (ten Have *et al.*, 2013; Yip *et al.*, 2015), stressful life event (Osvath *et al.*, 2004; Nock *et al.*, 2008a), poor physical health (Hawton and van Heeringen, 2009) or family psychiatric history (Saracli *et al.*, 2016) among others, also constitute risk factors for suicidality.

Spain has participated in the European Study of the Epidemiology of Mental Disorders [ESEMED] and presented results about prevalence for lifetime suicide ideation (4.4%) and attempts (1.5%) in the general population (Gabilondo *et al.*, 2007). Other community Spanish studies have found point prevalence of suicidal ideation of 2.3% in the city of Santander results from the European study “ODIN” (Casey *et al.*, 2008) and 6.5% on the island of Formentera (Gili-Planas *et al.*, 2001). Concerning risk factors for suicidality in Spain, there seems to be a significant relation being women, younger, lower education levels and having a mental disorders especially depression (Gabilondo *et al.*, 2007; Miret *et al.*, 2014). The data also suggest that the

factors associated with suicidality vary among age groups (Miret *et al.*, 2014). Despite these previous studies developed in Spain, little is known about the prevalence and associated factors of key behaviors related to suicide in the Spanish southern region of Andalusia, which is the largest autonomous community of Spain with a population of nearly 9 million inhabitants. Although, some studies, have been developed in this Spanish community, either concerning consumed suicide and its relationship with unemployment and the use of antidepressants (Alameda-Palacios *et al.*, 2014) or about suicide attempts attended by prehospital emergency services (Córdoba-Doña *et al.*, 2014; Mejías-Martín *et al.*, 2018; Moreno-Küstner *et al.*, 2019), none of them has been developed in the general population.

So, in response to the growing international interest in suicide prevention, from a public health perspective, and the importance to estimate prevalence data and calculate the impact of different risk domains from various geographical areas, we applied a community epidemiological approach to a wide and socio-economically diverse area in South of Spain, the PISMA-ep study (Epidemiological Study on Frequency and Risk Factors of Mental Disorders in Andalusia) that is the largest epidemiological mental health survey ever carried out in Andalusia. This research aims: 1) to estimate the prevalence of suicidality (including and differentiating a wide variety of suicidality components like death wish, self-harm ideas, suicidal thoughts, suicidal plan, and suicidal attempt); 2) to analyze a large number of demographic, psychosocial and clinical factors that could be associated to suicidal risk.

METHOD

Design and sample procedure

The PISMA-ep study is a cross-sectional household study conducted on a representative probability sample of non-institutionalized both male and female adult between 18-75 years old. Participants were selected from a multistage clustered sample, using different standard stratification levels. 1) Proportional stratification that took into account the geographical zones of western and eastern Andalusia; 2) In each geographical zone city size was classified under three categories: urban (over 10,000 inhabitants), intermediate (between 2,001 and 10,000 inhabitants) and rural (up to 2,000 inhabitants); 3) In addition, we also stratified each sampling area of the 8 provinces of Andalusia, and finally 4) Within each province a simple random method was used to select between one and 5 municipal entities (cities) for each type of locality size (urban, intermediate or rural). Then, using the same simple randomization method and taking into account the age and sex quotas, the census sections and the districts of each chosen location, the final sampling areas were also selected. These were people of both sexes, between 18 and

75 years old, who were interviewed if they lived in houses on predetermined street routes within the districts and census sections previously identified at random. Every fourth house of consecutive houses was called at. . The study interviews were held between 2013 and 2014, and went over a period of nearly a year. The PISMA-ep study was approved by the Research Ethics Committee of the University of Granada. A full description of the study methods can be found elsewhere (Cervilla *et al.*, 2016).

Assessment of suicidality

Non-lethal suicide behaviour was assessed with the suicidality module of the Spanish version of the Mini International Neuropsychiatric Interview (MINI-Plus 5.0) (Figure 1). The suicidality module was composed of six questions including: death wish, ideas of self-harm, suicidal thoughts, suicidal plans and attempts all in the past-month and also lifetime suicide attempts. We have calculated the prevalence rates for each of the above mentioned items.

For the associated factors analysis, the outcome variable suicidal risk is based on a MINI system that the subjects were categorized as being 'at risk of suicide' or 'no suicidal risk'. A "yes" answer on one of the six answered questions is considered a suicidal risk (Sheehan *et al.*, 1998).

Assessment of Associated Factors

Based on the literature about risk factors of suicidality and taking into account the purpose of our research, we also explored the following sets of potential correlates of suicidality: demographic, psychosocial and clinical factors. A full description of the methods has been registered elsewhere (Cervilla *et al.*, 2016).

Demographic factors such as gender, age, marital status, employment status, educational level, town size and province were collected using a questionnaire based on previous studies (Barona *et al.*, 1984; Bilbao and Seisdedos, 2004).

The following psychosocial factors were considered. Personal and social performance levels of functioning were assessed using the Personal and Social Performance Scale, PSP (Morosini *et al.*, 2000). Social support was measured with an inventory that evaluates what the person thinks about family and friends (Blaxter, 1990). Childhood experiences of physical, emotional or sexual abuse were assessed using the shortened version of the Childhood Trauma Questionnaire, CTQ (Fink *et al.*, 1995). Recent stressful life events were evaluated using a brief validated checklist (Brugha *et al.*, 1985). Personality traits such as neuroticism and impulsivity were evaluated using the Zuckerman-Kuhlman Personality Questionnaire, ZKPQ (Zuckerman, 2002).

Clinical variables such as family history of mental disorder and suicide in family members were assessed by an adaptation of the General Screening Questions and the Symptom Checklist of the Family Interview for Genetic Studies (NIMH, 2005). Presence of drugs dependence was assessed using the appropriate section of the MINI (Sheehan *et al.*, 1998). Alcohol consumption was assessed with the CAGE questionnaire (Ewing, 1984). Dependence of nicotine was evaluated by the Fagerström Test for Nicotine Dependence (Heatherton *et al.*, 1991).

Response rate

The homes originally selected which did not respond after 4 different attempts or did not have any member within the set age and gender range were replaced by the next available person within the predetermined route. A total of 70.8% of homes initially selected had to be replaced. 37.7% (N=2074) of homes chosen did not respond after all the attempts made and they had to be replaced due to a lack of response. Another third (33%) of homes originally chosen (N=1818) were replaced because there was nobody within the age, gender and educational range which was matched in the area.

Data quality

The error rates of all the provinces were far below the 1% acceptability rate. The study designs, field procedures and sample weightings are detailed elsewhere (Cervilla *et al.*, 2016).

Data analysis

Prevalence rates estimates of each item of suicidality (any items of suicide positive) were calculated with 95% confidence intervals. Explorative descriptive analyses including frequencies, percentages, mean and standard deviation values of all independent variables were calculated. Subsequently, we explored the univariate association between suicidal risk and different independent variables using *Chi square* test (two tails) and *Student's* test, according to the normal distribution of the data. After checking the significant finding for each risk factor, in the final multivariable regression model, all variables were entered simultaneously to examine independent effects and odds ratios (OR) are presented. Statistical significance was assumed when the *p* was <0.05. Descriptive and univariate analysis were applied using the SPSS 20 (SPSS, 2011). The software used for multivariate regression analysis was STATA 13 package (Statacorp, 2013).

RESULTS3

Sample and Response Rate

4507 residents in Andalusia were interviewed, 50.9% were male. Mean age was 42.8 (SD=15.22). A more detailed sample description can be found in Table 2 and a full description has been reported elsewhere (Cervilla *et al.*, 2016).

Prevalence of suicidality

Prevalence rate of suicidality was 6.4% (95% CI: 5.7-7.12). Regarding the prevalence of suicidality components, according to the items set by the MINI are: showed a death wish 4.4% (95% CI: 3.84-5.04); had ideas of self-harm 1.4% (95% CI: 1.07-1.77); had suicidal thoughts 2.4% (95% CI: 1.95-2.84); had a suicidal plan 1.1% (95% CI, 0.8-1.42); had attempted suicide 0.6% (95% CI, 0.41-0.88) and manifesting all during the month prior to the interview, and, lastly, 2.6% (95% CI, 2.13-3.06) reported to have previously had some sort of suicide attempt during his/her life.

Associated factor for suicidal risk

The univariate analysis showed that suicidal risk was associated with demographic factors such as female sex, older age, not having a stable partner, being unable to work because of illness or long-term disability, having unfinished primary educational level and to live in the province of Almeria. The psychosocial factors associated with suicidal risk are lower scores on social performance and social support, having had childhood abuse experiences (physical, emotional, and sexual), having a higher score on the stressful life events scale and a higher score on both, neuroticism and impulsivity traits. Suicidal risk was also associated with clinical factors including family history of mental disorder, any suicide in the family, drug dependence, higher alcohol consumption and nicotine dependence (Table 3).

A parsimonious multivariable model of association with suicidal risk was obtained including 3 demographic, 4 psychosocial and 3 clinical factors remained significantly correlated with suicidal risk: female, older age, not having a stable partner (being single/separated/divorced/widowed), less social support, physical childhood abuse, more stressful life events, higher neuroticism score, family history of mental disorder, drug dependence and nicotine dependence (Table 4). The largest OR among demographic factors was being single/separated/divorced/widowed [odds ratio (OR) 1.60, 95% CI 1.22-2.09, $p < 0.001$]. The psychosocial factor with greater power of association with suicidal risk was the experiences of physical abuse in childhood [odds ratio (OR) 2.43, 95% CI 1.66-3.56, $p < 0.001$]. Finally, drug dependence was the variable with the greatest association with suicidal risk among the different

clinical factors [odds ratio (OR) 2.96, 95% CI 1.70-5.16, $p < 0.001$], this last factor is also the one that shows the greatest OR for suicidal risk in whole multivariate model.

DISCUSSION

This cross-sectional study showed that suicidality defined as non-lethal suicide behaviour is a relatively highly prevalent situation in the Andalusian population, affecting over 1 in every 25 persons. We also found numerous potential risk factors of different domains such as demographics, psychological and clinical which appear to be associated with current suicidal risk. To the best of our knowledge, this study is the first epidemiological survey on the prevalence of suicidality in the region of Andalusia and one of very few in Spain. The main strength of the study is that it was performed on a large sample (N=4507) with a global response rate of 83.7% that is representative of the Andalusian population.

Definition of suicidality used

In our definition of suicidality, we included concepts that ranged from death wish, ideas of self-harm and suicidal thoughts, planning or attempts, and all which are included in the nomenclature of Silverman *et al.* (2007), of greater current recognition in the scientific community. We justify such a broad concept of suicidality given the seriousness of all such items and in accordance with the latest reports submitted by WHO on suicide prevention (2014, 2019). The latter suggest the difficulty of evaluating the real suicidal intentionality within each one of these measures due to the ambivalence and concealment surrounding suicidal behaviour. On the other hand, the inclusion of different levels, from the mildest (wish of death) to the most serious (suicide attempt), allows us to expand the range of possibilities, adapt preventive measures to people possibly at risk. We have also classified our outcome result of suicidal risk with a proper method following the MINI module of suicidality, because we wanted to analyse differences between the groups of persons who had any positive item of suicidality compared to the general population.

Prevalence of suicidality

We found a 1-month prevalence of 6.4% among adult persons with at least one point on the MINI Suicidality Module. A recent meta-analysis indicates an average prevalence of 3.96% for suicidality in Europe (Castillejos *et al.*, in press). Our prevalence would be above this figure. It is possible that the inclusion of more elements of suicidality in this study could imply an overestimation of prevalence if compared with other epidemiological studies, being difficult to establish equivalences and comparisons (Table 1). This could be due to a narrower focus used in

other studies. Death wish was the most prevalent element and attempted suicide was the less prevalent. On the whole, our study reports prevalence rates of suicidality and its components that generally compare favourably well with what could be expected from previous European reports (Table 1). Thus, prevalence of death wishes is higher if compared with Aschan *et al.* (2013) who found a much lower prevalence of 1.83 in the United Kingdom. Conversely, one of the more novel results is our prevalence figure of suicidal thoughts (2.4%) which is nearly fifty percent lower compared with the average of 5.28% found in the systematic review (Castillejos *et al.*, in press). Additionally, our prevalence figure for suicidal ideation is well within the range reported by an array of different European studies conducted during the last 10 years, that ranged between: 0.8 – 6.7% (Gili-Planas *et al.*, 2001; Ladwig *et al.*, 2008; Hintikka *et al.*, 2009; Michal *et al.*, 2010; Polatöz *et al.*, 2011; Forkman *et al.*, 2012; Economou *et al.*, 2013; Gisle and Van Oyen, 2013; McDonald, 2017). One explication concerning our lower rate of suicidal thoughts is that the suicidality module of the MINI included 2 questions concerning less severe suicidal behavior as “Think you would be better off dead or wish you were dead?” or “Want to harm yourself?”. So, in this situation, persons have more options to choose an answer according to their suicidal feeling, however in studies in which the only item is suicidal ideation, these people have to choose this option even when the feeling is less severe.

We have found 4 studies in European countries on current suicide attempts which report a prevalence range between 0.01 (McDonald, 2017) and 1.5% (Economou *et al.*, 2013), whilst our reported rate is one of 0.6% very similar to the average of the studies (0.63%) (Table 1).

Finally, regarding lifetime attempts, considered the most severe behavior of suicide, we have found 23 different reports from European studies ranging from 0.57% to 5.04% which places our 2.6% prevalence right in the median of this interval and very similar to the average of 2.9% (Castillejos *et al.*, in press). If compared with Spanish figures, our result exceeds figures from previous reports (Gabilondo *et al.*, 2007; Miret *et al.*, 2014). This can plausibly be attributable to regional differences, different timings (our survey occurred in the middle of a serious financial crisis) and different assessment tools. Indeed, relatively higher prevalence in our study when compared with other Spanish studies are in accordance with reports suggesting that economic crisis in Andalusia has had an impact on the increase of attempts of suicidal rates due to unemployment (Córdoba-Doña *et al.*, 2014; López-Bernal *et al.*, 2014). However, the relation between economic crisis and suicide are rather complex and open to controversy as reported by Masedo and Moreno-Küstner (2015). These data, nonetheless, provide an invitation to study those social factors such as the economic situation, unemployment or other cultural and demographic factors that could mediate higher rates of life prevalence of suicide attempts in

Andalusia if compared to the rest of Spain, with the aim of establishing adequate preventive measures adapted to each community.

Associated factor with suicidal risk

Demographic factors

Our results show that female gender associated with suicidal risk is in line with most others on suicidality (except for actual consummated suicide) (Gabilondo *et al.*, 2007; Nock *et al.*, 2008a; Borges *et al.*, 2010, Meltzer *et al.*, 2011; Gisle and Van Oyen, 2013; Hardt *et al.*, 2015; Hiswals *et al.*, 2015).

We found a trend for higher suicidal risk as participants' age increase. Previous studies have presented contradictory data. Thus, whilst some studies associated suicide behaviours with younger age groups (Bernal *et al.*, 2007; Nock *et al.*, 2008b; Scocco *et al.*, 2008; Hintikka *et al.*, 2009; Borges *et al.*, 2010; Kovess-Masfety *et al.*, 2011; Forkman *et al.*, 2012; Gisle and Van Oyen, 2013; Hiswals *et al.*, 2015; Lara *et al.*, 2015); others point towards an association of suicide behaviour with older age groups (Scocco *et al.*, 2008; Forkman *et al.*, 2012; Rancans *et al.*, 2016). Finally, there are others papers in which age showed no direct effects on suicidal risk (Hardt *et al.*, 2015).

Additionally, not having a partner (being single/separated/divorced/widowed) is also associated with people who present a higher suicidal risk as shown in a large number of previous reports (Weissman *et al.*, 1999; Kposowa, 2000; Bertolote *et al.*, 2005; Corcoran and Nagar, 2010; Meltzer *et al.*, 2011; Forkman *et al.*, 2012; Gisle and Van Oyen, 2013; Hiswals *et al.*, 2015; Yip *et al.*, 2015). This finding is supported by the notion that an intimate partner can act as a vehicle that facilitates greater social integration and more supportive social networks (Corcoran *et al.*, 2010).

Psychosocial variables

It is well-documented that different forms of child abuse significantly increases the risk for suicide (Felitti *et al.*, 1998; Bruffaerts *et al.*, 2015; Saracli *et al.*, 2016; Turecki and Brent, 2016) thus supporting our finding of an association between suicidal risk and physical child abuse (Hardt, 2008; Bruffaerts *et al.*, 2010; Borges *et al.*, 2010; ten Have *et al.*, 2013; Hardt *et al.*, 2015). Our finding is also in line with the hypothesis suggesting that early experience of violence like physical abuse, harsh physical punishment and other early experience of violence may lower the threshold to later act violently against oneself (Hardt *et al.*, 2008; Hardt *et al.*, 2015). Lower levels of social support and higher exposure to previous stressful life events are also associated with suicidal risk

as extensively reported earlier (Vijayakumar and Rajkumar, 1999; Phillips *et al.*, 2002; Osvath *et al.*, 2004; Chaqueta and Stiles 2007; Nock *et al.*, 2008b; You *et al.*, 2010; Kleiman *et al.*, 2012; Liu and Miller, 2014). It is striking that the variable sexual abuse in childhood had no significant relationship with suicidal behavior. It is likely that the sample size ($n = 48$) makes this association difficult. Finally, neuroticism has also previously been reported as associated to increased suicidal risk (Nordström *et al.*, 1995; Beautrais *et al.*, 1999; Fergusson *et al.*, 2003). This factor is generally higher especially in younger adults with serious intent, concurrent substance misuse and affective disorders (Brezo *et al.*, 2006). We suggest that clinical measuring of neuroticism, mainly in young people, can be of help as a preventative approach to suicide risk assessment.

Clinical variables

In our study, any suicide in the family and family history of mental disorders were both univariately associated with suicidal risk although only the effects of family history of mental disorders held significant as an independent variable associated with suicidal risk. This replicates previous reports (Kendler and Prescott, 2007; Nock *et al.*, 2008b). Indeed, living during the childhood and adolescence with a person suffering from a mental disorder could be a risk factor for future emotional problems that, in turn can trigger a higher degree of suicidal risk (Nock *et al.*, 2008b; Ayuso *et al.*, 2012). Similarly, drug dependence can be a sign of emotional problems that increase suicidality levels and many previous reports of an association between suicidality and drug or nicotine abuse exist (Kessler *et al.*, 1999; Breslau *et al.*, 2005; Gabilondo *et al.*, 2007; Kessler *et al.*, 2007; Bronisch *et al.*, 2008; Riala *et al.*, 2009; Meltzer *et al.*, 2011; Qin, 2011; Hardt *et al.*, 2015). Indeed, nicotine dependence is a variable strongly associated with different psychiatric disorders (Grant *et al.*, 2004; Pulay *et al.*, 2010), and the presence of any psychiatric disorder is a well established risk factor for suicidal behavior (Kessler, 2001; Nock *et al.*, 2008b; Yaworski *et al.*, 2011; Hardt *et al.*, 2015).

Limitations and strengths

There are limitations to this report. A first limitation is that the instrument employed to measure suicidality is a screening tool rather than a more in-depth diagnostic measure. Nevertheless, the suicidality module of the MINI has previously been used in suicidality studies both at clinical (Tajudeenet *et al.*, 2013; Guimarães *et al.*, 2014; Klug *et al.*, 2015) and general (Jeon *et al.*, 2014) population levels. Secondly, it has been suggested that when suicidality is enquired upon by an interviewer there is a tendency to infra-report it when compared with auto-administered suicidality reports (Turner *et al.*, 1998). Conversely, questions on death wish have been shown to

generate higher prevalence estimates for suicidality than more difficult probing questions on “seriously considering suicide” (Scocco and de Leo, 2002; Nock *et al.*, 2008b). Thirdly, it could be considered a limitation of the study, to include a suicide attempt as part of the current suicide risk, however, a history of attempted suicide was found to have high sensitivity (0.80) and very high specificity (0.97) in differentiating between suicidal and non-suicidal adult inpatients (Osman, 2001). In addition, it is the strongest and most robust predictor of suicide attempts and suicide completion in mental disorder (Harris, 1997; Hawton, 2009) and it is a risk factor highly ranked for suicide assessed by most practitioners (Osman, 2001; Truant, 1991). Moreover, we have used it as proposed by the developers of MINI and the aforementioned rationales, the item of previous suicide attempt of this MINI module was therefore included as a part of the current suicide risk. However, in our study we have not classified suicide risk according to severity of suicidal manifestations proposed by Sheean *et al.*(1998). Fourth, although we include a wide array of correlating factors, there are many other potential correlates of suicidal risk that were not included. Finally, this is a cross-sectional study which cannot establish causal relationships between exposures and outcomes, which weakens its ability to establish preventive recommendations based on the predictability of the model.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by a grant from the Consejería de Economía, Innovación y Ciencia de la Junta de Andalucía (10-CTS-6682).

Authors' contributions

Experiments conceived and designed by: BMK, BG, JC. Experiments performed by: BG, JC. Data analyzed by: JC, PH. Paper written by: PH, BMK, JC. Final draft of the manuscript read and agreed upon by: PH, BMK, BG, JC.

Acknowledgements: None

The authors declare no conflicts of interest.

References

- Alameda-Palacios, J., Ruiz-Ramos, M., García-Robredo, B., 2014. Suicide, antidepressant prescription and unemployment in Andalusia (Spain). *Gaceta Sanitaria* 28, 309-312. <https://doi.org/10.1016/j.gaceta.2013.12.005>.
- Aschan, L., Goodwin, L., Cross, S., Moran, P., Hotopf, M., Hatch, S.L., 2013. Suicidal behaviours in South East London: Prevalence, risk factors and the role of socio-economic status. *Journal of Affective Disorders* 150, 441-449. <https://doi.org/10.1016/j.jad.2013.04.037>.
- Atay, I.M., Eren, I., Gündoğar, D., 2012. The Prevalence of Death Ideation and Attempted Suicide and the Associated Risk Factors in Isparta, Turkey. *Turkish Journal of Psychiatry*, 23, 89-98.
- Ayuso, J., Baca-García, E., Bobes, J., Giner, J., Giner, L., Pérez, V., Sáiz, P.A., Saiz Ruiz, J., Grupo RECOMS, 2012. Recommendations for the prevention and management of suicidal behaviour. *Revista Psiquiatría y Salud Mental*, 5, 8-23. <https://doi.org/10.1016/j.rpsm.2012.01.001>.
- Barona, A., Reynolds, C.R., Chastain, R., 1984. A demographically based index premorbid intelligence for the WAIS-R. *Journal of Consulting and Clinical Psychology*, 52, 885-887.
- Beautrais, A.L., Joyce, P.R., Mulder, R.T., 1999. Personality traits and cognitive styles as risk factors for serious suicide attempts among young people. *Suicide & life-threatening behavior* 29, 37-47.
- Bebbington, P.E., Cooper, C., Minot, S., Brugha, T.S., Jenkins, R., Meltzer, H., Dennis, M., 2009. Suicide Attempts, Gender, and Sexual Abuse: Data From the 2000 British Psychiatric Morbidity Survey. *The American Journal of Psychiatry* 166, 1135-1140. <https://doi.org/10.1176/appi.ajp.2009.09030310>.
- Bernal, M., Haro, J.M., Bernert, S., Brugha, T., de Graaf, R., Bruffaerts, R., Lépine, J.P., de Girolamo, G., Vilagut, G., Gasquet, I., Torres, J.V., Kovees, V., Heider, D., Neeleman, J., Kessler, R., Alonso, J., 2007. Risk factors for suicidality in Europe: Results from the ESEMED study. *Journal of Affective Disorders* 101, 27-34. <https://doi.org/10.1016/j.jad.2006.09.018>.
- Bertolote, J.M., Fleischmann, A., de Leo, D., Bolhari, J., Botega, N., de Silva, D., Thanh, H.T.T., Philips, M., Schlegel, L., Värnik, A., Vijayakumar, L., Wasserman, D., 2005. Suicide attempts, plans, and ideation in culturally diverse sites: the WHO SUPRE-MISS community survey. *Psychological Medicine* 35, 1457-1465. <https://doi.org/10.1017/S0033291705005404>.
- Bilbao, A., Seisdedos, N., 2004. Eficacia de una fórmula de estimación de la inteligencia premórbida en la población española. *Revista de Neurología* 38, 431-434.
- Blaxter, M., 1990. *Health and Lifestyles*. Routledge: London.

- Borges, G., Nock, M.K., Haro Abad, J.M., Hwang, I., Sampson, N.A., Alonso, J., Andrade, L.H., Angermeyer, M.C., Beautrais, A., Bromet, E., Bruffaerts, R., de Girolamo, G., Florescu, S., Gureje, O., Hu, C., Karam, E.G., Kovess-Masfety, V., Lee, S., Levinson, D., Medina-Mora, M.E., Ormel, J., Posada-Villa, J., Sagar, R., Tomov, T., Uda, H., Williams, D.R., Kessler, R.C., 2010. Twelve Month Prevalence of and Risk Factors for Suicide attempts in the WHO World Mental Health Surveys. *Journal of Clinical Psychiatry* 71, 1617-1628. <https://doi.org/10.4088/JCP.08m04967blu>.
- Boyd, A., Van de Velde, S., Vilagut, G., de Graaf, R., O'Neill, S., Florescu, S., Alonso, J., Kovess-Masfety, V., 2015. Gender differences in mental disorders and suicidality in Europe: Results from a large cross-sectional population-based study. *Journal of Affective Disorders* 173, 245-254. <https://doi.org/10.1016/j.jad.2014.11.002>.
- Breslau, N., Schultz, L.R., Johnson, E.O., Peterson, E.L., Davis, G.C., 2005. Smoking and the Risk of Suicidal Behavior. *Archives of General Psychiatry* 62, 328-334. <https://doi.org/10.1001/archpsyc.62.3.328>.
- Brezo, J., Paris, J., Turecki, G., 2006. Personality traits as correlates of suicidal ideation, suicide attempts, and suicide completions: A systematic review. *Acta Psychiatrica Scandinavica* 113, 180-206. <https://doi.org/10.1111/j.1600-0447.2005.00702.x>.
- Bronisch, T., Höfler, M., Lieb, R., 2008. Smoking predicts suicidality: findings from a prospective community study. *Journal of Affective Disorders* 108, 135-145. <https://doi.org/10.1016/j.jad.2007.10.010>.
- Bruffaerts, R., Demyttenaere, K., Borges, G., Haro, J.M., Chiu, W.T., Hwang, I., Karam, E.G., Kessler, R.C., Sampson, N., Alonso, J., Andrade, L.H., Angermeyer, M., Benjet, C., Bromet, E., de Girolamo, G., de Graaf, R., Florescu, S., Gureje, O., Horiguchi, I., Hu, C., Kovess, V., Levinson, D., Posada-Villa, J., Sagar, R., Scott, K., Tsang, A., Vassilev, S.M., Williams, D.R., Nock, M.K., 2010. Childhood adversities as risk factors for onset and persistence of suicidal behaviour. *The British Journal of Psychiatry* 197, 20-27. <https://doi.org/10.1192/bjp.bp.109.074716>.
- Bruffaerts, R., Kessler, R.C., Demyttenaere, K., Bonnewyn, A., Nock, M.K., 2015. Examination of the population attributable risk of different risk factor domains for suicidal thoughts and behaviors. *Journal of Affective Disorders* 187, 66-72. <https://doi.org/10.1016/j.jad.2015.07.042>.
- Brugha, T., Bebbington, P., Tennant, C., Hurry, J., 1985. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychological Medicine* 15, 189-194. <https://doi.org/10.1017/S003329170002105X>.

- Casey, P., Dunn, G., Kelly, B.D., Lehtinen, V., Dalgard, O.S., Dowrick, C., Ayuso-Mateos, J.L., 2008. The prevalence of suicidal ideation in the general population: results from the Outcome of Depression International Network (ODIN) study. *Social Psychiatry and Psychiatric Epidemiology* 43, 299-304. <https://doi.org/10.1007/s00127-008-0313-5>.
- Castillejos, C., Huertas, P., Martín, P., Moreno-Küstner, B., Prevalence of suicidal behaviour in the european general population: a systematic review and meta analysis. *Archives of Suicide Research* (In press).
- Cerel, J., Brown, M., Maple, M., Singleton, M., van de Venne, J., Moore, M., 2019. How Many People Are Exposed to Suicide? Not Six. *Suicide and Life-Threatening Behavior* 49, 529-534. <https://doi.org/10.1111/sltb.12450>.
- Cervilla, J.A., Ruiz, I., Rodríguez-Barranco, M., Rivera, M., Ibáñez-Casas, I., Molina, E., Valmisa, E., Carmona-Calvo, J., Moreno-Küstner, B., Muñoz-Negro, J.E., Ching-López, A., Gutiérrez, B., 2016. Protocol and methodology of Study epidemiological mental health in Andalusia: PISMA-ep. *Revista de Psiquiatría y Salud Mental* 9, 185-194. <https://doi.org/10.1016/j.rpsm.2015.11.004>.
- Chaqueta, A.P., Stiles, T.C., 2007. The relationship between psychological buffers, hopelessness, and suicidal ideation: Identification of protective factors. *Crisis* 28, 67-73. <https://doi.org/10.1027/0227-5910.28.2.67>.
- Córdoba-Doña, J.A., San Sebastián, M., Escolar-Pujolar, A., Martínez-Faure, J., Gustafsson, P.E., 2014. Economic crisis and suicidal behavior: the role of unemployment, sex and age in Andalusia, Southern Spain. *International Journal for Equity in Health* 13, 55. <http://www.equityhealthj.com/content/13/1/55>.
- Corcoran, P., Nagar, A., 2010. Suicide and marital status in Northern Ireland. *Social Psychiatry and Psychiatric Epidemiology* 45, 795-800. <https://doi.org/10.1007/s00127-009-0120-7>.
- Economou, M., Angelopoulos, E., Peppou, L.E., Souliotis, K., Tzavara, C., Kontoangelos, K., Madianos, M., Stefanis, C., 2016. Enduring financial crisis in Greece: prevalence and correlates of major depression and suicidality. *Social Psychiatry and Psychiatric Epidemiology* 51, 1015-1024. <http://link.springer.com/10.1007/s00127-016-1238-z>.
- Encrenaz, G., Kovess-Masféty, V., Gilbert, F., Lagarde, E., Mishara, B., Messiah, A., 2012. Lifetime risk of suicidal behaviors and communication to a health professional about suicidal ideation. Results from a large survey of the French adult population. *Crisis* 33, 127-136. <https://doi.org/10.1027/0227-5910/a000113>.
- Ewing, J.A., 1984. Detecting Alcoholism. The CAGE Questionnaire. *JAMA* 252, 1905-1907. <https://doi.org/10.1001/jama.252.14.1905>.

- Felitti, V.J., Anda, R.F., Nordenberg, D., Williamson, D.F., Spitz, A.M., Edwards, V., Koss, M.P., Marks, J.S., 1998. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine* 14, 245-258. [https://doi.org/http://dx.doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/http://dx.doi.org/10.1016/S0749-3797(98)00017-8).
- Fergusson, D.M., Beautrais, A.L., Horwood, L.J., 2003. Vulnerability and resiliency to suicidal behaviours in young people. *Psychological Medicine* 33, 61-73. <https://doi.org/10.1017/S0033291702006748>.
- Fink, L.A., Bernstein, D., Handelsman, L., Foote, J., Lovejoy, M., 1995. Initial reliability and validity of the childhood trauma interview: a new multidimensional measure of childhood interpersonal trauma. *The American Journal of Psychiatry* 152, 1329-1335.
- Flaherty, C., 2019. How Many People Are Exposed to Suicide? Not Six, Suicide and Life-Threatening Behavior 49, 529-534. <https://doi.org/10.1111/sltb.12450>.
- Forkmann, T., Brähler, E., Guggel, S., Glaesmer, H., 2012. Prevalence of Suicidal Ideation and Related Risk Factors in the German General Population. *The Journal of Nervous and Mental Disease* 200, 401-405. <https://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00005053-201205000-00006>. <https://doi.org/10.1097/NMD.0b013e31825322cf>.
- Gabilondo, A., Alonso, J., Pinto-Meza, A., Vilagut, G., Fernández, A., Serrano-Blanco, A., Almansa, J., Codony, M., Haro, J.M., 2007. Prevalencia y factores de riesgo de las ideas, planes e intentos de suicidio en la población general española. Resultados del estudio ESEMeD. *Medicina Clínica* 129, 494-500. <https://doi.org/10.1157/13111370>.
- Gili-Planas, M., Roca-Bennasar, M., Ferrer-Pérez, V., Bernardo-Arroyo, M., 2001. Suicidal Ideation, Psychiatric Disorder, and Medical Illness in a Community Epidemiological Study. *Suicide and Life-Threatening Behavior* 31, 207-213.
- Gisle, L., Van Oyen, H., 2013. Household composition and suicidal behaviour in the adult population of Belgium. *Social Psychiatry and Psychiatric Epidemiology* 48, 1115-1124. <https://doi.org/10.1007/s00127-012-0621-7>.
- Grant, B.F., Hasin, D.S., Chou, S.P., Stinson, F.S. Dawson, D.A., 2004. Nicotine Dependence and Psychiatric Disorders in the United States, results from the national epidemiologic survey on alcohol and related conditions. *Archives of General Psychiatry* 61, 1107-1115. <https://doi.org/10.1001/archpsyc.61.11.1107>.
- Guimarães, P.M., Passos, S.P., Calvet, G.A., Hokerberg, Y.H., Lessa, J.L., de Andrade, C.A., 2014. Suicide risk and alcohol and drug abuse in outpatients with HIV infection and Chagas

disease. *Revista Brasileira de Psiquiatria* 36, 131-137. <https://doi.org/10.1590/1516-4446-2013-1219>.

Hardt, J., Bernet, S., Matschinger, H., Angermeier, M.C., Vilagut, G., Bruffaerts, R., de Girolamo, G., de Graaf, R., Haro, J.M., Kovess, V., Alonso, J., 2015. Suicidality and its relationship with depression, alcohol disorders and childhood experiences of violence: Results from the ESEMeD study. *Journal of Affective Disorders* 175, 168-174. <https://doi.org/10.1016/j.jad.2014.12.044>.

Hardt, J., Sidor, A., Nickel, R., Kappis, B., Petrak, F., Egle, U.T., 2008. Childhood Adversities and Suicide Attempts: A Retrospective Study. *Journal of Family Violence* 23, 713-718.

Harris E., Barraclough B., (1997). Suicide as an outcome for mental disorders: a meta-analysis. *British Journal of Psychiatry* 170, 205–228. <https://doi.org/10.1192/bjp.170.3.205>.

Hawton, K., van Heeringen, K., 2009. Suicide. *The Lancet* 373, 1372-1381. [https://doi.org/10.1016/S0140-6736\(09\)60372-X](https://doi.org/10.1016/S0140-6736(09)60372-X).

Heatherton, T.F., Kozlowski, L.T., Frecker, R.C., Fagerström, K.O., 1991. The Fagerström Test for Nicotine Dependence: a revision of the Fagerström Tolerance Questionnaire. *British Journal of Addiction* 86, 1119-1127.

Hintikka, J., Koivumaa-Honkanen, H., Lehto, S.M., Tolmunen, T., Honkalampi, K., Haatainen, K., Viinamäki, H., 2009. Are factors associated with suicidal ideation true risk factors? A 3-year prospective follow-up study in a general population. *Social Psychiatry and Psychiatric Epidemiology* 44, 29-33. <https://doi.org/10.1007/s00127-008-0401-6>.

Hiswals, A.S., Guilagaber, G., Wijk, K., Öberg, P., Soares, J., Macassa, G., 2015. Employment Status and Suicidal Ideation during Economic Recession. *Health Science Journal* 9, 1-13. <https://www.hsj.gr/archive>.

Jeon, H.J., Lee, C., Fava, M., Mischoulon, D., Shim, E.J., Heo, J.Y., Choi, H., Park, J.H., 2014. Childhood trauma, parental death, and their co-occurrence in relation to current suicidality risk in adults: a nationwide community sample of Korea. *The Journal of Nervous and Mental Disease* 202, 870-876. <https://doi.org/10.1097/NMD.0000000000000217>.

Kendler, K.S., Prescott, C.A., 2007. Genes, environment, and psychopathology: understanding the causes of psychiatric and substance use disorders. New York, NY: Guilford Press.

Kessler, R.C., Berglund, P.A., Bruce, M.L., Koch, J.R., Laska, E.M., Leaf, P.J., Manderscheid, R.W., Rosenheck, R.A., Walters, E.E., Wang, P.S., 2001. The prevalence and correlates of untreated serious mental illness. *Health services research* 36, 987-1007.

Kessler, R.C., Berglund, P.A., Borges, G., Castilla-Puentes, R.C., Glantz, M.D., Jaeger, S.A., Merikangas, K.R., Nock, M.K., Russo, L.J., Stang, P.E., 2007. Smoking and suicidal behaviors in

- the National Comorbidity Survey-Replication. *The Journal of Nervous and Mental Disease* 195, 369-377.
- Kessler, R.C., Borges, G., Walters, E.E., 1999. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Archives of General Psychiatry* 56, 617-626. <https://doi.org/10.1001/archpsyc.56.7.617>.
- Kleiman, E., Riskind, J., Schaefer, K., Weingarden, H., 2012. The moderating role of social support on the relationship between impulsivity and suicide risk. *Crisis* 33, 273-279. <https://doi.org/10.1027/0227-5910/a000136>.
- Klug, S.M., Dias, L., Coiro, B., 2015. AIDS Care: Psychological and Socio-medical Aspects of AIDS/HIV. *AIDS Care* 26, 1379-1382. <https://doi.org/10.1080/09540121.2014.913767>.
- Kovess-Masfety, V., Boyd, A., Haro, J.M., Bruffaerts, R., Villagut, G., Lépine, J.P., Alonso, J., 2011. High and low suicidality in Europe: a fine-grained comparison of France and Spain within the ESEMeD surveys. *Journal of Affective Disorders* 133, 247-256. <https://doi.org/10.1016/j.jad.2011.04.014>.
- Koyanagi, A., Stickley, A., Haro, J.M., 2015. Subclinical psychosis and suicidal behavior in England: Findings from the 2007 Adult Psychiatric Morbidity Survey. *Schizophrenia Research* 168, 62-67. <https://doi.org/10.1016/j.schres.2015.07.041>.
- Kposowa, A.J., 2000. Marital status and suicide in the National Longitudinal Mortality Study. *Journal of Epidemiology and Community Health* 54, 254-261. <https://doi.org/10.1136/jech.54.4.254>.
- Ladwig, K.H., Klupsch, D., Ruf, E., Meisinger, C., Baumert, J., Erazo, N., Wichmann, H.E., 2008. Sex- and age-related increase in prevalence rates of death wishes and suicidal ideation in the community: Results from the KORA-F3 Augsburg Study with 3154 men and women, 35 to 84 years of age. *Psychiatry Research* 161, 248-252. <https://doi.org/10.1016/j.psychres.2008.03.011>.
- Lara, E., Olaya, B., Garin, N., Ayuso-Mateos, J.L., Miret, M., Moneta, V., Haro, J.M., 2015. Is cognitive impairment associated with suicidality? A population-based study. *European Neuropsychopharmacology* 25, 203-213.
- Liu, R.T., Miller, I., 2014. Life events and suicidal ideation and behavior: A systematic review. *Clinical Psychology Review* 34, 181-192.
- López Bernal, L., Gasparrini, A., Artundo, C., McKee, M., 2014. The effect of the late 2000s financial crisis on suicides in Spain: an interrupted time-series analysis. *European Journal of Public Health*, 24, 184-185. <https://doi.org/10.1093/eurpub/ckt215>.
- Mann, J., Apter, A., Bertolome, J., Beautrais, A., Currier D., Haas, A., Hegerl, U., Lonngvisst, J., Malone, K., Marusic, A., Mehlum, L., Patton, G., Phillips, M., Rutz, W., Rihmer, Z., Schmidtke, A.,

- Shaffer, D., Silverman, M., Takahashi, Y., Varnik, A., Wasserman, D., Yip, P., Hendin, H., 2005. Suicide Prevention Strategies. *JAMA*, 294-297. <https://doi.org/10.1001/jama.294.16.2064>.
- Masedo-Gutierrez, A., Moreno-Küstner, B., 2015. Economic crisis and mortality by suicide: two concepts hard to link. *European Journal of Public Health*, 25, 900-901. <https://doi.org/doi:10.1093/eurpub/ckv080>.
- McDonald, K.C., Saunders, K.E.A., Geddes, J.R., 2017. Sleep problems and suicide associated with mood instability in the Adult Psychiatric Morbidity Survey, 2007. *Australian & New Zealand Journal of Psychiatry* 51, 822-828. <https://doi.org/10.1177/0004867416687398>.
- Mejías-Martín, Y., Martí-García, C., Rodríguez-Mejías, C., Valencia-Quintero, J.P., García-Caro, M.P., Luna, J., 2018. Suicide attempts in Spain according to prehospital healthcare emergency records, *PLoS One* 13, <https://doi.org/10.1371/journal.pone.0195370>.
- Meltzer, H., Bebbington, P., Brugha, T., Jenkins, R., McManus, S., Dennis, M.S., 2001. Personal debt and suicidal ideation. *Psychological Medicine* 41, 771-778. <https://doi.org/10.1017/S0033291710001261>.
- Michal, M., Wiltink, J., Till, Y., Wild, P.S., Münzel, T., Blankenberg, S., Beutel, M.E., 2010. Type-D personality and depersonalization are associated with suicidal ideation in the German general population aged 35-74: Results from the Gutenberg Heart Study. *Journal of Affective Disorders* 125, 227-233. <https://doi.org/10.1016/j.jad.2010.02.108>.
- Miret, M., Ayuso-Mateos, J.L., Sánchez-Moreno, J., Vieta, E., 2013. Depressive disorders and suicide: Epidemiology, risk factors, and burden. *Neuroscience and Biobehavioral Reviews* 37, 2372-2374. <https://doi.org/10.1016/j.neubiorev.2013.01.008>.
- Miret, M., Caballero, F.F., Huerta-Ramírez, R., Moneta, M.V., Olaya, B., Chatterji, S., Haro, J.M., Ayuso-Mateos, J.L., 2014. Factors associated with suicidal ideation and attempts in Spain for different age groups. Prevalence before and after the onset of the economic crisis. *Journal of Affective Disorders* 163, 1-9. <https://doi.org/10.1016/j.jad.2014.03.045>.
- Moreno-Küstner, B., del Campo-Ávila, J., Ruíz-Ibáñez, J., Martínez-García, A., Castro-Zamudio, S., Ramos-Jiménez, G., Guzmán-Parra, J., 2019. Epidemiology of Suicidal Behavior in Malaga (Spain): An Approach From the Prehospital Emergency Service, *Frontiers in Psychiatry* 10, 1-7. <https://doi.org/10.3389/fpsy.2019.00111>.
- Morosini, P.L., Magliano, L., Brambilla, L., Ugolini, S., Pioli, R., 2000. Development, reliability and acceptability of a new version of the DSM-IV Social and Occupational Functioning Assessment Scale (SOFAS) to assess routine social functioning. *Acta Psychiatrica Scandinavica* 101, 323-329.

- NIMH, 2005. Genetics Initiative. Family Interview for genetic studies 4.0 BP. https://www.nimhgenetics.org/interviews/figs/figs_4.0_bp.php.
- Nock, M.K., 2009. Why Do People Hurt Themselves? New Insights Into the Nature and Functions of Self-Injury. *Current Directions in Psychological Science* 18, 78-83. <https://doi.org/10.1111/j.1467-8721.2009.01613.x>.
- Nock, M.K., Borges, G., Bromet, E.J., Alonso, J., Angermeyer, M., Beautrais, A., Bruffaerts, R., Chiu, W.T., de Girolamo, G., Gluzman, S., de Graaf, R., Gureje, O., Haro, J.M., Huang, Y., Karam, E., Kessler, R.C., Lepine, J.P., Levinson, D., Medina-Mora, M.E., Ono, Y., Posada-Villa, J., Williams, D.R., 2008a. Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *British Journal of Psychiatry* 192, 98-105. <https://doi.org/10.1192/bjp.bp.107.040113>.
- Nock, M.K., Borges, G., Bromet, E.J., Cha, C.B., Kessler, R.C., Lee, S., 2008b. Suicide and suicidal behavior. *Epidemiologic Reviews* 30, 133-154. <https://doi.org/10.1093/epirev/mxn002>.
- Nordström, P., Schalling, D., Asberg, M., 1995. Temperamental vulnerability in attempted suicide. *Acta Psychiatrica Scandinavica* 92, 155-160. <https://doi.org/10.1111/j.1600-0447.1995.tb09560.x>.
- O'Neill, S., Ferry, F., Murphy, S., Corry, C., Bolton, D., Devine, B., Ennis, E., Bunting, B., 2014. Patterns of Suicidal Ideation and Behavior in Northern Ireland and Associations with Conflict Related Trauma. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0091532>.
- Osman, A., Bagge C.L., Gutierrez P.M., Konick L.C., Kopper B.A., Barrios F.X., 2001. The Suicidal Behaviors Questionnaire-Revised (SBQ-R): validation with clinical and nonclinical samples. *Assessment*, 8(4), 443-454.
- Osvath, P., Vörös, V., Fekete, S., 2004. Life Events and Psychopathology in a Group of Suicide Attempters. *Psychopathology* 37, 36-40. <https://doi.org/10.1159/000077018>.
- Package for the Social Sciences (SPSS) Statistics for Windows, Version 20.0, Armonk, NY: IBM Corp, 2011. IBM Corp.
- Phillips, M.R., Yang, G., Zhang, Y., Wang, L., Ji, H., Zhou, M., 2002. Risk factors for suicide in China: a national case-control psychological autopsy study. *The Lancet* 360, 1728-1736. [http://doi.org/10.1016/S0140-6736\(02\)11681-3](http://doi.org/10.1016/S0140-6736(02)11681-3).
- Pitman, A., Osborn, D., King, M., Erlangsen, A., 2014. Effects of suicide bereavement on mental health and suicide risk. *The Lancet Psychiatry* 1, 86-94. [https://doi.org/10.1016/S2215-0366\(14\)70224-X](https://doi.org/10.1016/S2215-0366(14)70224-X).
- Polatöz, Ö., Kuğu, N., Doğan, O., Akyüz, G., 2011. The Prevalence of Suicidal Behaviour and Its Correlation with Certain Sociodemographic Variables in Sivas Province. *Düşünen Adam The*

- Journal of Psychiatry and Neurological Sciences 24, 13-23.
<https://doi.org/10.5350/DAJPN2011240102>.
- Pulay, A.J., Stinson, F.S., Ruan, W.J., Smith, S.M., Pickering, R.P., Dawson, D.A., Grant, B.F., 2010. The relationship of DSM-IV personality disorders to nicotine dependence-results from a national survey. *Drug and Alcohol Dependence* 108, 141-145.
<https://doi.org/10.1016/j.drugalcdep.2009.12.004>.
- Qin, P., 2011. The impact of psychiatric illness on suicide: Differences by diagnosis of disorders and by sex and age of subjects. *Journal of Psychiatric Research* 45, 1445-1452.
<https://doi.org/10.1016/j.jpsychires.2011.06.002>
- Rancans, E., Pulmanis, T., Taube, M., Springe, L., Velika, B., Pudule, I., Grinberga, D., 2016. Prevalence and sociodemographic characteristics of self-reported suicidal behaviours in Latvia in 2010: A population-based study. *Nordic Journal of Psychiatry* 70, 195-201.
<https://doi.org/10.3109/08039488.2015.1077887>.
- Riala, K., Taanila, A., Hakko, H., Räsänen, P., 2009. Longitudinal smoking habits as risk factors for early-onset and repetitive suicide attempts: the Northern Finland 1966 Birth Cohort study. *Annals of Epidemiology* 19, 329-335. <https://doi.org/10.1016/j.annepidem.2008.12.004>.
- Saracli, O., Atasoy, N., Senormanci, O., Atik, L., Acikgoz, H.O., Dogan, V., Sankir, H., Köktürk, F., Öresl, S., 2016. Childhood trauma and suicide risk in the population living in Zonguldak Province. *Asia-Pacific Psychiatry* 8, 136-144. <https://doi.org/10.1111/appy.12214>.
- Scocco, P., de Girolamo, G., Vilagut, G., Alonso, J., 2008. Prevalence of suicide ideation, plans, and attempts and related risk factors in Italy: Results from the European Study on the Epidemiology of Mental Disorders-World Mental Health study. *Comprehensive Psychiatry* 49, 13-21. <https://doi.org/10.1016/j.comppsy.2007.08.004>.
- Scocco, P., de Leo, D., 2002. One-year prevalence of death thoughts, suicide ideation and behaviours in an elderly population. *International Journal of Geriatric Psychiatry* 17, 842-846.
<https://doi.org/10.1002/gps.69>.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., Herqueta, T., Baker, R., Dunbar, G.C., 1998. The Mini-International Neuropsychiatric Interview (M.I.N.I.). The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* 59, 22-33.
- Silverman, M. M., Berman, A. L., Sanddal, N. D., O'carroll, P. W., Joiner, T. E., 2007. Rebuilding the tower of Babel: a revised nomenclature for the study of suicide and suicidal behaviors. Part 1: Background, rationale, and methodology. *Suicide Life Threat.Behav.*, 37, 248-263.

- Simon, R., 2010. Preventing Patient Suicide: Clinical Assessment and Management. American Psychiatric Publishing, Inc: Washington, D.C.
- Statacorp, 2012. Stata Statistical Software. Release 13. College Station: Texas.
- Tajudeenet, F., Dupe, M., Jika, M., Bolakale, S., 2013. Suicide risk among adults with epilepsy in Kaduna, Nigeria. *General Hospital Psychiatry* 35, 517-520.
- Ten Have, M., van Dorsselaer, S., de Graaf, R., 2013. Prevalence and risk factors for first onset of suicidal behaviors in the Netherlands Mental Health Survey and Incidence Study-2. *Journal of Affective Disorders* 147, 205-211. <https://doi.org/10.1016/j.jad.2012.11.005>.
- Turecki, G., Brent, D.A., 2016. Suicide and suicidal behavior. *The Lancet* 387, 1227-1239. [https://doi.org/10.1016/S0140-6736\(15\)00234-2](https://doi.org/10.1016/S0140-6736(15)00234-2).
- Turner, C.F., Ku, L., Rogers, S.M., Lindberg, L.D., Pleck, J.H., Sonenstein, F.L., 1998. Adolescent sexual behavior, drug use, and violence: increased reporting with computer survey technology. *Science* 280, 867-873. <https://doi.org/10.1126/science.280.5365.867>.
- Vijayakumar, L., Rajkumar, S., 1999. Are risk factors for suicide universal? A case-control study in India. *Acta Psychiatrica Scandinavica* 99, 407-411. <https://doi.org/10.1111/j.1600-0447.1999.tb00985.x>.
- Truant G.S., O'Reilly R., Donaldson L., 1991. How psychiatrists weigh risk factors when assessing suicide risk. *Suicide Life and Threat Behavior* 21(2), 106–114.
- Weissman, M.M., Bland, R.C., Canino, G.J., Greenwald, S., Hwu, H.-G., Joyce, P.R., Karam, E.G., Lee, C.-K., Lellouch, J., Lepine, J.P., Newman, S.C., Rubio-Stipec, M., Wells, J.E., Wickramaratne, P.J., Wittchen, H.U., Yeh, E.-K., 1999. Prevalence of suicide ideation and suicide attempts in nine countries. *Psychological Medicine* 29, 9-17. <https://doi.org/10.1017/S0033291798007867>.
- World Health Organization, 2019. National suicide prevention strategies: progress, examples and indicators. https://www.who.int/mental_health/suicide-prevention/national_strategies_2019/en/.
- Yaworski, D., Robinson, J., Sareen, J., Bolton, J.M., 2011. The Relation Between Nicotine Dependence and Suicide Attempts in the General Population. *Canadian Journal of Psychiatry* 56, 161. <https://doi.org/10.1177/070674371105600306>.
- Yip, P.S.F., Yousuf, S., Chan, C.H., Yung, T., Wu, K.C.C., 2015. The roles of culture and gender in the relationship between divorce and suicide risk: A meta-analysis. *Social Science and Medicine* 128, 87-94. <https://doi.org/10.1016/j.socscimed.2014.12.034>.

You, S., Van Orden, K.A., Van, Conner, K.R.K., 2011. Social connections and suicidal thoughts and behavior. *Psychology of Addictive Behaviors: Journal of the Society of Psychologists in Addictive Behaviors* 25, 180-184. <https://doi.org/10.1037/a0020936>.

Zuckerman, M., 2002. Zuckerman-Kuhlman Personality Questionnaire (ZKPQ): an alternative five factorial model. In *Big five assessment* (ed. B. DeRaad & M. Perugini), pp.376-396. Hogrefe & Huber. Zuckerman, M: Seattle, WA.

Journal Pre-proof

Figure 1. Suicidality module of MINI (Mini International Neuropsychiatric Interview)

C. SUICIDALITY

In the past month did you:

- C1. Think you would be better off dead or wish you were dead?
- C2. Want to harm yourself?
- C3. Think about suicide?
- C4. Have a suicide plan?
- C5. Attempt suicide?
- C6. In your lifetime?

Did you ever make a suicide attempt?

IS AT LEAST 1 OF THE ABOVE CODED YES?
IF YES, SPECIFY THE LEVEL OF SUICIDE RISK AS FOLLOW:

No Yes
SUICIDE RISK CURRENT

Table 1. Prevalence rates of suicidality in Europe

First Author	Pub. Year	Country	N ¹	Population at risk	Prevalence ²	95%CI
Suicidality (Point)						
Castillejos	2019	Europe	1568	39605	3.96	2.37-5.56
Wishing to be dead (Point)						
Aschan	2013	UK	26	1698	1.8	0.95-2.12
Suicidal ideation (Point) [Range: 0.8-7.6%; Average : 5.28%]						
Mc Donald	2017	England	60	7393	0.80	0.60-1.00
Aschan	2013	UK	23	1698	1.35	0.80-1.90
Bebbington	2009	UK	192	8580	2.24	1.92-2.55
Casey	2008	Spain			2.3	2.00-4.80
Gisle	2013	Belgium	186	4459	4.17	3.58-4.76
Ladwig	2008	Germany	170	3154	5.39	4.60-6.18
Gili-Planas	2001	Spain	697	45	6.5	4.63-8.28
Michal	2010	Germany	374	4912	7.61	6.87-8.36
Forkman	2012	Germany	201	2509	8.01	6.95-9.07
Polatöz	2011	Turkey	104	1117	9.13	10.82-7.44
Hintikka	2009	Finland	141	1339	10.50	8.86-12.14
Economou	2013	Greece	226	2192	5.2	9.03-11.57
Economou	2013	Greece	151	2256	6.7	11.99-14.81
Attempted suicide (Point) [Range: 0.01-1.5 %; Average:0.63%]						
Mc Donald	2017	UK	1	7393	0.01	-0.01-0.04
Economou	2013	Greece	48	2192	1.1	1.59-2.81
Economou	2013	Greece	68	2256	1.5	2.30-3.70
Attempted suicide (Lifetime) [Range: 0.55-5.04%; Average : 2.88%]						
Boyd	2015	Italy	26	4712	0.55	0.34-0.77
Scocco	2008	Italy	26	4712	0.55	0.34-0.76
Boyd	2015	Bulgaria	30	5318	0.57	0.37-0.77
Boyd	2015	Romania	19	2357	0.81	0.45-1.17
Polatöz	2011	Turkey	16	1117	1.43	0.73-2.13
Miret	2014	Spain	67	4583	1.46	1.10-1.79
Boyd	2015	Spain	81	5473	1.48	1.16-1.80
Boyd	2015	Germany	61	3555	1.73	1.30-2.16
Hard	2015	ESEMED	167	8796	1.9	1.61-2.18
Ten Have	2013	The Netherlands	146	6646	2.20	1.84-2.55
Boyd	2015	The Netherlands	55	2372	2.30	1.70-2.90
Bruffaerts	2015	Belgium	26	1043	2.49	1.55-3.44
Boyd	2015	Belgium	61	2419	2.51	1.89-3.14
Boyd	2015	France	99	2894	3.43	2.76-4.09
O'Neill	2014	UK	149	4340	3.43	2.89-3.97
Saracli	2016	Turkey	31	891	3.50	2.29-4.71
Boyd	2015	UK	153	4340	3.52	2.98-4.07
Boyd	2015	Portugal	138	3849	3.59	3.00-4.17
Atay	2012	Turkey	26	600	4.33	2.70-5.96
Mc Donald	2017	UK	355	7393	4.79	4.31-5.28
Gisle	2013	Belgium	215	4459	4.82	4.19-5.45
Bebbington	2009	UK	50	8580	5.01	4.55-5.47
Encrenaz	2012	France	1116	22138	5.04	4.75-5.33

¹N: number of cases²Prevalence rate per 100 inhabitants

Table 2. Descriptive characteristics of study sample

Variables	<i>Total</i> N=4507
Gender	
Male	2214 (49.1)
Female	2293 (50.9)
Age	
18 -30	1106 (24.5)
31-45	1522 (33.8)
46-60	1135 (25.2)
61-75	744 (16.5)
Marital status	
Married/coupled	2747 (60.9)
Single/separated/divorced/widowed	1760 (39.1)
Employment status	
Employee or freelancer	1942 (43.1)
Student	316 (7.0)
Retired	504 (11.2)
Looking after the family or home	442 (9.8)
Unemployed	1221 (27.1)
Unable to work because of illness or long-term disability	81 (1.8)
Educational level	
Unfinished primary	614 (13.7)
Primary	1750 (38.8)
Secondary	1332 (29.6)
University	810 (18.0)
Town size	
Urban (+10000 inhabitants)	3593 (79.7)
Intermediate (2001-10000)	758 (16.8)
Rural (<2001)	156 (3.4)
Province	
Cordoba	430 (9.5)
Jaen	361 (8.0)
Malaga	870 (19.3)
Granada	496 (11.0)
Sevilla	1030 (22.8)
Huelva	280 (6.2)
Cadiz	665 (14.7)
Almeria	375 (8.3)

Data expressed as N (%).

Table 3. Univariate association for suicidal risk*

Variables	Total	Suicidal risk		Statistics ^a F or X ²	P value
		No	Yes		
Number of individuals, n (%)	4507 (100)	4218 (93.6)	289 (6.4)		
Gender, n (%)					
Male	2214 (49.1)	2102 (94.9)	112 (5.1)	13.285	<0.001
Female	2293 (50.9)	2116 (92.3)	177 (7.7)		
Age, Mean (SD)	42.80 (15.22)	42.67 (15.25)	44.70 (14.54)	-2.292	<0.001
Marital status, n (%)					
Married/coupled	2747 (60.9)	2613 (95.1)	134 (4.9)	27.591	<0.001
Single/separated/divorced/widowed	1760 (39.1)	1605 (92.2)	155 (8.8)		
Employment status, n (%)					
Employee or freelancer	1942 (43.1)	1857 (95.6)	85 (4.4)	47.727	<0.001
Student	316 (7.0)	300 (94.9)	16 (5.1)		
Retired	504 (11.2)	474 (94.0)	30 (6.0)		
Looking after the family or home	442 (9.8)	405 (91.6)	37 (8.4)		
Unemployed	1221 (27.1)	1116 (91.3)	106 (8.7)		
Unable to work because of illness or long-term disability	81 (1.8)	66 (81.5)	15 (18.5)		
Educational level, n (%)					
Unfinished primary	614 (13.7)	542 (88.3)	72 (8.7)	36.937	<0.001
Primary	1750 (38.8)	1652 (94.3)	99 (5.7)		
Secondary	1332 (29.6)	1250 (93.8)	82 (6.2)		
University	810 (18.0)	774 (95.6)	36 (4.4)		
Town size, n (%)					
Urban (+10000 inhabitants)	3593 (79.7)	3354 (93.3)	239 (6.7)	2.522	0.283
Intermediate (2001-10000)	758 (16.8)	714 (94.2)	44 (5.8)		
Rural (<2001)	156 (3.4)	150 (96.2)	6 (3.8)		
Province, n (%)					
Cordoba	430 (9.5)	414 (96.3)	16 (3.7)	30.846	<0.001
Jaen	361 (8.0)	345 (95.6)	16 (4.4)		
Malaga	870 (19.3)	824 (94.7)	46 (5.6)		
Granada	496 (11.0)	467 (94.2)	29 (5.8)		
Sevilla	1030 (22.8)	969 (94.1)	61 (5.9)		
Huelva	280 (6.2)	259 (92.5)	21 (7.5)		
Cadiz	665 (14.7)	607 (91.3)	58 (8.7)		
Almeria	375 (8.3)	333 (88.8)	42 (11.2)		
Personal and social performance Mean (SD)	95.43 (11.07)	96.29 (9.59)	82.89 (20.06)		
Social support, Mean (SD)	20.30 (1.86)	20.41 (1.66)	18.83 (3.40)	7.8	<0.001
Physical childhood abuse, n (%)					
No	4247 (94.2)	4015 (94.5)	232 (5.5)	110.616	<0.001
Yes	260 (5.8)	203 (71.8)	57 (21.9)		
Psychological childhood abuse, n (%)					
No	4155 (92.2)	3953 (95.1)	202 (4.9)	213.160	<0.001
Yes	352 (7.8)	265 (75.3)	87 (24.7)		
Sexual childhood abuse, n (%)					
No	4459 (98.9)	4190 (94.0)	269 (6.0)	100.482	<0.001
Yes	48 (1.1)	28 (58.3)	20 (41.7)		
Stressful life events, n (%)					
No	2110 (46.8)	2047 (97.0)	63 (3.0)	77.601	<0.001
Yes	2397 (53.2)	2171 (90.62)	226 (9.4)		
Neuroticism personal traits Mean (SD)	1.72 (1.49)	1.66 (1.48)	2.51 (1.50)	-11.746	<0.001
Impulsivity personal traits Mean (SD)	-0.98 (1.23)	-1.01 (1.21)	-0.52 (1.39)	-6.209	<0.001
Family history of mental disorder, n (%)					
No	3754 (83.3)	3571 (95.1)	183 (4.9)	88.991	<0.001

Yes	751 (16.7)	645 (85.9)	106 (14.1)		
Any suicide in the family, n (%)					
No	4470 (99.2)	4190 (93.7)	280 (6.3)	19.945	<0.001
Yes	37 (0.8)	28 (75.7)	9 (24.3)		
Drugs dependence, n (%)					
No	4408 (97.8)	4147 (94.1)	261 (5.9)	80.681	<0.001
Yes	99 (2.2)	71 (71.7)	28 (28.3)		
Alcohol consumption, Mean (SD)	0.17 (0.59)	0.16 (0.57)	0.36 (0.88)	-3.7	<0.001
Nicotine dependence n (%)					
No	4087 (90.7)	3854 (94.3)	233 (5.7)	37.191	<0.001
Yes	419 (9.3)	363 (86.60)	56 (13.4)		

^a Student's *t*-test or chi-square test (two tailed)

*Suicidality: any affirmative question of the suicidality module of MINI interview

Journal Pre-proof

Table 4. Multivariate logistic regression for suicidal risk

<i>Variables</i>	<i>OR</i>	<i>95% CI</i>	<i>P value</i>
Gender			
Male (reference)		1.04-1.82	0.02
Female	1.38		
Age			
	1.01	1.00-1.02	<0.001
Marital Status			
Married/coupled (reference)		1.22-2.09	<0.001
Single/separated/divorced/widowed	1.60		
Social support			
	0.86	0.82-0.90	<0.001
Physical childhood abuse			
No (reference)		1.66-3.56	<0.001
Yes	2.43		
Stressful life events			
No (reference)		1.30-1.63	<0.001
Yes	1.45		
Neuroticism, personal traits			
	1.34	1.25-1.43	<0.001
Family history of mental disorders			
No (reference)		1.40-2.51	<0.001
Yes	1.88		
Drugs dependence			
	2.96	1.70-5.16	<0.001
Alcohol consumption			
	1.17	0.98-1.38	0.07
Nicotine dependence			
No (reference)		1.17-2.41	<0.001
Yes	1.68		

*Suicidal risk: any affirmative question of the suicidality module of MINI interview