

REVIEW ARTICLE

Gamification and neurological motor rehabilitation in children and adolescents: a systematic review

M. Pimentel-Ponce^a, R.P. Romero-Galisteo^{b,*}, R. Palomo-Carrión^c, E. Pinero-Pinto^d, J. Antonio Merchán-Baeza^e, M. Ruiz-Muñoz^f, J. Oliver-Pecec^g, M. González-Sánchez^h

^a Ejercicio libre de la profesión, Spain

^b Departamento de Fisioterapia, Facultad de Ciencias de la Salud, Universidad de Málaga, Instituto de Investigación Biomédica de Málaga (IBIMA), Málaga, Spain

^c Departamento de Enfermería, Fisioterapia y Terapia Ocupacional, Facultad de Fisioterapia, Universidad de Castilla la Mancha, Toledo, Spain

^d Departamento de Fisioterapia, Facultad de Enfermería, Fisioterapia y Podología, Universidad de Sevilla, Sevilla, Spain

^e Grupo de investigación Methodology, Methods, Models and Outcomes of Health and Social Sciences (M30), Facultad de Ciencias de la Salud y Bienestar, Universidad de Vic-Universidad Central de Cataluña (UVIC-UCC), Vic, Barcelona, Spain

^f Departamento de Enfermería y Podología, Facultad de Ciencias de la Salud, Universidad de Málaga, Instituto de Investigación Biomédica de Málaga (IBIMA), Málaga, Spain

^g Departamento de Personalidad, Evaluación y Tratamiento Psicológico, Facultad de Psicología, Universidad de Málaga, Málaga, Spain

^h Departamento de Fisioterapia, Facultad de Ciencias de la Salud, Universidad de Málaga. Instituto de Investigación Biomédica de Málaga (IBIMA), Málaga, Spain

Received 4 September 2020; accepted 7 February 2021

KEYWORDS

Child;
Adolescent;
Neurological
rehabilitation;
Video game;
Systematic review

Abstract

Introduction: Gamification consists of the use of games in non-playful contexts. It is widely employed in the motor rehabilitation of neurological diseases, but mainly in adult patients. The objective of this review was to describe the use of gamification in the rehabilitation of children and adolescents with neuromotor impairment.

Methods: We performed a systematic review of clinical trials published to date on the MEDLINE (PubMed), Scielo, SCOPUS, Dialnet, CINAHL, and PEDro databases, following the PRISMA protocol. The methodological quality of the studies identified was assessed using the PEDro scale.

Results: From a total of 469 studies, 11 clinical trials met the inclusion criteria. We analysed the gamification systems used as part of the rehabilitation treatment of different neuromotor conditions in children and adolescents. Cerebral palsy was the most frequently studied condition (6 studies), followed by developmental coordination disorder (3), neurological gait disorders (1), and neurological impairment of balance and coordination (1).

DOI of refers to article: <https://doi.org/10.1016/j.nrl.2021.02.011>.

* Corresponding author.

E-mail address: rpromero@uma.es (R.P. Romero-Galisteo).

<https://doi.org/10.1016/j.nrleng.2023.12.006>

2173-5808/© 2023 Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Neurología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Conclusion: The use of gamification in rehabilitation is helpful in the conventional treatment of neuromotor disorders in children and adolescents, with increased motivation and therapeutic adherence being the benefits with the greatest consensus among authors. While strength, balance, functional status, and coordination also appear to improve, future research should aim to determine an optimal dosage.

© 2023 Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Neurología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Niño;
Adolescente;
Rehabilitación
neurológica;
Videojuego;
Revisión sistemática

Ludificación y neurorrehabilitación motora en niños y adolescentes: Revisión Sistemática

Resumen

Introducción: La ludificación consiste en emplear el juego en contextos no lúdicos. Su uso en la rehabilitación motora de patologías neurológicas está muy extendido, pero sobre todo en pacientes adultos. El objetivo de esta revisión fue describir el uso de la ludificación en los tratamientos de rehabilitación en niños y adolescentes con afectación neuromotora.

Métodos: Se realizó una revisión sistemática de ensayos clínicos en diferentes bases de datos: Medline (a través de Pubmed), Scielo, SCOPUS, Dialnet, Cinahl y PEDro de la literatura científica publicada hasta la fecha siguiendo el protocolo PRISMA. La calidad metodológica de los estudios identificados se evaluó a través de la escala PEDro.

Resultados: De un total de 469 estudios localizados se seleccionaron 11 ensayos clínicos que cumplieron los criterios de inclusión. Se analizaron los sistemas de ludificación utilizados como parte del tratamiento rehabilitador en distintas afecciones neuromotoras en niños y adolescentes. La Parálisis Cerebral fue la afección con mayor número de estudios (n = 6) seguida del Trastorno del Desarrollo de la Coordinación (n = 3). También se estudió la alteración de la marcha por causa neurológica (n = 1) y la alteración del equilibrio y coordinación por causa neurológica (n = 1).

Conclusión: El uso de la ludificación en rehabilitación aporta beneficios al tratamiento convencional de las alteraciones neuromotoras en niños y adolescentes siendo el incremento de la motivación y adherencia terapéutica los que mayor consenso han alcanzado entre autores. Fuerza, equilibrio, funcionalidad y coordinación son otras variables analizadas que, si bien sugieren mejoras, necesitarían futuras investigaciones para determinar una óptima dosificación.
© 2023 Publicado por Elsevier España, S.L.U. en nombre de Sociedad Española de Neurología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Neurorehabilitation is a complex healthcare process that aims to restore, minimise, or compensate to the greatest possible extent for functional deficits in individuals with severe disability secondary to central nervous system lesions.¹ In recent years, contributions from neuroscience research have led to an important paradigm shift in therapeutic approaches to neurological disease. In paediatric rehabilitation, physiotherapists working as part of interdisciplinary teams are incorporating new approaches into the treatment of motor disorders. This perspective includes task-oriented functional training, practising activities in a context relevant to the child, with a movement towards “hands-off” therapy.^{2–5} Treatment adherence and motivation are fundamental elements of this new approach, and have played a particularly important role in paediatric treatment since the publication of the International Classification of Functioning, Disability and Health for Children and Youth (ICF-CY),⁶ which describes in detail the health status of children and adolescents in the first 2 decades of life.

Gamification is defined as the use of game-related elements in a non-play setting.^{7,8} Such settings could include the workplace, education, or healthcare contexts.⁹ Most studies on gamification in

healthcare are specifically related to the rehabilitation of chronic diseases, physical activity, and mental health.^{10–12} The majority of neurorehabilitation research addresses adult patients.^{13–16}

In the paediatric field, interactive games should be used in a context in which the therapist observes and treats the child, using the game as a tool to boost motivation and training intensity in a given task. Motivation is needed to modify motor learning programmes.¹⁷ van der Kooij et al.¹⁸ showed that the inclusion of gamification techniques in the development of a given exercise made the task more pleasant, leading to better results. It should be noted that the therapeutic use of gamification differs from the use of environments enriched with toys, which are understood as complex environments that facilitate cognitive, motor, and sensory stimulation.^{19,20}

In physiotherapy, the use of computer games to attract, motivate, and engage children and adolescents does not guarantee that the desired therapeutic effect will be achieved. However, the characteristics of video games involve dopaminergic reward systems in the brain. This facilitates learning through the generation of synaptic connections.²¹ The use of games in treatment sessions also triggers cognitive patterns necessary to the treatment and recovery of numerous neurological disorders, assisting in the development of new skills or training in specific activities.²² Treatment adherence,

or engagement, and motivation therefore represent essential pillars in the rehabilitation provided by gamification.²³ This new approach helps therapists in creating challenging, attractive environments that encourage children and adolescents to practice skills.^{21,24} Gamification offers clear objectives and new challenges through the use of increasing levels of difficulty, and shows their achievements and provides feedback using a story that offers “rewards.”²⁵

The use of conventional games consoles presents several disadvantages due to the difficulty of adapting the game to the child’s abilities, as games are typically designed for users with normal motor function. Furthermore, the quantification of motor performance is inexact, which may be a hindrance for reliable, systematic follow-up.²⁶ However, said limitations should not prevent the use of these techniques, as several studies suggest that the use of physiotherapy treatments combined with gamification improves treatment success rates.^{27–29}

Identifying and disseminating knowledge on gamification among neurorehabilitation professionals will lead to new treatment opportunities. It will also facilitate the accomplishment of some treatment objectives, taking into account certain important elements alluded to in the ICF-CY, such as participation, activity, or environmental and personal factors. Among physiotherapists, in particular, clinical practice may be optimised by incorporating task-oriented activities, motivation for motor learning, encouraging engagement, etc. This study reports a systematic review of the literature published to date on the use of gamification in physiotherapy treatments in children and adolescents with motor disorders, with an analysis of the methodological quality of the studies identified.

Material and methods

We conducted a systematic review, following the PRISMA guidelines for the selection of articles.³⁰ The literature search was conducted on the PubMed, Scielo, Scopus, Dialnet, CINAHL, and PEDro databases in July 2020.

The search strategy was adapted to the different databases using the Boolean operators AND and OR. The literature search used the following MeSH terms: “games, experimental” OR “video games” OR “games, recreational” AND “neurological rehabilitation,” “physical therapy modalities,” “child*,” “infant,” “pediatrics,” and “adolescent.” We also added the terms “gamification,” “serious game,” and “game base.”

Inclusion and exclusion criteria

Inclusion criteria were: clinical trials using some physiotherapy intervention to treat patients aged 0–18 years with some neuro-motor disorder, published in English or Spanish.

Results were not filtered by date of publication, and we only excluded articles that did not report results related to the physiotherapy intervention.

Selection of articles

Two reviewers (MP and MG) independently searched the different databases, eliminating duplicates and verifying that articles met the selection criteria. Both reviews were submitted to a third reviewer (RR), who resolved any discrepancies. A total of 11 studies were finally included for analysis (Fig. 1).

Assessment of methodological quality

The PEDro scale was used to determine the methodological quality of the articles selected.³¹ This scale measures the internal validity and interpretability of results of clinical trials using 11 items, scored 1 or 0 depending on whether each is present in the study. The max-

imum possible score is 10 points, as the first item is not counted for the final score. Methodological quality is considered high for studies scoring 6 points or more, moderate for scores of 4–5, and low for studies scoring below 4.³²

The procedure for the assessment of studies meeting the eligibility criteria mentioned above was the same as that used to select the final results: 2 reviewers (MP and MG) independently evaluated articles, and a third reviewer (RR) resolved any discrepancies.

Results

The initial literature search yielded 469 articles; 404 remained after elimination of 65 duplicate results. Subsequently, results were screened by applying the search terms to the titles and abstracts only; 47 remained after screening and were read in full text to apply the eligibility criteria. A total of 10 studies were finally included for qualitative analysis (Fig. 1).

In the qualitative assessment of studies, PEDro scores ranged from 2 points³⁶ to 8 points.^{33,40} Seven studies scored 6 points or higher, and were therefore considered to be of high quality.^{33–35,37–39} Only the study by Hammond et al.⁴⁰ was classed as being of moderate quality (5/10 points). The 2 remaining studies were of low quality, meeting only one³⁶ and 3 criteria,⁴¹ respectively.

It should be noted that 2 items were not present in any of the studies analysed: blinding of patients and blinding of therapists. On the other hand, one item was fulfilled by all studies: comparison between groups.

Table 1 presents the structural characteristics of the studies reviewed. The studies reviewed included a total of 344 participants, although sample sizes varied greatly, ranging from 6 participants³⁶ in one study to 62 in another.³³ Participants’ ages ranged from 5 years^{35,37,38} to 17 years.^{35,39}

Cerebral palsy (CP) was the neurological disorder addressed in the greatest number of studies analysing gamification (n=6).^{33–38} Secondly, developmental coordination disorder (DCD) was explicitly addressed in 3 studies.^{39,40} Only one study analysed gamification in children and adolescents with probable DCD and balance problems.⁴¹

All intervention groups completed treatment programmes that included the use of new technologies. The duration of the intervention ranged from 14 sessions³⁹ to 60 sessions.³⁷ Participants completed between one³⁹ and 6 sessions³⁵ per week. Finally, the duration of interventions ranged from 3 to 14 weeks.³⁹ Table 1 presents in greater detail the remaining structural characteristics of the reviewed studies.

Table 2 shows the results of all the studies reviewed, classifying measurements into 3 time periods: short term (< 6 weeks), medium term (6–12 weeks), and long term (> 12 weeks). A total of 21 outcome variables were analysed, although not all were measured over all 3 time periods. Specifically, 8 variables were studied over the short term, 11 over the medium term, and 9 over the long term.

Variables can be classified into 3 different categories: quantitatively analysed functional capacities (9 variables); qualitatively analysed functional capacities (9); and subjective assessment using questionnaires (4). Some variables (e.g., motor performance) were analysed with more than one instrument, and therefore may appear in more than one of these categories. Table 2 presents detailed information on the results for each variable.

Ferguson et al.⁴³ compared the effects of 2 task-oriented intervention strategies in children aged 6–10 years with DCD. One strategy used the Nintendo Wii console. The variables measured included motor performance, aerobic/anaerobic capacity, and muscle strength. Both groups presented significant improvements, with anaerobic capacity being the only one for which gamification was associated with statistically significant differences compared to neuromotor task training based on cognitive neuroscience and

Table 1 Main characteristics of the final sample of articles included in the study.

Authors, year	Sample size (no. girls)	Age	Disorder	Interventions	Sessions	Outcome variables	Time of measurement
Arnoni et al., ³⁸ 2019	15	5–13	CP	IG: conventional therapy + virtual reality (20 + 20 min/session) CG: conventional therapy (50 min/session stretching and strength exercises)	2 sessions/week (8 weeks)	Balance	Pre-/post-intervention
	IG: 7 (1)					GMFM	
Bonney et al., ³⁹ 2017	43	13–16	DCD	Nintendo Wii with Wii Balance Board and the Wii Fit game Task-oriented functional training	1 session/week (14 weeks)	-Isometric muscle strength	Pre-/post-intervention
	Nintendo Wii: 21 NTT: 22					-Motor coordination (MABC-2) -Self-efficacy in activities of daily living -Coordination (tracking task)	
Chiu et al., ³³ 2014	62	6–13	CP	IG: conventional therapy + training at home with Wii Sport Resort (10 min/game; 4 games = 40 min.)	IG: 3 days/week (6 weeks)	-Strength (dynamometer) -Hand function (Nine-Hole Peg Test and Jebsen-Taylor Hand Function Test) -Caregiver perception of hand function (Functional Use Survey)	Pre-intervention and at 6/12 weeks
	IG: 32 (17)						
Ferguson et al., ⁴³ 2013	56	6–10	DCD	CG: conventional therapy Training with Wii Fit, using the Wii Balance Board and 18 games (30 min/session) Task oriented training (NTT); 45–60 min/session	3 sessions/week (6 weeks)	-Motor coordination (MABC-2)	Pre-/post-intervention
	Wii: 19					-Hand strength (FSM)	
	NTT: 37					-Grip strength (dynamometer)	

Table 1 (Continued)

Authors, year	Sample size (no. girls)	Age	Disorder	Interventions	Sessions	Outcome variables	Time of measurement
Gatica-Rojas et al., ³⁴ 2017	32 IG: 16 (6) CG: 16 (7)	7–14	CP	IG: Wii Fit Plus game with Nintendo Balance Board. 2 series of one game and 1 series of Yoga; 30 min/session CG: conventional physiotherapy; 40 min/session	3 sessions/week (6 weeks)	-Muscle strength (MPST) -Aerobic capacity (20mSRT) -Standing balance (posturography)	Pre-intervention; 2, 4, and 6 weeks. Follow-up at weeks 8 and 10
Hammond et al., ⁴⁰ 2013	18 IG: 10 (2) CG: 8 (2)	7–10	DCD	IG: training with Wii Fit, using the Wii Balance Board and 9 games (phase 1: 10 min/day) CG: conventional therapy (1 h/week)	Phase 1: 3 days/week (4 weeks) Phase 2 (2.5 months after phase 1): 3 days/week (4 weeks)	-Motor proficiency (short-form BOT-2): fine manual control, coordination, strength, and agility -Self-perceived ability and satisfaction with motor tasks (CSQ) -Emotional and behavioural development (SDQ) for parents	Pre-intervention, between phases, after phase 2
Hsieh, ³⁷ 2018	40 IG: 20 CG: 20	5–10	CP	IG: computer game + standing platform (40 min/session) CG: computer game + mouse (40 min/session)	5 days/week (12 weeks)	-Postural balance (centre-of-pressure sway) -Functional balance (BBS) -Standing and dynamic balance (FAB)	Pre-/post-intervention
Jelsma et al., ⁴¹ 2014	34	6–12	BD	Playing 2 Wii Fit Plus games, selected from 18 options	3 days/week (6 weeks)	-Functional mobility and dynamic balance (TUG) -Motor coordination (MABC-2)	Pre-/post-intervention

Table 1 (Continued)

Authors, year	Sample size (no. girls)	Age	Disorder	Interventions	Sessions	Outcome variables	Time of measurement
Kassee et al., ³⁶ 2017	IG: 14	7–12	CP	Wii Sports Resort game (40 min/day; 5 days/week) 6 upper-limb resistance exercises (12 repetitions each)	5 days/week (6 weeks)	-Motor proficiency (BOT-2): fine manual control, coordination, running speed and agility	Pre-/post-intervention
	CG: 20					-Standing balance (Wii Fit Ski Slalom test)	
	Wii: 3					-Level of enjoyment (Enjoyment Scale)	
Sajan et al., ³⁵ 2016	Resistance: 3	5–16	CP	Nintendo Wii game (boxing + tennis) (20 min/session) + conventional multidisciplinary therapy (25 min/session) Conventional multidisciplinary therapy	6 days/week (3 weeks)	-Upper limb function (MA2; ABILHAND-Kids)	Pre-/post-intervention
	IG: 9					-Grip strength (dynamometer)	
	CG: 9					-Motivation and feasibility (parent-reported questionnaire)	
						-Posture control (static posturography)	
						-Balance (paediatric BBS)	
						-Manual dexterity (Box and Block Test)	
						-Quality of upper limb use (QUEST)	
						-Visual perception (TVPS-3)	
						-Functional ambulation (walking distance and speed)	

20mSRT: 20-m Shuttle Run Test; BBS: Berg Balance Scale; BD: balance disorder; BOT-2: Bruininks-Oseretsky Test-2; CG: control group; CP: cerebral palsy; CSQ: Coordination Skills Questionnaire; DCD: developmental coordination disorder; FAB: Fullerton Advanced Balance scale; FSM: Functional Strength Measure; GMFM: Gross Motor Function Measure; IG: intervention group; MA2: Melbourne Assessment of Unilateral Upper Limb Function-2; MABC-2: Movement Assessment Battery for Children-2; MPST: Muscle Power Sprint Test; NTT: Neuromotor Task Training; QUEST: Quality of Upper Extremity Skills Test; SDQ: Strengths and Difficulties Questionnaire; TUG: Timed Up and Go; TVPS-3: Test for Visual-Perceptual Skills-3rd edition.

Table 2 Short-, medium-, and long-term results for the outcome variables analysed in the different studies.

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)
Quality of upper limb movement	Kassee et al., ³⁶ 2017		MA2	Intervention group	77.94	–	87.27	82.59
				Control group	82.02	–	85.77	84.64
			ABILHAND	Intervention group	1.84	–	2.93	0.87
				Control group	1.50	–	1.38	0.11
				Intervention group	2.00	–	3.08	3.42
				Control group	2.83	–	2.25	2.50
Posture control	Sajan et al., ³⁵ 2016	Static posturography	Sway velocity-eyes open (mm/s)	Intervention group	137.67	83.06	–	–
				Control group	137.66	127.93	–	–
			Sway velocity-eyes closed (mm/s)	Intervention group	131.29	86.92	–	–
				Control group	120.33	123.93	–	–
			Paediatric BBS	Intervention group	15.70	18.7	–	–
				Control group	20.44	25.00	–	–
Compliance, motivation, and feasibility	Kassee et al., ³⁶ 2017	Compliance and parent feedback questionnaire	Q1	Intervention group	–	–	4.00	–
				Control group	–	–	2.00	–
			Q2	Intervention group	–	–	3.00	–
				Control group	–	–	.67	–
			Q3	Intervention group	–	–	2.33	–
				Control group	–	–	.67	–
Q4	Intervention group	–	–	5.00	–			
	Control group	–	–	4.67	–			
Ambulation	Sajan et al., ³⁵ 2016		Walking speed (m/min)	Intervention group	12.61	34.31	–	–
				Control group	23.89	24.61	–	–
			Walking endurance (distance in metres)	Intervention group	131.10	317.22	–	–
				Control group	151.00	218.44	–	–
Gross manual dexterity	Sajan et al., ³⁵ 2016		Box and Block Test	Intervention group	46.90	68.00	–	–
				Control group	55.2	59.33	–	–
			Total CoP displacement (cm)	Intervention group	–	–15.93	–	–
				Control group	–	–14.53	–	–
			AP CoP displacement (cm)	Intervention group	–	1.45	–	–
				Control group	–	–2.22	–	–
ML CoP displacement (cm)	Intervention group	–	2.45	–	–			
	Control group	–	–2.44	–	–			
Balance	Arnoni et al., ³⁸ 2019	Force plate	AP amplitude of CoP displacement (cm)	Intervention group	–	–3.66	–	–
				Control group	–	2.03	–	–
			ML amplitude of CoP displacement (cm)	Intervention group	–	6.23	–	–
				Control group	–	–4.85	–	–

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)
Balance	Gatica-Rojas et al., ³⁴ 2017	AMTI OR6–7 force plate	Area (cm ²)	Intervention group	–	23.32	–	–
				Control group	–	13.29	–	–
			Mean velocity (cm/s)	Intervention group	–	20.22	–	–
				Control group	–	46.84	–	–
			CoP sway, eyes open (cm ²)	Intervention group	3.75	3.58	3.06	2.19
				Control group	3.92	4.08	4.84	4.68
			CoP sway, eyes closed (cm ²)	Intervention group	4.17	4.73	4.85	4.83
				Control group	6.64	6.26	5.36	3.69
			SD _{ML} , eyes open (cm)	Intervention group	0.43	0.39	0.42	0.36
				Control group	0.46	0.42	0.46	0.44
			SD _{ML} , eyes closed (cm)	Intervention group	0.45	0.42	0.48	0.45
				Control group	0.51	0.53	0.50	0.43
			SD _{AP} , eyes open (cm)	Intervention group	0.43	0.61	0.47	0.47
				Control group	0.48	0.55	0.60	0.64
			SD _{AP} , eyes closed (cm)	Intervention group	0.45	0.70	0.64	0.60
				Control group	0.51	0.63	0.60	0.60
			ML velocity, eyes open (cm/s)	Intervention group	0.85	1.00	0.90	0.84
				Control group	0.97	0.91	0.94	0.88
			ML velocity, eyes closed (cm/s)	Intervention group	0.92	1.15	1.07	0.96
				Control group	1.05	1.03	0.96	0.91
			AP velocity, eyes open (cm/s)	Intervention group	0.94	1.00	0.95	1.00
				Control group	1.10	0.97	1.02	1.06
			AP velocity, eyes closed (cm/s)	Intervention group	1.11	1.18	1.13	1.22
				Control group	1.32	1.14	1.05	1.15
	AP sway (mm)	Intervention group		9.94 (2.27)	9.17 (1.69)	–	–	
		Control group		10.87 (1.41)	10.55 (1.29)	–	–	
	ML sway (mm)	Intervention group		6.62 (0.69)	6.35 (1.00)	–	–	
		Control group		7.13 (1.08)	6.72 (0.70)	–	–	
	Sway area (cm ²)	Intervention group		14.42 (2.35)	13.71 (2.24)	–	–	
		Control group		14.33 (2.39)	13.93 (2.12)	–	–	
Sway velocity (cm/s)	Intervention group		4.04 (0.45)	3.56 (0.47)	–	–		
	Control group		3.69 (0.37)	3.67 (0.23)	–	–		
BBS (score)	Intervention group		44.74 (2.75)	48.81 (4.74)	–	–		
	Control group		44.39 (2.33)	45.37 (2.68)	–	–		
FAB (score)	Intervention group		21.32 (1.47)	23.41 (2.09)	–	–		
	Control group		22.07 (2.23)	22.25 (1.90)	–	–		
	Hsieh, ³⁷ 2018	CoP kinematics						
	Hsieh, ³⁷ 2018	Specific tests						

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)
Strength	Chiu et al., ³³ 2014	PowerTrack II™	TUG (s)	Intervention group	16.43 (2.12)	17.51 (1.70)	–	–
				Control group	15.60 (1.10)	15.91 (1.87)	–	–
			Pressure	Intervention group	23.5 (21.5)	–	28.3 (22.3)	29.7 (23.3)
				Control group	26.0 (17.8)	–	26.9 (23.3)	29.4 (21.4)
			Total FSM	Intervention group	10.59 (4.50)	–	6.24 (2.72)	–
				Control group	6.87 (3.58)	–	6.95 (3.95)	–
			Over-hand throwing	Intervention group	2.16 (0.73)	–	2.04 (0.55)	–
				Control group	1.98 (0.45)	–	2.09 (0.53)	–
			Long-jump	Intervention group	1.09 (0.31)	–	1.09 (0.18)	–
				Control group	1.03 (0.25)	–	1.05 (0.36)	–
Functional strength	Ferguson et al., ⁴³ 2013	FSM	Under-hand throwing	Intervention group	2.72 (0.77)	–	2.80 (0.60)	–
				Control group	2.50 (0.50)	–	2.70 (0.60)	–
			Lateral step up right	Intervention group	27.85 (6.47)	–	34.48 (5.49)	–
				Control group	32.47 (4.97)	–	33.00 (4.24)	–
			Lateral step up left	Intervention group	27.48 (6.47)	–	35.19 (5.76)	–

Table 2 (Continued)								
Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term (< 6 weeks) Mean (SD)	Medium term (6–12 weeks) Mean (SD)	Long term (> 12 weeks) Mean (SD)
Isometric strength	Ferguson et al., ⁴³ 2013	Hand-held dynamometer	Chest pass	Control group	33.05 (5.06)	–	32.42 (4.87)	–
				Intervention group	1.69 (0.52)	–	1.83 (0.38)	–
			Sit to stand	Control group	1.69 (0.25)	–	1.67 (0.29)	–
				Intervention group	20.30 (5.14)	–	25.56 (4.66)	–
			Lift box	Control group	25.95 (6.72)	–	26.58 (5.50)	–
				Intervention group	16.52 (6.56)	–	22.15 (6.88)	–
			Stairs	Control group	18.79 (3.63)	–	20.89 (4.37)	–
				Intervention group	61.04 (13.11)	–	69.46 (11.81)	–
			Right elbow flexors	Control group	63.84 (9.10)	–	64.79 (7.56)	–
				Intervention group	104.62 (25.06)	–	109.21 (33.24)	–
			Left elbow flexors	Control group	97.48 (20.11)	–	90.53 (21.13)	–
				Intervention group	101.71 (26.91)	–	105.43 (28.48)	–

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)
				Control group	92.50 (24.28)	–	89.69 (19.79)	–
			Right elbow extensors	Intervention group	84.47 (21.51)	–	80.96 (19.44)	–
				Control group	70.54 (20.59)	–	76.32 (17.36)	–
			Left elbow extensors	Intervention group	86.72 (21.16)	–	81.41 (21.10)	–
				Control group	71.77 (20.83)	–	79.58 (27.94)	–
			Right knee extensors	Intervention group	152.25 (51.17)	–	150.65 (48.75)	–
				Control group	127.22 (32.72)	–	139.37 (34.67)	–
			Left knee extensors	Intervention group	151.71 (51.46)	–	145.54 (51.55)	–

Table 2 (Continued)								
Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term (< 6 weeks) Mean (SD)	Medium term (6–12 weeks) Mean (SD)	Long term (> 12 weeks) Mean (SD)
Muscle strength	Bonney et al., ³⁹ 2017		Right grip force	Control group	120.43 (35.35)	–	137.57 (31.71)	–
				Intervention group	43.33 (12.37)	–	47.03 (13.32)	–
			Left grip force	Control group	35.56 (7.57)	–	49.42 (12.92)	–
				Intervention group	37.83 (10.21)	–	43.58 (12.64)	–
			Knee extensors (N)	Control group	35.03 (8.64)	–	42.94 (8.83)	
				Intervention group	139.5 (27.1)	–		263.6 (49.2)
			Ankle	Control group	157.9 (22.9)	–		302.3 (68.1)
				Intervention group	98.9 (17.7)	–		281.0 (34.5)
			Ankle plantar flexors (N)	Control group	109.4 (20.0)	–		271.0 (51.3)
				Intervention group	119.3 (17.8)	–		213.0 (34.9)
Hand function	Chiu et al., ³³ 2014		Nine-Hole Peg Test	Control group	128.7 (12.9)	–		229.4 (38.9)
				Intervention group	0.10 (0.09)	–	0.10 (0.19)	0.11 (0.11)
			Jebsen-Taylor Hand Function Test	Control group	0.12 (0.10)	–	0.13 (0.11)	0.13 (0.11)
				Intervention group	0.21 (0.14)	–	0.26 (0.18)	0.31 (0.19)
				Control group	0.22 (0.15)	–	0.27 (0.20)	0.34 (0.21)

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)
Upper limb function	Sajan et al., ³⁵ 2016	QUEST	QUEST: grasp domain	Intervention group	67.82	73.35	–	–
				Control group	81.05	83.95	–	–
			QUEST: dissociated movements domain	Intervention group	77.94	80.40	–	–
				Control group	92.61	92.96	–	–
			QUEST: total score	Intervention group	72.86	76.38	–	–
				Control group	86.83	88.45	–	–
Generalised self-efficacy	Bonney et al., ³⁹ 2017		CSAPPA total score	Intervention group	52.7 (10.7)	–	–	55.6 (11.2)
				Control group	50.0 (12.8)	–	–	53.9 (9.9)
			Adequacy score	Intervention group	19.8 (4.5)	–	–	19.9 (4.9)
				Control group	17.6 (5.4)	–	–	19.6 (5.4)
			Enjoyment score	Intervention group	9.8 (2.5)	–	–	9.5 (2.2)
				Control group	9.7 (2.5)	–	–	9.4 (1.6)
GMFM	Arnoni et al., ³⁸ 2019	GMFM	Predilection score	Intervention group	23.2 (6.5)	–	–	26.1 (5.9)
				Control group	22.8 (7.3)	–	–	24.9 (4.3)
			Standing	Intervention group	–	–2.310	–	–
				Control group	–	NA	–	–

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term	
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)	
Visual-perceptual skills	Sajan et al., ³⁵ 2016	TVPS-3	Walking, running, and jumping	Intervention group	–	–2.672	–	–	
				Control group	–	NA	–	–	
			TVPS (total score)	Intervention group	32.10	43.44	–	–	
				Control group	36.50	37.30	–	–	
MABC-2 (motor proficiency)	Bonney et al., ³⁹ 2017		Total standard score	Intervention group	61.4 (10.4)	–	–	74.0 (10.3)	
				Control group	63.0 (5.6)	–	–	78.1 (8.8)	
			Manual dexterity	Intervention group	5.5 (2.1)	–	–	8.9 (2.7)	
				Control group	5.9 (1.5)	–	–	9.7 (1.5)	
			Aiming and catching	Intervention group	9.0 (2.3)	–	–	8.9 (2.5)	
				Control group	9.3 (2.6)	–	–	10.1 (2.9)	
	Ferguson et al., ⁴³ 2013			Balance	Intervention group	7.0 (2.9)	–	–	8.9 (2.6)
					Control group	6.9 (2.1)	–	–	9.1 (2.5)
				Total standard score	Intervention group	4.26 (1.02)	–	8.67 (2.51)	–
					Control group	5.32 (1.45)	–	6.05 (2.83)	–
			Manual dexterity	Intervention group	6.04 (2.23)	–	9.04 (2.56)	–	
				Control group	6.05 (2.42)	–	6.16 (3.02)	–	

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term		
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)		
Caregiver perception	Chiu et al., ³³ 2014	Functional Use Survey	Aiming and catching	Intervention group	7.56 (2.72)	–	8.52 (2.44)	–		
				Control group	6.53 (3.27)	–	6.84 (3.18)	–		
			Balance	Intervention group	4.26 (2.44)	–	9.48 (2.31)	–		
				Control group	7.00 (2.31)	–	8.11 (2.47)	–		
			Quantity (0–65)	Intervention group	28.1 (17.9)	–	32.7 (15.9)	35.6 (15.8)		
				Control group	34.0 (16.9)	–	32.6 (13.4)	33.8 (15.5)		
			Quality (0–65)	Intervention group	30.1 (17.6)	–	34.0 (16.7)	34.8 (13.4)		
				Control group	34.7 (14.9)	–	34.5 (14.9)	35.3 (14.5)		
			Aerobic and anaerobic performance	Ferguson et al., ⁴³ 2013	MPST	Intervention group	104.92 (45.91)	–	139.70 (63.63)	–
						Control group	115.88 (36.91)	–	136.54 (41.64)	–
Intervention group	1.28 (0.40)	–				1.74 (1.05)	–			
Control group	1.63 (0.86)	–				1.61 (0.92)	–			
BOT-2 Running and agility (SS)	Intervention group	14.1 (3.9)				–	–	18.5 (3.2)		
Control group	14.4 (2.9)	–				–	18.3 (1.6)			
Functional performance	Bonney et al., ³⁹ 2017	10 × 5 m straight sprint (s)	Intervention group	24.8 (3.9)	–	–	22.2 (3.0)			
			Control group	25.0 (3.5)	–	–	21.7 (2.1)			
			Intervention group	24.3 (4.1)	–	–	22.2 (3.1)			
			Control group	24.6 (3.8)	–	–	20.6 (2.0)			
			Intervention group	67.5 (14.9)	–	–	74.8 (10.4)			
			Control group	–	–	–	–			

Table 2 (Continued)

Outcome variable	Study	Measuring instrument	Measurement variable	Group	Pre-intervention	Short term	Medium term	Long term
						(< 6 weeks) Mean (SD)	(6–12 weeks) Mean (SD)	(> 12 weeks) Mean (SD)
Motor proficiency	Hammond et al., ⁴⁰ 2013	BOT-2	Fine motor precision	Control group	72.0 (7.3)	–	–	78.0 (7.9)
				Intervention group	9.2 (1.95)	10.3 (2.39)	10.2 (2.92)	–
			Fine motor integration	Control group	10.4 (3.26)	10.0 (2.51)	12.25 (2.38)	–
				Intervention group	7.5 (2.17)	8.9 (2.45)	7.6 (2.06)	–
			Manual dexterity	Control group	9.0 (0.82)	8.25 (1.83)	9.25 (1.16)	–
				Intervention group	4.3 (1.23)	5.3 (1.04)	5.3 (1.39)	–
			Bilateral coordination	Control group	4.14 (1.07)	4.5 (1.31)	5.5 (0.76)	–
				Intervention group	4.6 (1.83)	6.4 (0.91)	5.8 (1.46)	–
			Balance	Control group	6.29 (0.76)	5.5 (0.93)	7.38 (1.06)	–
				Intervention group	5.9 (2.84)	7.8 (2.91)	7.5 (3.28)	–
			Running speed and agility	Control group	7.57 (0.79)	6.88 (1.46)	8 (0)	–
				Intervention group	1.2 (4.12)	2.3 (3.86)	1.3 (3.88)	–
			Upper limb coordination	Control group	1.57 (0.53)	1.5 (0.53)	1.75 (0.46)	–
				Intervention group	7.8 (2.99)	8.9 (2.43)	8.2 (2.27)	–
			Strength	Control group	6.71 (1.5)	9.13 (1.89)	10 (2)	–
Intervention group	5.7 (2.31)	6.8 (1.93)		7.2 (2.35)	–			
Aerobic endurance	Bonney et al., ³⁹ 2017	Shuttle run (level)	Intervention group	5.29 (1.5)	7.63 (4.31)	8.88 (3.09)	–	
			Control group	2.3 (1.5)	–	–	2.4 (1.6)	
				Control group	1.9 (0.9)	–	–	2.3 (1.1)

20mSRT: 20-m Shuttle Run Test; AP: anterior-posterior; BBS: Berg Balance Scale; BOT-2: Bruininks-Oseretsky Test-2; CoP: centre of pressure; CSAPPA: Children's Self-Perceptions of Adequacy in and Predilection for Physical Activity scale; FAB: Fullerton Advanced Balance scale; FSM: Functional Strength Measure; GMFM: Gross Motor Function Measure; MA2: Melbourne Assessment of Unilateral Upper Limb Function-2; MABC-2: Movement Assessment Battery for Children-2; ML: medial-lateral; MPST: Muscle Power Sprint Test; NA: not available; QUEST: Quality of Upper Extremity Skills Test; SD_{AP}: standard deviation of AP centre of pressure; SD_{ML}: standard deviation of ML centre of pressure; SS: scale scores; TUG: Timed Up and Go; TVPS-3: Test for Visual-Perceptual Skills-3rd edition.




Figure 1 Article selection process.

motor learning theories. Outcome variables were measured with the most recent version of the Movement Assessment Battery for Children-2 (MABC-2), the 20-m Shuttle Run Test, the Muscle Power Sprint Test, and hand-held dynamometer measurement of grip strength.

Hammond et al.⁴⁰ used the Nintendo Wii Fit game, on the Nintendo Wii console, to compare efficacy against the conventional therapy. They selected 9 balance- and coordination-based games, which members of the intervention group played for 10 minutes, 3 times weekly. In the second phase, both groups switched interventions for 4 weeks. The outcome variables studied were motor skills, self-perceived ability and satisfaction with the motor task, and emotional and behavioural development. Regarding motor skills, both groups presented improvements in the short-form Bruininks-Oseretsky Test (BOT-2), a measure of manual dexterity. However, the change was only statistically significant in the intervention group. Similarly, both groups showed improvements in self-perceived ability and satisfaction with the motor task at the end of phase 1, and the improvement persisted after phase 2; however, these results were not statistically significant. These variables were measured with the Coordination Skills Questionnaire. Emotional and behavioural development was analysed with the parent-reported Strengths and Difficulties Questionnaire; only 7 families (for a total of 18 participating patients) responded, with the intervention group showing a quantifiable improvement.

Chiu et al.³³ used the Nintendo Wii console with the Wii Sports Resort game in a sample of 62 children with hemiplegic CP, comparing this intervention against conventional physiotherapy in the paretic upper limb. The authors analysed coordination and hand strength and function, as well as caregivers' perception of hand function. These variables were measured before the intervention, at the end of the intervention, and at 6 weeks post-intervention. The games used (bowling, air sports, Frisbee, and basketball) were selected because of the ability to select progressive difficulty levels. Improvements were observed in the intervention group, but were not statistically significant; therefore, the study does not demonstrate that the use of video games offers advantages over conventional therapy in such variables as coordination, strength, or hand function (measured with the Nine-Hole Peg Test, Jebsen-Taylor Hand Function Test, and dynamometry measurement) in children with hemiplegia. However, perceived hand function, measured with the parent-reported Functional Use Survey, did show improvements in the intervention group.

Jelsma et al.⁴¹ studied the effect on balance of training with the Nintendo Wii, with the Wii Fit Plus game and the Wii Balance Board. The sample was made up of children with balance problems and typically developing children. Although the study used the MABC-2 to assess balance, the authors did not assert that children scoring below percentile 16 presented DCD. All participants used these platforms for sessions of 30 minutes, 3 times weekly. During that time, they could choose from 18 games available in Wii Fit Plus, with the exception of the Ski Slalom game, which was used for assessment. Participants had to play each game twice before they could choose a different one. Before and after the intervention, participants were assessed to test motor ability (MABC-2), motor proficiency (BOT-2), standing balance (Wii Fit Ski Slalom), and enjoyment (Enjoyment Scale). Analysis of the results showed that the typically developing children scored better for the balance test. Children with balance problems also showed a significant improvement after the intervention, with positive results persisting after the non-intervention period. All children enjoyed the intervention, and continued to show motivation at 6 weeks post-intervention.

Gatica-Rojas et al.³⁴ analysed standing balance in children with CP, comparing the results after a conventional physiotherapy intervention (stretching, strengthening, flexibility, and balance) against use of Wii Fit Plus with the Nintendo Wii Balance Board. Participants completed 2 sets on the Snowboard, Penguin Slide, and Super Hula Hoop games. The intervention group showed statistically significant

improvements in all variables, with the effect persisting 6 weeks after completion of the intervention. However, the improvement in centre-of-pressure sway did not persist in the long term. Outcome variables were measured using posturography.

The study by Sajan et al.³⁵ measured posture control, manual dexterity, upper limb function, and ambulation in children and adolescents with CP. Both the control group and the intervention group, who completed a game-based intervention with the Nintendo Wii system, showed similar results in the different study variables. Only manual dexterity as measured with the Quality of Upper Extremity Skills Test showed significant improvements in the intervention group after treatment. The variables postural control, balance, unilateral gross manual dexterity, visual-perceptual skills, and functional ambulation (studied with static posturography, the Berg Balance Scale, the Box and Block Test, the Test for Visual-Perceptual Skills, and walking speed and distance, respectively) showed no significant difference after adding conventional physiotherapy to the intervention.

Bonney et al.³⁹ compared the efficacy of Nintendo Wii training against task-oriented functional training in 43 girls aged 13–16 years with DCD. They measured strength in the knee flexors, ankle plantar flexors, and dorsiflexors, coordination, and self-efficacy, among other variables. No differences between the 2 interventions were observed in the different outcome measures used, including the MABC-2, BOT-2, and Participation in Activities of Daily Living for Adolescents Questionnaire.

Kassee et al.³⁶ compared upper limb functional ability and strength and treatment adherence in 6 children with CP. The intervention consisted in Nintendo Wii training, and was compared against resistance training with a series of 6 exercises to be performed each week. Both interventions were followed at home. This study found significant differences between groups, with greater improvements in the intervention group for all variables except grip strength, for which dynamometry showed better results in the conventional training group. Motivation and feasibility of home training were studied with a questionnaire completed by participants' parents.

In a sample of 40 children with CP, Hsieh³⁷ studied 2 interventions to treat balance alterations, measuring centre-of-pressure sway, the Berg Balance Scale, the Fullerton Advanced Balance Scale, and the Timed Up and Go test. Results were compared between the intervention group, who used a platform through which children used their whole body to play computer games, and the control group, who played the same games with a computer mouse rather than the standing platform. The author concluded that the platform could be used to improve balance in children with CP.

Finally, Arnoni et al.³⁸ compared the effect on balance and gross motor function of conventional therapy and a virtual reality game in 15 children and adolescents with CP. Outcome variables were measured with a force platform and with the Gross Motor Function Measure tool. The intervention group showed statistically significant improvements in gross motor function but not in standing stability.

Discussion

The aim of this study was to review the scientific literature on the use of gamification in the neurorehabilitation of children. After gathering and critically analysing clinical trials of children with motor disorders, we observed that gamification is a treatment strategy that should be considered. Short-, medium-, and long-term improvements were observed in motivation, balance, strength, function, coordination, and family satisfaction, among other variables, when gamification was incorporated into the treatment of children and adolescents with neuromotor disorders. Our conclusions are supported by the high methodological quality of the majority of the studies analysed.

Game mechanics are essential, and lead to engagement and motivation in a field that was not originally related to the game, such as neurorehabilitation. However, several studies included motivation as an outcome variable.^{36,40–42} This is highly relevant, as treatment adherence is often a challenge in the treatment of children and adolescents.

A systematic review by Sardi et al.,¹⁰ addressing the use of gamification in healthcare contexts, analysed 46 articles mainly including patients with chronic diseases and mental illness. Video games have been incorporated into the neurorehabilitation treatment of adults by authors including Yates et al.,¹³ Psychouli et al.,¹⁴ Tseklevs et al.,¹⁵ and Trombetta et al.¹⁶

It is essential to consider both personal and environmental factors in the development of treatments, in accordance with the recommendations of the ICF-CY.⁶ Thus, some researchers apply interventions in the environments in which children and adolescents naturally develop, such as the home^{33,36} or at school.⁴⁰ Thus, promotion and monitoring of community participation³⁹ and involvement of parents in treatment^{33,36,40} are approaches that follow these recently developed principles, which are not always present in intervention proposals.

The use of conventional games consoles presents several challenges, such as the difficulty of adapting the game to the child's abilities, as games are usually designed for users without motor alterations. Furthermore, the quantification of motor performance is inexact, which may constitute a hindrance to reliable, systematic follow-up.²⁶ However, authors including Ferguson et al.,⁴³ Hammond et al.,⁴⁰ Chiu et al.,³³ Jelsma et al.,⁴¹ Gatica-Rojas et al.,³⁴ Sajjan et al.,³⁵ Bonney et al.,³⁹ and Kassee et al.³⁶ did not hesitate to use commercially available games and platforms; other authors designed video games for specific purposes.⁴² Nonetheless, these limitations should not prevent the use of these games, as several studies suggest that the use of physiotherapy treatments combined with gamification improves treatment success rates.^{27–29}

Short-term effects

The short-term results of interventions in patients with CP show improvements in the vast majority of the variables analysed. Thus, intervention groups participating in rehabilitation programmes including gamification strategies showed short-term improvements in such areas as posture control,³⁵ ambulation,³⁵ gross manual dexterity,³⁵ and visual perceptual skills,³⁵ compared to controls. However, variables including upper limb function³⁵ and balance^{34,37} showed similar values after the intervention, although patients with CP did show short-term improvements compared to controls in one balance parameter (centre-of-pressure sway, open-eyes).³⁴

Patients with DCD displayed significant short-term improvements in motor performance.⁴⁰ The results reported support the need to develop specific treatment protocols for the outcome variable of interest, as very good short-term benefits have been observed in patients participating in 3 or 6 sessions per week,^{35,40} whereas patients undergoing programmes with 3 or 5 sessions per week showed similar outcomes to controls.^{34,37}

Medium-term effects

As was the case with short-term effects, the majority of variables analysed in patients with CP showed greater medium-term benefits in intervention groups than in controls (Table 2). Specifically, intervention programmes including gamification showed greater compliance, motivation, and feasibility,³⁶ with participating patients showing better upper limb quality of movement³⁶ and balance (with eyes open),³⁴ and grip strength,³³ compared to controls. Furthermore, analysis of a measure as subjective as caregiver

perception revealed an increase of over 10% compared to baseline, in both qualitative and quantitative assessments.³³

However, results for hand function presented a similar trend to that observed in the control group.³³

One important consideration is the fact that the study by Gatica-Rojas et al.³⁴ was the only one to include assessment and follow-up in the medium term. In that study, the short-term benefits observed persisted in the medium term. In this sense, balance (eyes open) improved significantly more in the intervention group than in the control group. However, this trend was not observed in the other variables analysed, with both groups showing comparable results.

In patients with DCD, 4 outcome variables were analysed in the medium term. Specifically, neither isometric strength⁴³ nor motor performance⁴⁰ showed significant differences between the intervention and the control groups. However, functional strength⁴³ and motor proficiency⁴³ did show significant differences between groups, with better results in the conventional physiotherapy group. Specifically, motor proficiency⁴³ showed an increase in total test score, particularly conditioned by scores for 2 components (manual dexterity and balance).

Long-term effects

Eight outcome variables were analysed in the long term, in 2 studies (assessing 4 variables) that included patients with CP^{33,35} and one (4 variables) that included individuals with DCD.³⁹ Specifically, patients with CP showed poorer upper limb quality of movement in 2 of the variables analysed, whereas a third (grip strength) improved with respect to baseline.³⁵ This long-term improvement is consistent with that observed by Chiu et al.,³³ who also observed significant differences compared to baseline, although the magnitude of the effect was not significantly larger than that observed in the control group.³³ This similarity in the progression of outcome variables in the intervention and the control groups was also observed for hand function. Caregiver perception not only showed persistence of the improvement observed in the medium term, but even a slight increase.³³

Another 4 variables were analysed in a study of patients with DCD.³⁹ Three of these (muscle force, self-efficacy, and functional performance) behaved similarly in both the intervention and the control groups. Furthermore, analysis of running performance after the intervention showed better results in the control group than in the intervention group in the long term.³⁹

This review is not free of potential limitations due to the databases selected, filtering for language of publication, and the inclusion criteria applied in the literature search. The heterogeneity of the studies reviewed in terms of methodology and the outcome measures studied prevented us from conducting a meta-analysis.

In this sense, raising awareness of gamification among neurorehabilitation professionals in general, and physiotherapists in particular, will lead to new therapeutic opportunities. Thus, realistic use of gamification in physiotherapy may enable optimisation of clinical practice, and represents a promising approach.²¹

Future studies should use larger patient samples in order to obtain more robust results. It would also be interesting to establish the most beneficial dose of neurorehabilitation treatments incorporating gamification.

In conclusion, the literature consulted on gamification in paediatric neurology supports its use in clinical practice, either in isolation or as a supplement to conventional therapy; these techniques are safe to use in natural environments such as the home or school settings, and offer motivation for children and adolescents with neuromotor impairment.

Funding

This study has received no specific funding from any public, commercial, or non-profit organisation.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Vidal-Samsó J. La neurorrehabilitación, un proceso de alta complejidad. *Rev Neurol*. 2020;70:433, <http://dx.doi.org/10.33588/rn.7012.2019481>.
- Ketelaar M, Vermeer A, Hart TH, Van Petegem-van Beek E, Hadders PJM. Effects of a functional therapy program on motor abilities of children with cerebral palsy. *Phys Ther*. 2001;81:1534–45, <http://dx.doi.org/10.1093/ptj/81.9.1534>.
- Hadders PJM, Engelbert RHH, Custers JWH, Gorter JW, Takken T, Van Der Net J. Creating and being created: the changing panorama of paediatric rehabilitation. *Pediatr Rehabil*. 2003;6:5–12, <http://dx.doi.org/10.1080/1363849031000095260>.
- Ahl LE, Johansson E, Granat T, Carlberg EB. Functional therapy for children with cerebral palsy: an ecological approach. *Dev Med Child Neurol*. 2005;47:613–9, <http://dx.doi.org/10.1017/S0012162205001210>.
- Law M, Darrach J, Pollock N, Rosenbaum P, Russell D, Walter SD, et al. Focus on function — a randomized controlled trial comparing two rehabilitation interventions for young children with cerebral palsy. *BMC Pediatr*. 2007;7:1–12, <http://dx.doi.org/10.1186/1471-2431-7-31>.
- World Health Organization. International Classification of functioning, disability and health. Children & youth version. s. f.
- Tixes F. La generación Y. In: Tixes F, editor. *Gamificación: fundamentos y aplicaciones*. Barcelona: UOC; 2014. p. 34–60.
- Deterding S, Khaled R, Nacke LE, Dixon D. Gamification: toward a definition. *Stud Comput Intell*. 2010;300:1–361, http://dx.doi.org/10.1007/978-3-642-13959-8_1.
- Lopes S, Pereira A, Magalhães P, Oliveira A, Rosário P. Gamification: focus on the strategies being implemented in interventions: a systematic review protocol. *BMC Res Notes*. 2019;12:100, <http://dx.doi.org/10.1186/s13104-019-4139-x>.
- Sardi L, Idri A, Fernández-Alemán JL. A systematic review of gamification in e-Health. *J Biomed Inform*. 2017;71:31–48, <http://dx.doi.org/10.1016/j.jbi.2017.05.011>.
- Levac D, McCormick A, Levin MF, Brien M, Mills R, Miller E, et al. Active video gaming for children with cerebral palsy: does a clinic-based virtual reality component offer an additive benefit? A pilot study. *Phys Occup Ther Pediatr*. 2018;38:74–87, <http://dx.doi.org/10.1080/01942638.2017.1287810>.
- Cheng VWS, Davenport T, Johnson D, Vella K, Hickie IB. Gamification in apps and technologies for improving mental health and well-being: systematic review. *J Med Internet Res*. 2019;21:1–15, <http://dx.doi.org/10.2196/13717>.
- Yates M, Kelemen A, Sik Lanyi C. Virtual reality gaming in the rehabilitation of the upper extremities post-stroke. *Brain Inj*. 2016;30:855–63, <http://dx.doi.org/10.3109/02699052.2016.1144146>.
- Psychouli P, Katzis K, Elliott M. Home-based training support for stroke patients using the leap motion and StandInExercise stand. *Stud Health Technol Inform*. 2018;251:55–8, <http://dx.doi.org/10.3233/978-1-61499-880-8-55>.
- Tsekleves E, Paraskevopoulos IT, Warland A, Kilbride C. Development and preliminary evaluation of a novel low cost VR-based upper limb stroke rehabilitation platform using Wii technology. *Disabil Rehabil Assist Technol*. 2016;11:413–22, <http://dx.doi.org/10.3109/17483107.2014.981874>.
- Trombetta M, Bazzanello Henrique PP, Brum MR, Colussi EL, De Marchi ACB, Rieder R. Motion rehab AVE 3D: a VR-based exergame for post-stroke rehabilitation. *Comput Methods Programs Biomed*. 2017;151:15–20, <http://dx.doi.org/10.1016/j.cmpb.2017.08.008>.
- Ungerleider LG, Doyon J, Karni A. Imaging brain plasticity during motor skill learning. *Neurobiol Learn Mem*. 2002;78:553–64, <http://dx.doi.org/10.1006/nlme.2002.4091>.
- van der Kooij K, van Dijksseldonk R, van Veen M, Steenbrink F, de Weerd C, Overvliet KE. Gamification as a sustainable source of enjoyment during balance and gait exercises. *Front Psychol*. 2019;10:1–12, <http://dx.doi.org/10.3389/fpsyg.2019.00294>.
- Novak I, McIntyre S, Morgan C, Campbell L, Dark L, Morton N, et al. A systematic review of interventions for children with cerebral palsy: state of the evidence. *Dev Med Child Neurol*. 2013;55:885–910, <http://dx.doi.org/10.1111/dmcn.12246>.
- Purpura G, Tinelli F, Bargagna S, Bozza M, Bastiani L, Cioni G. Effect of early multisensory massage intervention on visual functions in infants with Down syndrome. *Early Hum Dev*. 2014;90:809–13, <http://dx.doi.org/10.1016/j.earlhumdev.2014.08.016>.
- Janssen J, Verschuren O, Renger WJ, Ermers J, Ketelaar M, Van Ee R. Gamification in physical therapy: more than using games. *Pediatr Phys Ther*. 2017;29:95–9, <http://dx.doi.org/10.1097/PEP.0000000000000326>.
- Lohse K, Shirzad N, Verster A, Hodges N, Van der Loos H. Video games and rehabilitation: using design principles to enhance engagement in physical therapy. *J Neurol Phys Ther*. 2013;37:166–75, <http://dx.doi.org/10.1097/NPT.0000000000000017>.
- Shah N, Basteris A, Amirabdollahian F. Design parameters in multimodal games for rehabilitation. *Games Health J*. 2014;3:13–20, <http://dx.doi.org/10.1089/g4h.2013.0044>.
- Bratton SC, Ray D, Rhine T, Jones L. The efficacy of play therapy with children: a meta-analytic review of treatment outcomes. *Prof Psychol Res Pract*. 2005;36:376–90, <http://dx.doi.org/10.1037/0735-7028.36.4.376>.
- Cugelman B. Gamification: what it is and why it matters to digital health behavior change developers. *J Med Internet Res*. 2013;15:1–6, <http://dx.doi.org/10.2196/games.3139>.
- Plasencia-Robledo M. Recursos tecnológicos en rehabilitación pediátrica. In: Macías L, Fagoaga J, editors. *Fisioterapia en Pediatría*. Madrid: Médica Panamericana; 2018. p. 479.
- Acar G, Altun GP, Yurdalan S, Polat MG. Efficacy of neurodevelopmental treatment combined with the nintendo® wii in patients with cerebral palsy. *J Phys Ther Sci*. 2016;28:774–80, <http://dx.doi.org/10.1589/jpts.28.774>.
- Ilg W, Schatton C, Schicks J, Giese MA, Schöls L, Synofzik M. Video game-based coordinative training improves ataxia in children with degenerative ataxia. *Neurology*. 2012;79:2056–60, <http://dx.doi.org/10.1212/WNL.0b013e3182749e67>.
- Facchin P, Rosa-Rizzotto M, Dalla Pozza LV, Turconi AC, Pagliano E, Signorini S, et al. Multisite trial comparing the efficacy of constraint-induced movement therapy with that of bimanual intensive training in children with hemiplegic cerebral palsy: postintervention results. *Am J Phys Med Rehabil*. 2011;90:539–53, <http://dx.doi.org/10.1097/PHM.0b013e3182247076>.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: expla-

- nation and elaboration. *J Clin Epidemiol.* 2009;62:1–34, <http://dx.doi.org/10.1016/j.jclinepi.2009.06.006>.
31. de Morton NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother.* 2009;55:129–33, [http://dx.doi.org/10.1016/S0004-9514\(09\)70043-1](http://dx.doi.org/10.1016/S0004-9514(09)70043-1).
 32. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther.* 2003;83:713–21, <http://dx.doi.org/10.1093/ptj/83.8.713>.
 33. Chiu HC, Ada L, Lee HM. Upper limb training using Wii Sports Resort™ for children with hemiplegic cerebral palsy: a randomized, single-blind trial. *Clin Rehabil.* 2014;28:1015–24, <http://dx.doi.org/10.1177/0269215514533709>.
 34. Gatica-Rojas V, Méndez-Rebolledo G, Guzman-Muñoz E, Soto-Poblete A, Cartes-Velásquez R. Does Nintendo Wii Balance Board improve standing balance? A randomized controlled trial in children with cerebral palsy. *Eur J Phys Rehabil Med.* 2017;53:535–45, <http://dx.doi.org/10.23736/S1973-9087.16.04447-6>.
 35. Sajan JE, John JA, Grace P, Sabu SS, Tharion G. Wii-based interactive video games as a supplement to conventional therapy for rehabilitation of children with cerebral palsy: a pilot, randomized controlled trial. *Dev Neurorehabil.* 2017;20:361–7, <http://dx.doi.org/10.1080/17518423.2016.1252970>.
 36. Kasee C, Hunt C, Holmes MWR, Lloyd M. Home-based Nintendo Wii training to improve upper-limb function in children ages 7 to 12 with spastic hemiplegic cerebral palsy. *J Pediatr Rehabil Med.* 2017;10:145–54, <http://dx.doi.org/10.3233/PRM-170439>.
 37. Hsieh HC. Effects of a gaming platform on balance training for children with cerebral palsy. *Pediatr Phys Ther.* 2018;30:303–8, <http://dx.doi.org/10.1097/PEP.0000000000000521>.
 38. Arnoni JLB, Pavão SL, dos Santos Silva FP, Rocha NACF. Effects of virtual reality in body oscillation and motor performance of children with cerebral palsy: a preliminary randomized controlled clinical trial. *Complement Ther Clin Pract.* 2019;35:189–94, <http://dx.doi.org/10.1016/j.ctcp.2019.02.014>.
 39. Bonney E, Ferguson G, Smits-Engelsman B. The efficacy of two activity-based interventions in adolescents with Developmental Coordination Disorder. *Res Dev Disabil.* 2017;71:223–36, <http://dx.doi.org/10.1016/j.ridd.2017.10.013>.
 40. Hammond J, Jones V, Hill EL, Green D, Male I. An investigation of the impact of regular use of the Wii Fit to improve motor and psychosocial outcomes in children with movement difficulties: a pilot study. *Child Care Health Dev.* 2014;40:165–75, <http://dx.doi.org/10.1111/cch.12029>.
 41. Jelsma D, Geuze RH, Mombarg R, Smits-Engelsman BCM. The impact of Wii Fit intervention on dynamic balance control in children with probable Developmental Coordination Disorder and balance problems. *Hum Mov Sci.* 2014;33:404–18, <http://dx.doi.org/10.1016/j.humov.2013.12.007>.
 42. Brüttsch K, Schuler T, Koenig A, Zimmerli L, Mérillat S, Lünenburger L, et al. Influence of virtual reality soccer game on walking performance in robotic assisted gait training for children. *J Neuroeng Rehabil.* 2010;7, <http://dx.doi.org/10.1186/1743-0003-7-15>.
 43. Ferguson GD, Jelsma D, Jelsma J, Smits-Engelsman BCM. The efficacy of two task-orientated interventions for children with Developmental Coordination Disorder: Neuromotor Task Training and Nintendo Wii Fit training. *Res Dev Disabil.* 2013;34:2449–61, <http://dx.doi.org/10.1016/j.ridd.2013.05.007>.