

COUP-TFII EXPRESSION DEFINES TWO DIFFERENT SEPTUM TRANSVERSUM CELL COMPARTMENTS CRUCIAL TO CARDIAC SEPTATION AND COMPACT VENTRICULAR WALL GROWTH

J.A.Guadix^{1,2}, A.A.S.Pires-Gomes^{1,2}, S.Cano^{1,2}, C.Pogontke^{1,2}, P.Palmquist^{1,2}, A.Rojas³, J.M.Pérez-Pomares^{1,2}

1. Department of Animal Biology, Faculty of Sciences, Instituto de Investigación Biomédica de Málaga (IBIMA), University of Málaga, Campus de Teatinos s/n, 29071 Málaga, Spain.
2. Andalusian Center for Nanomedicine and Biotechnology (BIONAND), 29590 Campanillas (Málaga), Spain.
3. Centro Andaluz de Biología Molecular y Medicina Regenerativa (CABIMER), Avda. Americo Vesputio s/n. centro de Investigación Biomédica en red de Diabetes y Enfermedades metabólicas Asociadas (CIBERDEM), Parque Científico Isla de la Cartuja, 41092 Sevilla, Spain.

COUP-TFII encodes for an orphan nuclear receptor expressed by multiple embryonic tissues. *COUP-TFII* functions include the regulation of mesodermal progenitor differentiation and cell fate specification, and is required for completion of cardiovascular development as shown by the early death (E9.5) of *COUP-TFII*-null mice. In this study, we show that *COUP-TFII*, which is strongly expressed in the atrial myocardium, is also expressed in two different compartments of the *septum transversum* (ST, E9.5), a mesodermal folding adjacent to cardiac inflow myocardium. The first ST compartment is *COUP-TFII*⁺/*Isl1*⁺; cells in this compartment concentrate in the posterior part of the ST, overlap with SHF, and are continuous with the dorsal mesenchymal protrusion (DMP, also known as *spina vestibuli*). The second compartment is characteristically *COUP-TFII*⁺/*Isl1*⁻, and comprises the majority of proepicardial cells. To dissect the role of *COUP-TFII*⁺ ST cells in cardiac development, we conditionally deleted *COUP-TFII* in the ST using two different Cre constructs (*Wt1Cre*; *G2-Gata4Cre*). We show that *COUP-TFII* deletion in the ST is most severe in *G2-Gata4Cre;COUP-TFII*^{-/-} mice, containing various cardiovascular progenitor lineages. Mutant mice display atrial septation and atrioventricular septal defects as well as a severe disruption of compact ventricular myocardial growth and coronary vascularization. We conclude that *COUP-TFII* plays critical, pleiotropic, tissue-dependent roles during cardiac septation, growth and vascularization.