

Poster preferred

Understanding Cross-Conjugation for Organic Electronics

 Samara Medina,^a Jean-François Morin,^b Xiaozhang Zhu^c, Juan Casado^a
^a Univ. Malaga, Dept. Phys. Chem., Campus Teatinos S-N, E-29071 Malaga, Spain, samara@uma.es.

^b Département de Chimie, Université Laval, 1045 Ave de la Médecine, Québec, G1V 0A6, Canada.


^c Chinese Acad. Sci., Inst. Chem., CAS Key Lab Organ Solids, Beijing 100080, Peoples R China.

Abstract:

π -Conjugated organic molecules have been the focus of interest since they have been probed as potential semiconducting materials,^[1] suitable for replacing the widely used silicon technologies. Their structural, optical and conductive properties are now under study to improve their application in organic electronics and to make possible their *ad hoc* synthesis. In this sense, the knowledge of the π -electron delocalization is crucial to establish the relation between the properties and the function, enabling the development of a synthesis guide based on the specific application.

The most acknowledged conjugated organic materials are those which present extended, linearly-conjugated π -systems.^[1,2] However, this is not the only way of π -electron delocalization: homoconjugation, cross-conjugation, curved-conjugation, etc. constitute different electronic designs to achieve new organic materials.

There is a relative high abundance in the organic world of cross-conjugated but limited comprehension.^[1,2,3] Thus, the understanding of how cross-conjugation works in π -electronic systems is of importance. Following this idea, in this project we show 4 different structures which present two perpendicular π -conjugated paths and how the cross-conjugated property is revealed. On the one hand, two molecules based on thieno[3,4-c]pyrrole-4,6-dione quaterthiophenes^[2,3] allow us to accomplish the subject from the aromatic/quinoidal outlook, and, on the other hand, two molecules with an anthanthrone core make possible the study from the perspective of the substituent groups.^[4]



Cross-conjugated paths in a quaterthiophene-based molecule.

References:

1. Gholami, M.; Tykwinskyi, R. R., *Chem. Rev.*, **2006**, *106*, 4997–5027.
2. Ponce Ortiz, R. *et al.*, *Adv. Funct. Mater.*, **2009**, *19*, 386–394.
3. Yuan, D. *et al.*, *Chem. Eur. J.*, **2018**, *24*, 13523–13534.
4. Klokkenburg, M. *et al.*, *Chem. Eur. J.*, **2013**, *9*, 3544–3554.
5. Desroches, M. *et al.*, *Angew. Chem. Int. Ed.*, **2017**, *56*, 16212–16217.