

Stress, depression and the hippocampus: modulatory effects of continuous LPA treatment

Moreno-Fernández, R¹., Bacq A²., Rosell-Valle C¹., Zanoletti², O., Cifuentes, M³., Pérez-Martín, M³., García-Fernández, M⁴., Estivill-Torrús, G⁵., Rodríguez de Fonseca, F⁶., Santín, LJ¹., Sandi, C²., Pedraza, C¹.

¹ Departamento de Psicobiología y Metodología de las CC, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga; Málaga 29071, Spain.

² Laboratory of Behavioral Genetics, Brain Mind Institute, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

³ Departamento de Biología Celular, Genética y Fisiología, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga; Málaga 29071, Spain.

⁴ Departamento de Fisiología y Medicina Deportiva, Instituto de Investigación Biomédica de Málaga (IBIMA), Universidad de Málaga; Málaga 29071, Spain.

⁵ Unidad Gestión Clínica de Neurociencias, Instituto de Investigación Biomédica de Málaga (IBIMA), Hospital Regional Universitario de Málaga, Málaga 29010, Spain.

⁶ Unidad de Gestión Clínica de Salud Mental, Instituto de Investigación Biomédica de Málaga (IBIMA), Hospital Regional Universitario de Málaga; Málaga 29010, Spain.

E-mail correspondence: mdpedraza@uma.es

The LPA₁, one of the six characterized G protein-coupled receptors (LPA₁₋₆) through which lysophosphatidic acid acts may be involved in promoting normal emotional behaviors. Evidence also imply a role for the LPA₁ receptor in mediating the consequences of stress on the hippocampus. However, to date, there is not available information regarding the mechanisms whereby the LPA₁ receptor mediates this adaptation. Changes in glutamate/GABA cycling could be one possible mechanism. To gain further insight into how LPA-LPA₁ may prevent the negative consequences of chronic stress, we assessed the effects of chronic ICV administration of LPA on depressive-like behaviours induced by a chronic restraint stress protocol. Then, gene expression for molecular markers for excitatory and inhibitory neurotransmission was determined. In addition, the hippocampal expression of mineralocorticoid receptor and glucocorticoid receptor genes and proteins were determined, as well as plasma corticosterone levels. Contrary to expectations, the continuous delivery of LPA in chronically stressed animals instead of inhibiting, potentiated some, though not all, negative effects of stress. Furthermore, this treatment induced as well altered the excitatory/inhibitory balance in the ventral hippocampus. In conclusion, the results of this study reinforce the assumption that LPA, mainly through the LPA₁ receptor, regulates hippocampal-dependent behaviour and functions.

Key words: lysophosphatidic acid, LPA, chronic stress, animal models of depression, hippocampus

Funding: This study was supported by Consejería de Economía, Innovación, Ciencia y Empleo, Junta de Andalucía (SEJ1863 to C.P.; CTS-643 to G.E.-T), Ministerio de Economía y Competitividad and the European Regional Development Fund (PSI2017-83408-P to C.P.), Consejería de Salud de la Junta de Andalucía (NICOLÁS MONARDE Program to G.E:T) Ministerio de Educación, Cultura y Deporte (FPU14/01610 to R.D.M.-F.; FPD12010 to C.R.-V. (Junta de Andalucía) and intramural funding from the EPFL to C.S. Congress Attendance C.P_ *I plan propio de investigación y transferencia Univesidad de Málaga*